

4-(Dodecyloxy)benzotrile

Huey Chong Kwong, Mohamad Zaki Ab. Rahman,
Mohamed Ibrahim Mohamed Tahir and Sidik Silong*

Department of Chemistry, Faculty of Science, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

Correspondence e-mail: sidik@science.upm.edu.my


Received 14 September 2011; accepted 10 October 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.046; wR factor = 0.127; data-to-parameter ratio = 16.9.

In the title compound, $\text{C}_{19}\text{H}_{29}\text{NO}$, the C—C and C—N bond distances of the benzotrile group are 1.445 (2) and 1.157 (2) Å, respectively. The aliphatic fragment adopts a bent zigzag arrangement which differs from the planar zigzag arrangement normally observed in n -alkanes or long-chain alkylbenzenes. In the crystal, inversion dimers linked by pairs of C—H \cdots O hydrogen bonds occur. A C—H \cdots N interaction also occurs. In the crystal, molecules are packed with the nitrile and aliphatic groups oriented in a head-to-tail fashion involving, forming a ripple-like motif along the a axis.

Related literature

For standard bond lengths, see Allen *et al.* (1987). For related structures, see: Merz (2002); Britton *et al.* (2004); Kwong *et al.* (2011); Boese *et al.* (1999). The title compound was synthesised by reacting hydroxybenzotrile with bromoalkane, see Rahman *et al.* (2009).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{29}\text{NO}$	$V = 1708.4$ (3) Å ³
$M_r = 287.45$	$Z = 4$
Monoclinic, $P2_1/n$	Cu $K\alpha$ radiation
$a = 5.7080$ (6) Å	$\mu = 0.52$ mm ⁻¹
$b = 7.3644$ (8) Å	$T = 100$ K
$c = 40.642$ (5) Å	$0.17 \times 0.14 \times 0.09$ mm
$\beta = 90^\circ$	

Data collection

Oxford Diffraction Gemini E diffractometer	9444 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2006)	3214 independent reflections
$T_{\min} = 0.930$, $T_{\max} = 0.955$	2575 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	190 parameters
$wR(F^2) = 0.127$	H-atom parameters constrained
$S = 0.98$	$\Delta\rho_{\text{max}} = 0.24$ e Å ⁻³
3202 reflections	$\Delta\rho_{\text{min}} = -0.25$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

D—H \cdots A	D—H	H \cdots A	D \cdots A	D—H \cdots A
C10—H102 \cdots N7 ⁱ	0.98	2.67	3.468 (2)	139
C3—H31 \cdots O1 ⁱⁱ	0.94	2.67	3.569 (5)	159

Symmetry codes: (i) $-x, -y + 2, -z + 1$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: *Gemini* (Oxford Diffraction, 2006); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2006); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *CRYSTALS* (Betteridge *et al.*, 2003); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *CRYSTALS*.

The authors would like to acknowledge the Ministry of Science, Technology and Innovation (MOSTI) Malaysia for funding (Research Grant Nos. 04-01-04-SF0144 and 05-02-10-0934RU).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2353).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Prpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Altomare, A., Casciaro, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.
- Boese, R., Weiss, H.-C. & Blaeser, D. (1999). *Angew. Chem. Int. Ed.* **38**, 988–992.
- Britton, D., Sowa, J. R. & Mann, K. R. (2004). *Acta Cryst.* **C60**, o418–o420.
- Kwong, H. C., Rahman, M. Z. A., Mohamed Tahir, M. I. & Silong, S. (2011). *Acta Cryst.* **E67**, o612.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Merz, K. (2002). *Acta Cryst.* **E58**, o450–o451.
- Oxford Diffraction (2006). *CrysAlis PRO*, *CrysAlis RED* and *Gemini*. Oxford Diffraction Ltd, Abingdon, England.
- Rahman, M. Z. A., Salisu, A. A., Silong, S., Luffor, M. R. & Ayub, M. B. A. (2009). *Asian J. Appl. Sci.* **2**, 177–183.

supplementary materials

Acta Cryst. (2011). E67, o3000 [doi:10.1107/S1600536811041602]

4-(Dodecyloxy)benzotrile

H. C. Kwong, M. Z. A. Rahman, M. I. Mohamed Tahir and S. Silong

Comment

The titled compound (I), 4-(dodecyloxy)benzotrile (Fig. 1) was synthesised by reacting hydroxybenzotrile with bromoalkane (Rahman *et al.*, 2009). Bond distance and angles of (I) are in normal range (Allen *et al.* 1987). Bond distance of the benzotrile group C5—C6 and C6—N7 are 1.445 (2) Å and 1.157 (2) Å, respectively and these bond lengths are comparable with those in *p*-decylbenzotrile of 1.446 (3) Å and 1.153 (3) Å, respectively (Britton *et al.*, 2004).

In this molecule, the plane formed by benzotrile ring and O1 was almost planar, the largest deviation from the least-squares plane is 0.0187 (12) Å at O1. The benzene ring and the alkane carbon skeleton (C9—C2—O1—C10) form the torsion angle of 1.62 (2)°. In this structure the alkane carbon skeleton has a bended zigzag arrangement; this arrangement is in agreement with previously reported alkoxy benzenes [4-hexyloxybenzamide, Kwong *et al.*, 2011] However, the mean C(H3)—C(H2) and C(H2)—C(H2) distances, and C(H3)—C(H2)—C and C(H2)—C(H2)—C angles, are in accordance of those determined for *n*-alkanes and long-chain alkylbenzene, 1.521 (1) Å and 112.8 (1)–113.5 (1)°, respectively. (Boese *et al.*, 1999; Merz, 2002; Britton *et al.*, 2004).

In the crystal packing, the centrosymmetric hydrogen bond C3—H31[⋯]O1 is formed generating a hydrogen bonded ring (Table 1 and Fig. 2). Packing of the titled compound shows a ripple-like motif (Fig. 2) with nitrile and aliphatic groups oriented head-to-tail. The stacking interaction between the aromatic rings with the separation distances of their centres of gravity Cg1[⋯]Cg1ⁱ (-x,1-y,1-z) of 3.573 (1) and Cg1[⋯]Cg1ⁱⁱ (-x,2-y,1-z) of 3.808 (1) Å and slippage of 1.395 and 1.865 Å, respectively, were observed.

Experimental

The titled compound (I) was synthesised by reacting hydroxybenzotrile with bromoalkane with conventional heating (Rahman *et al.*, 2009). Crystals of (I) were grown from hexane using a slow evaporation.

Refinement

The H atoms were all located in a difference map, but those attached to carbon atoms were repositioned geometrically. The H atoms were initially refined with soft restraints on the bond lengths and angles to regularize their geometry (C—H in the range 0.93–0.98, N—H in the range 0.86–0.90 Å) and $U_{\text{iso}}(\text{H})$ (in the range 1.2–1.5 times U_{eq} of the parent atom), after which the positions were refined with riding constraints.

Figures



Fig. 1. Molecular structure of (I) with atom numbering and displacement ellipsoids at 50% probability level.



Fig. 2. The packing diagram of (I) showing a ripple-like motif viewing along a axis; hydrogen bonds were shown as dashed lines [b axis green; c axis blue].

4-(Dodecyloxy)benzotrile

Crystal data

$C_{19}H_{29}NO$	$F(000) = 632$
$M_r = 287.45$	$D_x = 1.117 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Cu $K\alpha$ radiation, $\lambda = 1.54180 \text{ \AA}$
$a = 5.7080 (6) \text{ \AA}$	Cell parameters from 3853 reflections
$b = 7.3644 (8) \text{ \AA}$	$\theta = 3\text{--}71^\circ$
$c = 40.642 (5) \text{ \AA}$	$\mu = 0.52 \text{ mm}^{-1}$
$\beta = 90^\circ$	$T = 100 \text{ K}$
$V = 1708.4 (3) \text{ \AA}^3$	Plate-like, colourless
$Z = 4$	$0.17 \times 0.14 \times 0.09 \text{ mm}$

Data collection

Oxford Diffraction Gemini E diffractometer	3214 independent reflections
Radiation source: sealed x-ray tube graphite	2575 reflections with $I > 2\sigma(I)$
$\omega/2\theta$ scans	$R_{\text{int}} = 0.020$
Absorption correction: multi-scan (<i>Crys.Alis PRO</i> ; Oxford Diffraction, 2006)	$\theta_{\text{max}} = 71.6^\circ$, $\theta_{\text{min}} = 4.4^\circ$
$T_{\text{min}} = 0.930$, $T_{\text{max}} = 0.955$	$h = -6 \rightarrow 6$
9444 measured reflections	$k = -8 \rightarrow 9$
	$l = -37 \rightarrow 49$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Hydrogen site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.046$	H-atom parameters constrained
$wR(F^2) = 0.127$	Method = Modified Sheldrick $w = 1/[\sigma^2(F^2) + (0.05P)^2 + 1.35P]$, where $P = [\max(F_o^2, 0) + 2F_c^2]/3$
$S = 0.98$	$(\Delta/\sigma)_{\text{max}} = 0.007$
3202 reflections	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
190 parameters	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$
0 restraints	

Special details

Refinement. Refinement. For this compound, 9444 numbers of reflections were collected and measured during the refinement. Symmetry related reflections were measured more than once and after merging the symmetry equivalent reflections there were only 3214

reflection left. 12 more reflections were filtered, as σ cutoff was set as 3 and (\sin^2/λ) set to >0.01 (to eliminate reflection measured near the vicinity of beam stop) therefore numbers of reflection reduced to 3202.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.33735 (19)	0.61108 (16)	0.45325 (3)	0.0243
C2	0.1734 (3)	0.6793 (2)	0.47454 (4)	0.0214
C3	0.2277 (3)	0.6631 (2)	0.50786 (4)	0.0221
C4	0.0719 (3)	0.7247 (2)	0.53150 (4)	0.0228
C5	-0.1403 (3)	0.8033 (2)	0.52206 (4)	0.0219
C6	-0.3025 (3)	0.8685 (2)	0.54668 (4)	0.0234
N7	-0.4333 (3)	0.9209 (2)	0.56623 (4)	0.0295
C8	-0.1922 (3)	0.8209 (2)	0.48877 (4)	0.0221
C9	-0.0351 (3)	0.7606 (2)	0.46502 (4)	0.0220
C10	0.2944 (3)	0.6284 (2)	0.41859 (4)	0.0242
C11	0.5019 (3)	0.5506 (2)	0.40024 (4)	0.0252
C12	0.4902 (3)	0.5918 (2)	0.36351 (4)	0.0246
C13	0.6978 (3)	0.5162 (2)	0.34403 (4)	0.0257
C14	0.7033 (3)	0.5790 (2)	0.30831 (4)	0.0251
C15	0.9124 (3)	0.5066 (2)	0.28868 (4)	0.0258
C16	0.9230 (3)	0.5776 (2)	0.25349 (4)	0.0257
C17	1.1323 (3)	0.5068 (2)	0.23378 (4)	0.0257
C18	1.1450 (3)	0.5806 (2)	0.19882 (4)	0.0258
C19	1.3543 (3)	0.5106 (2)	0.17902 (4)	0.0259
C20	1.3684 (3)	0.5847 (3)	0.14407 (4)	0.0282
C21	1.5784 (3)	0.5133 (3)	0.12487 (4)	0.0322
H31	0.3704	0.6093	0.5144	0.0260*
H41	0.1098	0.7150	0.5546	0.0259*
H81	-0.3353	0.8725	0.4821	0.0251*
H91	-0.0710	0.7743	0.4422	0.0250*
H102	0.2751	0.7577	0.4131	0.0287*
H101	0.1498	0.5632	0.4128	0.0284*
H111	0.6426	0.6050	0.4091	0.0296*
H112	0.5060	0.4186	0.4040	0.0302*
H122	0.4875	0.7226	0.3605	0.0291*
H121	0.3473	0.5416	0.3544	0.0290*
H131	0.8392	0.5561	0.3546	0.0305*
H132	0.6919	0.3834	0.3445	0.0313*
H142	0.7085	0.7111	0.3080	0.0303*
H141	0.5615	0.5402	0.2974	0.0293*
H151	1.0537	0.5428	0.3000	0.0304*
H152	0.9037	0.3730	0.2881	0.0309*
H162	0.9305	0.7093	0.2541	0.0313*
H161	0.7809	0.5426	0.2420	0.0302*
H172	1.2745	0.5408	0.2450	0.0312*
H171	1.1235	0.3737	0.2328	0.0314*
H181	1.1543	0.7134	0.1998	0.0308*
H182	1.0027	0.5487	0.1872	0.0310*

supplementary materials

H192	1.4972	0.5440	0.1907	0.0307*
H191	1.3442	0.3777	0.1780	0.0315*
H201	1.3788	0.7175	0.1451	0.0338*
H202	1.2254	0.5535	0.1326	0.0335*
H212	1.5813	0.5587	0.1021	0.0465*
H211	1.7246	0.5486	0.1352	0.0467*
H213	1.5746	0.3804	0.1239	0.0474*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0228 (6)	0.0292 (7)	0.0208 (6)	0.0029 (5)	-0.0019 (5)	-0.0005 (5)
C2	0.0216 (8)	0.0179 (8)	0.0247 (8)	-0.0037 (7)	-0.0010 (6)	-0.0006 (7)
C3	0.0195 (8)	0.0198 (8)	0.0269 (9)	-0.0008 (7)	-0.0045 (6)	0.0010 (7)
C4	0.0238 (9)	0.0212 (9)	0.0235 (8)	-0.0035 (7)	-0.0044 (7)	0.0001 (7)
C5	0.0208 (8)	0.0199 (8)	0.0251 (8)	-0.0043 (7)	0.0002 (6)	0.0002 (7)
C6	0.0224 (9)	0.0215 (8)	0.0261 (9)	-0.0022 (7)	-0.0055 (7)	0.0019 (7)
N7	0.0269 (8)	0.0338 (9)	0.0279 (8)	0.0001 (7)	-0.0017 (6)	-0.0015 (7)
C8	0.0173 (8)	0.0207 (9)	0.0283 (9)	-0.0022 (6)	-0.0044 (6)	0.0016 (7)
C9	0.0223 (8)	0.0211 (9)	0.0224 (8)	-0.0037 (7)	-0.0039 (6)	0.0016 (6)
C10	0.0231 (9)	0.0274 (9)	0.0221 (8)	-0.0011 (7)	-0.0031 (7)	0.0000 (7)
C11	0.0227 (9)	0.0283 (9)	0.0247 (9)	0.0004 (7)	-0.0018 (7)	-0.0009 (7)
C12	0.0226 (8)	0.0264 (9)	0.0249 (9)	-0.0001 (7)	-0.0024 (7)	0.0007 (7)
C13	0.0257 (9)	0.0274 (9)	0.0240 (9)	0.0034 (7)	-0.0025 (7)	-0.0012 (7)
C14	0.0228 (9)	0.0282 (9)	0.0244 (9)	0.0010 (7)	-0.0026 (7)	-0.0001 (7)
C15	0.0249 (9)	0.0277 (9)	0.0249 (9)	0.0026 (7)	-0.0036 (7)	-0.0016 (7)
C16	0.0251 (9)	0.0277 (9)	0.0242 (9)	0.0011 (7)	-0.0032 (7)	0.0001 (7)
C17	0.0254 (9)	0.0283 (9)	0.0234 (9)	0.0010 (7)	-0.0041 (7)	-0.0013 (7)
C18	0.0239 (9)	0.0284 (9)	0.0251 (9)	0.0009 (7)	-0.0034 (7)	0.0008 (7)
C19	0.0255 (9)	0.0281 (9)	0.0240 (9)	0.0003 (7)	-0.0040 (7)	-0.0016 (7)
C20	0.0270 (9)	0.0313 (10)	0.0264 (9)	0.0000 (8)	-0.0026 (7)	0.0006 (7)
C21	0.0308 (10)	0.0387 (11)	0.0269 (9)	-0.0008 (8)	-0.0012 (8)	-0.0007 (8)

Geometric parameters (\AA , $^\circ$)

O1—C2	1.3698 (19)	C13—H132	0.979
O1—C10	1.4355 (19)	C14—C15	1.531 (2)
C2—C3	1.394 (2)	C14—H142	0.973
C2—C9	1.387 (2)	C14—H141	0.966
C3—C4	1.385 (2)	C15—C16	1.524 (2)
C3—H31	0.943	C15—H151	0.967
C4—C5	1.396 (2)	C15—H152	0.985
C4—H41	0.965	C16—C17	1.530 (2)
C5—C6	1.445 (2)	C16—H162	0.972
C5—C8	1.391 (2)	C16—H161	0.970
C6—N7	1.157 (2)	C17—C18	1.523 (2)
C8—C9	1.390 (2)	C17—H172	0.963
C8—H81	0.941	C17—H171	0.982
C9—H91	0.954	C18—C19	1.530 (2)

C10—C11	1.512 (2)	C18—H181	0.980
C10—H102	0.984	C18—H182	0.968
C10—H101	0.984	C19—C20	1.524 (2)
C11—C12	1.525 (2)	C19—H192	0.976
C11—H111	0.967	C19—H191	0.981
C11—H112	0.984	C20—C21	1.524 (2)
C12—C13	1.530 (2)	C20—H201	0.981
C12—H122	0.971	C20—H202	0.967
C12—H121	0.969	C21—H212	0.985
C13—C14	1.524 (2)	C21—H211	0.969
C13—H131	0.961	C21—H213	0.980
C2—O1—C10	118.08 (12)	C15—C14—H142	108.6
O1—C2—C3	115.50 (14)	C13—C14—H141	109.3
O1—C2—C9	124.62 (14)	C15—C14—H141	108.1
C3—C2—C9	119.89 (15)	H142—C14—H141	108.4
C2—C3—C4	120.20 (15)	C14—C15—C16	113.61 (14)
C2—C3—H31	120.0	C14—C15—H151	107.8
C4—C3—H31	119.8	C16—C15—H151	108.7
C3—C4—C5	120.14 (15)	C14—C15—H152	108.7
C3—C4—H41	120.4	C16—C15—H152	108.8
C5—C4—H41	119.4	H151—C15—H152	109.2
C4—C5—C6	120.23 (15)	C15—C16—C17	113.93 (14)
C4—C5—C8	119.37 (15)	C15—C16—H162	108.6
C6—C5—C8	120.40 (15)	C17—C16—H162	108.6
C5—C6—N7	179.56 (17)	C15—C16—H161	109.0
C5—C8—C9	120.54 (15)	C17—C16—H161	108.1
C5—C8—H81	120.1	H162—C16—H161	108.3
C9—C8—H81	119.4	C16—C17—C18	113.81 (14)
C8—C9—C2	119.85 (15)	C16—C17—H172	108.8
C8—C9—H91	120.1	C18—C17—H172	107.9
C2—C9—H91	120.0	C16—C17—H171	108.7
O1—C10—C11	108.46 (13)	C18—C17—H171	108.8
O1—C10—H102	109.2	H172—C17—H171	108.7
C11—C10—H102	110.0	C17—C18—C19	114.02 (14)
O1—C10—H101	109.5	C17—C18—H181	108.6
C11—C10—H101	110.8	C19—C18—H181	108.4
H102—C10—H101	108.9	C17—C18—H182	109.1
C10—C11—C12	111.90 (14)	C19—C18—H182	108.5
C10—C11—H111	108.0	H181—C18—H182	108.0
C12—C11—H111	108.6	C18—C19—C20	114.25 (14)
C10—C11—H112	108.4	C18—C19—H192	108.1
C12—C11—H112	110.4	C20—C19—H192	108.7
H111—C11—H112	109.4	C18—C19—H191	108.2
C11—C12—C13	113.60 (14)	C20—C19—H191	108.8
C11—C12—H122	108.7	H192—C19—H191	108.7
C13—C12—H122	108.0	C19—C20—C21	113.30 (15)
C11—C12—H121	109.5	C19—C20—H201	108.7
C13—C12—H121	108.5	C21—C20—H201	108.5
H122—C12—H121	108.5	C19—C20—H202	108.7

supplementary materials

C12—C13—C14	113.50 (14)	C21—C20—H202	109.5
C12—C13—H131	107.9	H201—C20—H202	108.0
C14—C13—H131	108.5	C20—C21—H212	112.2
C12—C13—H132	109.1	C20—C21—H211	111.4
C14—C13—H132	108.8	H212—C21—H211	107.5
H131—C13—H132	109.0	C20—C21—H213	110.4
C13—C14—C15	114.01 (14)	H212—C21—H213	107.5
C13—C14—H142	108.3	H211—C21—H213	107.7

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C10—H102 \cdots N7 ⁱ	0.98	2.67	3.468 (2)	139.
C3—H31 \cdots O1 ⁱⁱ	0.94	2.67	3.569 (5)	159.

Symmetry codes: (i) $-x, -y+2, -z+1$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

