

4-[4-(1*H*-Tetrazol-5-yl)phenoxy]-benzaldehyde

Jing Lu, Jiao Xu, Li-Wei Ni, Wei-Li Ma and Zhen-Ting Du*

College of Science, Northwest A&F University, Yangling Shaanxi 712100, People's Republic of China

Correspondence e-mail: duzt@nwsuaf.edu.cn


Received 19 October 2011; accepted 24 October 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.056; wR factor = 0.142; data-to-parameter ratio = 12.7.

The asymmetric unit of the title compound, $\text{C}_{14}\text{H}_{10}\text{N}_4\text{O}_2$, contains two independent molecules with similar structures. In one molecule, the tetrazole ring is oriented at dihedral angles of $17.71(16)$ and $57.13(17)^\circ$, respectively, to the central benzene ring and the terminal benzene ring; in the other molecule, the corresponding dihedral angles are $16.46(18)$ and $75.87(18)^\circ$. Intermolecular $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds and weak $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds occur in the crystal structure.

Related literature

For the synthesis of 5-substituted 1*H*-tetrazoles, see: Ostrovskii *et al.* (2008); Saikia & Phukan (2009); Nasrollahzadeh *et al.* (2009); Teimouri & Najafi Chermahini (2011). For related structures, see: Li *et al.* (2008); Hu *et al.* (2009); Xu *et al.* (2010); Deng *et al.* (2010).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{10}\text{N}_4\text{O}_2$	$\gamma = 99.737(4)^\circ$
$M_r = 266.26$	$V = 1279.8(9)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 4$
$a = 9.854(4)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.948(4)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$c = 14.139(6)\text{ \AA}$	$T = 296\text{ K}$
$\alpha = 98.537(4)^\circ$	$0.23 \times 0.21 \times 0.19\text{ mm}$
$\beta = 106.668(4)^\circ$	

Data collection

Bruker APEXII CCD diffractometer
9274 measured reflections

4683 independent reflections
2623 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.142$
 $S = 1.01$
4683 reflections
369 parameters
30 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.41\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.30\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N \cdots N8 ⁱ	0.91 (3)	1.94 (3)	2.849 (4)	174 (2)
N5—H2N \cdots N4 ⁱⁱ	0.90 (3)	2.03 (3)	2.924 (3)	176 (3)
C4—H4 \cdots O4	0.93	2.54	3.420 (4)	158
C17—H17 \cdots N3 ⁱⁱ	0.93	2.58	3.365 (4)	143
C23—H23 \cdots O2 ⁱⁱⁱ	0.93	2.42	3.262 (6)	151

Symmetry codes: (i) $x + 1, y - 1, z + 1$; (ii) $x - 1, y, z - 1$; (iii) $-x, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

Financial support from the Fundamental Research Funds for the Central Universities in NWSUAF (No. QN2009048) and the opening project of Xinjiang Production & Construction Corps Key Laboratory of Protection and Utilization of Biological Resources in Tarim Basin (BRTD1004) as well as the National Natural Science Foundation of China (20802058) are greatly appreciated.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5357).

References

- Bruker (2001). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2005). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
Deng, J.-H., Yuan, X.-L. & Mei, G.-Q. (2010). *Inorg. Chem. Commun.* **13**, 1585–1589.
Hu, D.-Y., Chu, X.-W. & Qu, Z.-R. (2009). *Acta Cryst. E65*, o2463.
Li, G.-Q., Wu, A.-Q., Li, Y., Zheng, F.-K. & Guo, G.-C. (2008). *Acta Cryst. E64*, o1368.
Nasrollahzadeh, M., Bayat, Y., Habibi, D. & Moshaei, S. (2009). *Tetrahedron Lett.* **50**, 4435–4438.
Ostrovskii, V. A., Koldobskii, G. I., Trifonov, R. E., Alan, R. K., Christopher, A. R., Eric, F. V. S. & Richard, J. K. T. (2008). *Comprehensive Heterocyclic Chemistry III*, p. 257. Oxford: Elsevier.
Saikia, I. & Phukan, P. (2009). *Tetrahedron Lett.* **50**, 5083–5087.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Teimouri, A. & Najafi Chermahini, A. (2011). *Polyhedron*, **30**, 2606–2610.
Xu, L., Gu, L.-Y., Zhao, D.-Y., Wang, B. & Kang, T.-G. (2010). *Acta Cryst. E66*, o742.

supplementary materials

Acta Cryst. (2011). E67, o3106 [doi:10.1107/S1600536811044254]

4-[4-(1*H*-Tetrazol-5-yl)phenoxy]benzaldehyde

J. Lu, J. Xu, L.-W. Ni, W.-L. Ma and Z.-T. Du

Comment

Tetrazoles play a variety of roles in coordination chemistry, medicinal chemistry, materials chemistry etc. (Ostrovskii *et al.*, 2008). We intended to find a new way for preparation of tetrazoles using a new solid acid which is regarded as a green catalyst. In order to confirm the verity of the final product, the single crystal X-ray analysis was performed and the structure is reported here.

Compared with the other phenyltetrazole coplanar structures reported by Li *et al.* (2008) and Xu *et al.* (2010), there is torsion between tetrazole rings and neighboring phenyl rings, with a dihedral angle 17.7 (1) $^{\circ}$ and 16.4 (4) $^{\circ}$, respectively.

Experimental

A solution of 4-(4-formylphenoxy)benzonitrile (892 mg, 4 mmol) and sodium azide (780 mg, 12 mmol) in dry DMF (15 mL) at the presence of sulfuric acid on silica gel (5% load, 400 mg) was heated to 80°C. When the reaction was completed, the solid acid was filtered, 2 mL water was added to the filtration, and then was extracted with ethyl acetate (20 mL \times 3). The ethyl acetate layers were combined and washed by 20 mL water, and then 15 mL saturated sodium chloride and dried over anhydrous sodium sulfate. The solution was evaporated and the residue was separated on silica gel column chromatography with a gradient of petroleum ether and ethyl acetate as eluent to yield 570 mg the title compound. The compound was then dissolved in methanol, and colorless crystals were formed on slow evaporation at room temperature over one week.

Refinement

The H1N and H2N atoms were located in a difference Fourier map and refined isotropically. Other H atoms were placed in calculated positions with C—H = 0.93 Å and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

4-[4-(1*H*-Tetrazol-5-yl)phenoxy]benzaldehyde

Crystal data

C₁₄H₁₀N₄O₂

$M_r = 266.26$

$Z = 4$

$F(000) = 552$

supplementary materials

Triclinic, $P\bar{1}$	$D_x = 1.382 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.854 (4) \text{ \AA}$	Cell parameters from 1475 reflections
$b = 9.948 (4) \text{ \AA}$	$\theta = 2.2\text{--}21.0^\circ$
$c = 14.139 (6) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$\alpha = 98.537 (4)^\circ$	$T = 296 \text{ K}$
$\beta = 106.668 (4)^\circ$	Block, colourless
$\gamma = 99.737 (4)^\circ$	$0.23 \times 0.21 \times 0.19 \text{ mm}$
$V = 1279.8 (9) \text{ \AA}^3$	

Data collection

Bruker APEXII CCD diffractometer	2623 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.035$
graphite	$\theta_{\text{max}} = 25.5^\circ, \theta_{\text{min}} = 2.2^\circ$
φ and ω scans	$h = -11\text{--}11$
9274 measured reflections	$k = -10\text{--}12$
4683 independent reflections	$l = -17\text{--}17$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.056$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.142$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.01$	$w = 1/[(\sigma^2(F_o^2) + (0.0462P)^2 + 0.4163P)]$
4683 reflections	where $P = (F_o^2 + 2F_c^2)/3$
369 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
30 restraints	$\Delta\rho_{\text{max}} = 0.41 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.8863 (3)	0.1795 (3)	1.0915 (2)	0.0414 (7)
C2	0.7457 (3)	0.1392 (3)	1.0126 (2)	0.0415 (7)
C3	0.6554 (3)	0.2318 (3)	1.0004 (2)	0.0514 (8)
H3	0.6862	0.3205	1.0416	0.062*
C4	0.5208 (3)	0.1945 (3)	0.9283 (2)	0.0602 (8)
H4	0.4598	0.2568	0.9209	0.072*
C5	0.4773 (3)	0.0638 (3)	0.8672 (2)	0.0543 (8)
C6	0.5645 (3)	-0.0298 (3)	0.8772 (2)	0.0597 (8)
H6	0.5335	-0.1177	0.8349	0.072*
C7	0.6986 (3)	0.0072 (3)	0.9504 (2)	0.0533 (8)
H7	0.7580	-0.0564	0.9583	0.064*
C8	0.3025 (3)	0.0912 (3)	0.7196 (2)	0.0558 (8)
C9	0.1567 (3)	0.0772 (4)	0.6724 (3)	0.0728 (10)
H9	0.0880	0.0209	0.6914	0.087*
C10	0.1142 (4)	0.1482 (4)	0.5964 (3)	0.0820 (11)
H10	0.0155	0.1401	0.5647	0.098*
C11	0.2142 (4)	0.2309 (4)	0.5662 (3)	0.0718 (9)
C12	0.3588 (4)	0.2389 (3)	0.6127 (2)	0.0668 (9)
H12	0.4274	0.2921	0.5918	0.080*
C13	0.4051 (3)	0.1701 (3)	0.6898 (2)	0.0629 (9)
H13	0.5037	0.1769	0.7208	0.075*
C14	0.1805 (5)	0.3200 (5)	0.4876 (3)	0.1064 (14)
H14	0.2555	0.3725	0.4720	0.128*
C15	-0.0013 (3)	0.7097 (3)	0.1508 (2)	0.0430 (7)
C16	0.1047 (3)	0.7403 (3)	0.2513 (2)	0.0455 (7)
C17	0.1222 (4)	0.6369 (3)	0.3065 (2)	0.0667 (9)
H17	0.0665	0.5468	0.2782	0.080*
C18	0.2201 (4)	0.6649 (3)	0.4020 (2)	0.0747 (11)
H18	0.2318	0.5936	0.4373	0.090*
C19	0.3005 (3)	0.7972 (3)	0.4453 (2)	0.0587 (8)
C20	0.2850 (3)	0.9026 (3)	0.3926 (2)	0.0606 (9)
H20	0.3397	0.9927	0.4219	0.073*
C21	0.1884 (3)	0.8738 (3)	0.2964 (2)	0.0530 (8)
H21	0.1788	0.9451	0.2609	0.064*
C22	0.4087 (4)	0.7458 (3)	0.6065 (2)	0.0612 (9)
C23	0.2899 (4)	0.7028 (3)	0.6363 (3)	0.0678 (9)
H23	0.2027	0.7280	0.6082	0.081*
C24	0.3016 (3)	0.6222 (3)	0.7079 (3)	0.0627 (9)
H24	0.2217	0.5927	0.7282	0.075*
C25	0.4307 (3)	0.5845 (3)	0.7501 (2)	0.0551 (8)
C26	0.5488 (4)	0.6304 (3)	0.7202 (3)	0.0644 (9)
H26	0.6365	0.6065	0.7489	0.077*
C27	0.5387 (4)	0.7110 (3)	0.6487 (3)	0.0663 (9)
H27	0.6189	0.7416	0.6290	0.080*
C28	0.4455 (4)	0.4991 (3)	0.8268 (3)	0.0734 (10)

supplementary materials

H28	0.5360	0.4795	0.8538	0.088*
N1	0.9674 (3)	0.0935 (3)	1.12958 (18)	0.0499 (6)
N2	1.0895 (3)	0.1684 (2)	1.20221 (19)	0.0581 (7)
N3	1.0818 (3)	0.2977 (2)	1.20764 (19)	0.0584 (7)
N4	0.9565 (2)	0.3084 (2)	1.13952 (18)	0.0503 (6)
N5	-0.0625 (3)	0.5829 (2)	0.09329 (18)	0.0466 (6)
N6	-0.1563 (3)	0.5921 (2)	0.00568 (18)	0.0545 (6)
N7	-0.1514 (3)	0.7233 (2)	0.00983 (18)	0.0565 (7)
N8	-0.0556 (3)	0.7999 (2)	0.09919 (18)	0.0520 (6)
O1	0.3385 (2)	0.0200 (2)	0.79639 (17)	0.0707 (7)
O2	0.0601 (4)	0.3220 (4)	0.4469 (3)	0.1609 (14)
O3	0.4057 (2)	0.8339 (2)	0.53920 (17)	0.0789 (7)
O4	0.3497 (3)	0.4526 (3)	0.8575 (2)	0.0967 (9)
H2N	-0.053 (3)	0.499 (3)	0.107 (2)	0.058 (9)*
H1N	0.954 (3)	-0.001 (3)	1.117 (2)	0.081 (11)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0474 (17)	0.0281 (14)	0.0483 (17)	0.0103 (13)	0.0123 (14)	0.0106 (13)
C2	0.0441 (16)	0.0303 (14)	0.0494 (17)	0.0101 (12)	0.0107 (14)	0.0121 (12)
C3	0.0529 (19)	0.0350 (15)	0.0599 (19)	0.0138 (14)	0.0082 (16)	0.0058 (14)
C4	0.0524 (19)	0.0473 (18)	0.076 (2)	0.0198 (16)	0.0077 (17)	0.0148 (17)
C5	0.0447 (18)	0.0490 (18)	0.0581 (19)	0.0046 (15)	0.0008 (15)	0.0153 (15)
C6	0.065 (2)	0.0373 (16)	0.061 (2)	0.0086 (15)	0.0021 (17)	0.0003 (14)
C7	0.0558 (19)	0.0388 (16)	0.0597 (19)	0.0161 (14)	0.0080 (16)	0.0082 (14)
C8	0.053 (2)	0.0499 (18)	0.0533 (19)	0.0130 (16)	0.0014 (16)	0.0058 (15)
C9	0.046 (2)	0.094 (3)	0.073 (2)	0.0132 (19)	0.0095 (18)	0.024 (2)
C10	0.052 (2)	0.119 (3)	0.070 (2)	0.031 (2)	0.004 (2)	0.022 (2)
C11	0.069 (2)	0.093 (3)	0.060 (2)	0.041 (2)	0.0175 (19)	0.0136 (18)
C12	0.071 (2)	0.069 (2)	0.061 (2)	0.0207 (19)	0.0215 (19)	0.0127 (18)
C13	0.0494 (19)	0.068 (2)	0.063 (2)	0.0149 (17)	0.0072 (17)	0.0101 (18)
C14	0.090 (3)	0.133 (3)	0.086 (3)	0.070 (2)	-0.007 (2)	0.011 (2)
C15	0.0487 (17)	0.0287 (14)	0.0513 (17)	0.0090 (13)	0.0157 (14)	0.0082 (13)
C16	0.0487 (17)	0.0318 (15)	0.0509 (17)	0.0096 (13)	0.0088 (14)	0.0062 (13)
C17	0.083 (2)	0.0308 (16)	0.062 (2)	-0.0002 (16)	-0.0051 (18)	0.0075 (15)
C18	0.096 (3)	0.0411 (18)	0.062 (2)	0.0041 (18)	-0.009 (2)	0.0154 (16)
C19	0.058 (2)	0.0500 (19)	0.0533 (19)	0.0019 (16)	0.0007 (16)	0.0103 (16)
C20	0.0534 (19)	0.0372 (17)	0.073 (2)	-0.0035 (14)	0.0022 (17)	0.0066 (16)
C21	0.0505 (18)	0.0336 (15)	0.067 (2)	0.0050 (14)	0.0064 (16)	0.0148 (14)
C22	0.059 (2)	0.0532 (19)	0.0505 (19)	-0.0055 (17)	-0.0011 (17)	0.0063 (16)
C23	0.053 (2)	0.065 (2)	0.071 (2)	0.0110 (17)	0.0019 (18)	0.0117 (19)
C24	0.053 (2)	0.061 (2)	0.071 (2)	0.0094 (17)	0.0178 (17)	0.0098 (18)
C25	0.058 (2)	0.0457 (18)	0.0545 (19)	0.0123 (16)	0.0111 (17)	0.0031 (15)
C26	0.053 (2)	0.062 (2)	0.070 (2)	0.0134 (17)	0.0078 (18)	0.0096 (18)
C27	0.051 (2)	0.072 (2)	0.068 (2)	0.0037 (17)	0.0157 (18)	0.0101 (19)
C28	0.085 (3)	0.060 (2)	0.080 (3)	0.026 (2)	0.030 (2)	0.011 (2)
N1	0.0518 (16)	0.0326 (14)	0.0574 (16)	0.0109 (12)	0.0044 (13)	0.0101 (12)

N2	0.0553 (16)	0.0395 (14)	0.0648 (17)	0.0116 (12)	-0.0032 (13)	0.0093 (12)
N3	0.0588 (17)	0.0371 (14)	0.0655 (17)	0.0109 (12)	-0.0001 (14)	0.0086 (12)
N4	0.0510 (15)	0.0313 (13)	0.0593 (15)	0.0086 (11)	0.0048 (13)	0.0077 (11)
N5	0.0557 (15)	0.0295 (13)	0.0505 (15)	0.0124 (12)	0.0078 (12)	0.0106 (11)
N6	0.0652 (17)	0.0411 (14)	0.0505 (16)	0.0131 (12)	0.0072 (13)	0.0103 (12)
N7	0.0693 (17)	0.0396 (14)	0.0541 (16)	0.0144 (13)	0.0083 (14)	0.0107 (12)
N8	0.0654 (16)	0.0334 (13)	0.0548 (15)	0.0156 (12)	0.0108 (13)	0.0132 (12)
O1	0.0513 (13)	0.0620 (14)	0.0781 (16)	-0.0014 (11)	-0.0063 (12)	0.0215 (12)
O2	0.142 (3)	0.222 (4)	0.138 (3)	0.066 (3)	0.038 (3)	0.081 (3)
O3	0.0761 (16)	0.0672 (15)	0.0615 (15)	-0.0176 (12)	-0.0099 (13)	0.0190 (12)
O4	0.129 (2)	0.0865 (19)	0.111 (2)	0.0483 (18)	0.071 (2)	0.0389 (17)

Geometric parameters (\AA , $^\circ$)

C1—N4	1.325 (3)	C16—C21	1.388 (4)
C1—N1	1.335 (3)	C17—C18	1.373 (4)
C1—C2	1.455 (4)	C17—H17	0.9300
C2—C3	1.381 (3)	C18—C19	1.366 (4)
C2—C7	1.389 (4)	C18—H18	0.9300
C3—C4	1.372 (4)	C19—C20	1.378 (4)
C3—H3	0.9300	C19—O3	1.381 (3)
C4—C5	1.373 (4)	C20—C21	1.374 (4)
C4—H4	0.9300	C20—H20	0.9300
C5—C6	1.366 (4)	C21—H21	0.9300
C5—O1	1.396 (3)	C22—C27	1.375 (4)
C6—C7	1.374 (4)	C22—C23	1.378 (4)
C6—H6	0.9300	C22—O3	1.384 (4)
C7—H7	0.9300	C23—C24	1.374 (4)
C8—C13	1.375 (4)	C23—H23	0.9300
C8—C9	1.372 (4)	C24—C25	1.381 (4)
C8—O1	1.378 (3)	C24—H24	0.9300
C9—C10	1.377 (4)	C25—C26	1.378 (4)
C9—H9	0.9300	C25—C28	1.464 (4)
C10—C11	1.380 (5)	C26—C27	1.373 (4)
C10—H10	0.9300	C26—H26	0.9300
C11—C12	1.369 (4)	C27—H27	0.9300
C11—C14	1.520 (5)	C28—O4	1.200 (4)
C12—C13	1.381 (4)	C28—H28	0.9300
C12—H12	0.9300	N1—N2	1.349 (3)
C13—H13	0.9300	N1—H1N	0.91 (3)
C14—O2	1.166 (4)	N2—N3	1.294 (3)
C14—H14	0.9300	N3—N4	1.363 (3)
C15—N8	1.325 (3)	N5—N6	1.343 (3)
C15—N5	1.332 (3)	N5—H2N	0.90 (3)
C15—C16	1.455 (4)	N6—N7	1.291 (3)
C16—C17	1.385 (4)	N7—N8	1.362 (3)
N4—C1—N1	107.5 (2)	C18—C17—H17	119.4
N4—C1—C2	126.2 (2)	C16—C17—H17	119.4
N1—C1—C2	126.3 (2)	C19—C18—C17	120.1 (3)

supplementary materials

C3—C2—C7	119.0 (3)	C19—C18—H18	119.9
C3—C2—C1	119.8 (2)	C17—C18—H18	119.9
C7—C2—C1	121.1 (2)	C18—C19—C20	120.1 (3)
C4—C3—C2	120.8 (3)	C18—C19—O3	124.1 (3)
C4—C3—H3	119.6	C20—C19—O3	115.7 (3)
C2—C3—H3	119.6	C21—C20—C19	119.7 (3)
C5—C4—C3	119.0 (3)	C21—C20—H20	120.1
C5—C4—H4	120.5	C19—C20—H20	120.1
C3—C4—H4	120.5	C20—C21—C16	121.2 (3)
C6—C5—C4	121.5 (3)	C20—C21—H21	119.4
C6—C5—O1	117.8 (3)	C16—C21—H21	119.4
C4—C5—O1	120.6 (3)	C27—C22—C23	120.9 (3)
C5—C6—C7	119.3 (3)	C27—C22—O3	117.3 (3)
C5—C6—H6	120.3	C23—C22—O3	121.6 (3)
C7—C6—H6	120.3	C24—C23—C22	119.3 (3)
C6—C7—C2	120.3 (3)	C24—C23—H23	120.4
C6—C7—H7	119.8	C22—C23—H23	120.4
C2—C7—H7	119.8	C23—C24—C25	120.7 (3)
C13—C8—C9	121.2 (3)	C23—C24—H24	119.6
C13—C8—O1	122.7 (3)	C25—C24—H24	119.6
C9—C8—O1	116.1 (3)	C24—C25—C26	119.1 (3)
C8—C9—C10	118.7 (3)	C24—C25—C28	121.5 (3)
C8—C9—H9	120.7	C26—C25—C28	119.4 (3)
C10—C9—H9	120.7	C27—C26—C25	120.9 (3)
C9—C10—C11	121.6 (3)	C27—C26—H26	119.5
C9—C10—H10	119.2	C25—C26—H26	119.5
C11—C10—H10	119.2	C26—C27—C22	119.2 (3)
C12—C11—C10	118.1 (3)	C26—C27—H27	120.4
C12—C11—C14	115.6 (4)	C22—C27—H27	120.4
C10—C11—C14	126.3 (4)	O4—C28—C25	124.9 (4)
C11—C12—C13	121.7 (3)	O4—C28—H28	117.5
C11—C12—H12	119.2	C25—C28—H28	117.5
C13—C12—H12	119.2	C1—N1—N2	109.5 (2)
C8—C13—C12	118.6 (3)	C1—N1—H1N	133 (2)
C8—C13—H13	120.7	N2—N1—H1N	117 (2)
C12—C13—H13	120.7	N3—N2—N1	106.1 (2)
O2—C14—C11	119.7 (5)	N2—N3—N4	110.5 (2)
O2—C14—H14	120.2	C1—N4—N3	106.4 (2)
C11—C14—H14	120.2	C15—N5—N6	109.9 (2)
N8—C15—N5	107.1 (2)	C15—N5—H2N	129.8 (18)
N8—C15—C16	127.5 (2)	N6—N5—H2N	120.1 (18)
N5—C15—C16	125.4 (2)	N7—N6—N5	106.0 (2)
C17—C16—C21	117.8 (3)	N6—N7—N8	110.5 (2)
C17—C16—C15	120.6 (2)	C15—N8—N7	106.5 (2)
C21—C16—C15	121.6 (3)	C8—O1—C5	118.2 (2)
C18—C17—C16	121.2 (3)	C19—O3—C22	119.9 (2)
N4—C1—C2—C3	-17.7 (4)	C19—C20—C21—C16	0.5 (5)
N1—C1—C2—C3	161.4 (3)	C17—C16—C21—C20	-0.1 (4)
N4—C1—C2—C7	163.8 (3)	C15—C16—C21—C20	178.0 (3)

N1—C1—C2—C7	−17.1 (4)	C27—C22—C23—C24	−1.0 (5)
C7—C2—C3—C4	0.3 (4)	O3—C22—C23—C24	−176.1 (3)
C1—C2—C3—C4	−178.3 (3)	C22—C23—C24—C25	0.1 (5)
C2—C3—C4—C5	−0.8 (5)	C23—C24—C25—C26	0.8 (5)
C3—C4—C5—C6	0.5 (5)	C23—C24—C25—C28	179.7 (3)
C3—C4—C5—O1	176.9 (3)	C24—C25—C26—C27	−0.8 (5)
C4—C5—C6—C7	0.3 (5)	C28—C25—C26—C27	−179.8 (3)
O1—C5—C6—C7	−176.1 (3)	C25—C26—C27—C22	−0.1 (5)
C5—C6—C7—C2	−0.9 (5)	C23—C22—C27—C26	1.0 (5)
C3—C2—C7—C6	0.6 (4)	O3—C22—C27—C26	176.3 (3)
C1—C2—C7—C6	179.1 (3)	C24—C25—C28—O4	1.9 (5)
C13—C8—C9—C10	−2.4 (5)	C26—C25—C28—O4	−179.2 (3)
O1—C8—C9—C10	178.8 (3)	N4—C1—N1—N2	0.0 (3)
C8—C9—C10—C11	0.8 (6)	C2—C1—N1—N2	−179.2 (3)
C9—C10—C11—C12	1.3 (5)	C1—N1—N2—N3	0.0 (3)
C9—C10—C11—C14	−176.4 (3)	N1—N2—N3—N4	−0.1 (3)
C10—C11—C12—C13	−1.8 (5)	N1—C1—N4—N3	−0.1 (3)
C14—C11—C12—C13	176.1 (3)	C2—C1—N4—N3	179.1 (3)
C9—C8—C13—C12	1.9 (5)	N2—N3—N4—C1	0.1 (3)
O1—C8—C13—C12	−179.4 (3)	N8—C15—N5—N6	−0.5 (3)
C11—C12—C13—C8	0.3 (5)	C16—C15—N5—N6	179.3 (3)
C12—C11—C14—O2	−177.7 (4)	C15—N5—N6—N7	0.3 (3)
C10—C11—C14—O2	0.0 (7)	N5—N6—N7—N8	−0.1 (3)
N8—C15—C16—C17	162.4 (3)	N5—C15—N8—N7	0.4 (3)
N5—C15—C16—C17	−17.4 (4)	C16—C15—N8—N7	−179.4 (3)
N8—C15—C16—C21	−15.7 (5)	N6—N7—N8—C15	−0.2 (3)
N5—C15—C16—C21	164.5 (3)	C13—C8—O1—C5	21.7 (4)
C21—C16—C17—C18	−0.8 (5)	C9—C8—O1—C5	−159.5 (3)
C15—C16—C17—C18	−178.9 (3)	C6—C5—O1—C8	−120.5 (3)
C16—C17—C18—C19	1.4 (6)	C4—C5—O1—C8	63.0 (4)
C17—C18—C19—C20	−1.0 (5)	C18—C19—O3—C22	−17.6 (5)
C17—C18—C19—O3	−177.2 (3)	C20—C19—O3—C22	166.1 (3)
C18—C19—C20—C21	0.1 (5)	C27—C22—O3—C19	128.5 (3)
O3—C19—C20—C21	176.6 (3)	C23—C22—O3—C19	−56.3 (4)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···N8 ⁱ	0.91 (3)	1.94 (3)	2.849 (4)	174 (2)
N5—H2N···N4 ⁱⁱ	0.90 (3)	2.03 (3)	2.924 (3)	176 (3)
C4—H4···O4	0.93	2.54	3.420 (4)	158
C17—H17···N3 ⁱⁱ	0.93	2.58	3.365 (4)	143
C23—H23···O2 ⁱⁱⁱ	0.93	2.42	3.262 (6)	151

Symmetry codes: (i) $x+1, y-1, z+1$; (ii) $x-1, y, z-1$; (iii) $-x, -y+1, -z+1$.

supplementary materials

Fig. 1

