

Bis[bis(2,2'-bipyridine- κ^2N,N')chlorido-copper(II)] bis(μ -2,6-pyridinedicarboxylato)- $\kappa^4O^2,N,O^6:O^6;\kappa^4O^2:O^2,N,O^6$ -bis[aquadichloridobismuthate(III)] pentahydrate

Hong-Wei Wang,^a Wan-Lan Liu^b and Yu-Quan Feng^{a*}

^aCollege of Chemistry and Pharmacy Engineering, Nanyang Normal University, Nanyang 473061, People's Republic of China, and ^bDepartment of Equipment and Lab Administration, Nanyang Normal University, Nanyang 473061, People's Republic of China

Correspondence e-mail: yqfeng2008@126.com


Received 17 October 2011; accepted 28 October 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.007$ Å; H-atom completeness 81%; R factor = 0.025; wR factor = 0.063; data-to-parameter ratio = 13.9.

In the title compound, $[CuCl(C_{10}H_8N_2)_2][Bi_2Cl_4(C_7H_3NO_4)_2(H_2O)_2] \cdot 5H_2O$, the dianion $[Bi_2Cl_4(C_7H_3NO_4)_2(H_2O)_2]^{2-}$ is located about an inversion center. The Cu^{II} atom of the cation is coordinated by four N atoms of the two chelating 2,2'-bipyridine ligands and one Cl^- ion, completing a distorted trigonal-bipyramidal coordination environment. In the anion, each Bi^{III} atom is seven-coordinate and is bonded to a tridentate pyridine-2,6-dicarboxylate ligand, a water molecule, two chloride ions and a bridging carboxylate O atom of another carboxylate ligand. The coordination geometry of Bi^{III} is distorted pentagonal-bipyramidal with the Cl^- ions located in axial positions. The structure of the dianion is additionally stabilized by an intramolecular $O-H \cdots O$ hydrogen bond between the coordinated water molecule and carboxylate O atom. In the crystal, $O-H \cdots O$ hydrogen bonds occur. The H atoms of the solvent water molecules could not be located.

Related literature

For examples of bismuth(III) coordination compounds, see: Sun *et al.* (2004); Jiang *et al.* (2006); Meng & Zhang (2011).



Experimental

Crystal data

$[CuCl(C_{10}H_8N_2)_2][Bi_2Cl_4(C_7H_3NO_4)_2(H_2O)_2] \cdot 5H_2O$
 $M_r = 1838.72$
 Monoclinic, $C2/c$
 $a = 22.880$ (6) Å
 $b = 22.044$ (6) Å
 $c = 13.628$ (4) Å

$\beta = 110.130$ (4)°
 $V = 6454$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 6.41$ mm⁻¹
 $T = 296$ K
 $0.18 \times 0.16 \times 0.15$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2008)
 $T_{min} = 0.392$, $T_{max} = 0.447$

16290 measured reflections
 5684 independent reflections
 4963 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.063$
 $S = 1.04$
 5684 reflections
 408 parameters
 3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{max} = 0.88$ e Å⁻³
 $\Delta\rho_{min} = -0.74$ e Å⁻³

Table 1

Selected bond lengths (Å).

Bi1—O3	2.297 (3)	Cl3—Cu1	2.2962 (14)
Bi1—N1	2.385 (3)	Cu1—N4	1.983 (3)
Bi1—O1	2.485 (3)	Cu1—N3	1.994 (4)
Bi1—O5	2.531 (4)	Cu1—N5	2.107 (3)
Bi1—Cl2	2.6174 (15)	Cu1—N2	2.118 (4)
Bi1—Cl1	2.7479 (16)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O5—H1 \cdots O3W	0.83 (2)	1.92 (3)	2.733 (7)	169 (11)
O5—H2 \cdots O2 ⁱ	0.83 (2)	2.04 (6)	2.777 (5)	147 (10)

Symmetry code: (i) $-x, y, -z + \frac{3}{2}$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2419).

References

- Bruker (2008). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Jiang, Q. Y., Shen, J. & Zhong, G. Q. (2006). *Chin. J. Prog. Chem.* **18**, 1634–1645.
 Meng, Z.-H. & Zhang, S.-S. (2011). *Acta Cryst.* **E67**, m1402–m1403.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sun, H., Zhang, L. & Szeto, K. Y. (2004). *Met. Ions Biol. Syst.* **41**, 333–378.

supplementary materials

Acta Cryst. (2011). E67, m1651 [doi:10.1107/S1600536811045259]

Bis[bis(2,2'-bipyridine- κ^2N,N')chloridocopper(II)] bis(μ -2,6-pyridinedicarboxylato)- $\kappa^4O^2,N,O^6:O^6;\kappa^4O^2:O^2,N,O^6$ -bis[aquadichloridobismuthate(III)] pentahydrate

H.-W. Wang, W.-L. Liu and Y.-Q. Feng

Comment

Increasing attention has been paid to bismuth(III) coordination compounds in recent years due to their fascinating structural architectures and potential applications. As an expansion of bismuth(III) coordination compounds, recently, we have successfully isolated a novel compound $[\text{Cu}(\text{C}_{10}\text{H}_8\text{N}_2)_2\text{Cl}]_2[\text{Bi}_2(\text{C}_7\text{H}_3\text{NO}_4)_2(\text{H}_2\text{O})_2\text{Cl}_4] \cdot 5\text{H}_2\text{O}$.

The title compound consists of the coordination cations $[\text{Cu}(\text{C}_{10}\text{H}_8\text{N}_2)_2\text{Cl}]^+$, coordination dianions $[\text{Bi}_2(\text{C}_7\text{H}_3\text{NO}_4)_2(\text{H}_2\text{O})_2\text{Cl}_4]^{2-}$ and solvent water molecules (Fig. 1). In coordination cation $[\text{Cu}(\text{C}_{10}\text{H}_8\text{N}_2)_2\text{Cl}]^+$, the Cu^{II} center is coordinated by one Cl atom and four N atoms from two $\text{C}_{10}\text{H}_8\text{N}_2$ ligands. In dianion $[\text{Bi}_2(\text{C}_7\text{H}_3\text{NO}_4)_2(\text{H}_2\text{O})_2\text{Cl}_4]^{2-}$, each Bi^{III} center is coordinated by one N atom, two Cl atoms, three O atoms and one water molecule. It is worth noting that one of the carboxylate groups acts as bridging group between two Bi centers. The structure of the dianion is additionally stabilized by intramolecular hydrogen bonds $[\text{O}5\text{—H}2\cdots\text{O}2(-x,y,-z + 3/2)$ and $\text{O}(5)\text{—H}(1)\cdots\text{O}(3\text{ W})]$. There exist the electrostatic interactions between the coordination cations and dianions, the crystal packing of the title compound along the *c* axis is shown in Fig. 2. Obviously, these electrostatic interactions and hydrogen bonds play a crucial role in the chemical stability of the title compound.

Experimental

All chemicals were of reagent grade quality obtained from commercial sources and used without further purification. The title compound was synthesized from a mixture of $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$ (0.49 g, 1 mmol), $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ (0.26 g, 1.5 mmol), 2,6-pyridinedicarboxylic acid (0.17 g, 1 mmol), 2,2'-bipyridine (0.24 g, 1.5 mmol) and H_2O (12 g, 667 mmol) in a molar ratio of 1: 1.5: 1: 1.5: 667 by hydrothermal reaction. The mixture was stirred for half an hour, and then transferred into a Teflon-lined stainless steel autoclave (50 ml) and treated at 170 °C for 5 days. After the mixture was slowly cooled to room temperature, blue crystals were obtained. Yield: 75% (based on Bi).

Refinement

The H atoms bonded to C were positioned geometrically and refined using a riding model, with $\text{C—H} = 0.93 \text{ \AA}$ and with $U_{\text{iso}}(\text{H}) = 1.2$ times $U_{\text{eq}}(\text{C})$. The H atoms bonded to O5 atom were located from Fourier difference maps and refined with distance restraints of $\text{O}5\text{—H}1 = 0.83$ (2), $\text{O}5\text{—H}2 = 0.83$ (2), $\text{H}1\cdots\text{H}2 = 1.37$ (2) Å. The H atoms bonded to O1W, O2W and O3W were not located in Fourier difference maps, most probably due to their disorder.

Figures




Fig. 1. View of the title compound with displacement ellipsoids drawn at the 30% probability level (symmetry code (i): $-x, y, -z+3/2$).




Fig. 2. Crystal packing of the title compound along the c axis.

**Bis[bis(2,2'-bipyridine- κ^2N,N')chloridocopper(II)] bis(μ -2,6-pyridinedicarboxylato)- $\kappa^4O^2,N,O^6:O^6$;
 $\kappa^4O^2:O^2,N,O^6$ - bis[aquadichloridobismuthate(III)] pentahydrate**

Crystal data

$[\text{CuCl}(\text{C}_{10}\text{H}_8\text{N}_2)_2]_2[\text{Bi}_2\text{Cl}_4(\text{C}_7\text{H}_3\text{NO}_4)_2(\text{H}_2\text{O})_2] \cdot 5\text{H}_2\text{O}$ $F(000) = 3528$

$M_r = 1838.72$

$D_x = 1.882 \text{ Mg m}^{-3}$

Monoclinic, $C2/c$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Hall symbol: $-C 2yc$

Cell parameters from 8126 reflections

$a = 22.880 (6) \text{ \AA}$

$\theta = 2.4\text{--}28.1^\circ$

$b = 22.044 (6) \text{ \AA}$

$\mu = 6.41 \text{ mm}^{-1}$

$c = 13.628 (4) \text{ \AA}$

$T = 296 \text{ K}$

$\beta = 110.130 (4)^\circ$

Block, blue

$V = 6454 (3) \text{ \AA}^3$

$0.18 \times 0.16 \times 0.15 \text{ mm}$

$Z = 4$

Data collection

Bruker APEXII CCD
 diffractometer

5684 independent reflections

Radiation source: fine-focus sealed tube
 graphite

4963 reflections with $I > 2\sigma(I)$

φ and ω scans

$R_{\text{int}} = 0.025$

$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 1.8^\circ$

Absorption correction: multi-scan
 (SADABS; Bruker, 2008)

$h = -27 \rightarrow 23$

$T_{\text{min}} = 0.392$, $T_{\text{max}} = 0.447$

$k = -26 \rightarrow 26$

16290 measured reflections

$l = -16 \rightarrow 14$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.025$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.063$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0312P)^2 + 11.1454P]$
5684 reflections	where $P = (F_o^2 + 2F_c^2)/3$
408 parameters	$(\Delta/\sigma)_{\max} = 0.002$
3 restraints	$\Delta\rho_{\max} = 0.88 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.74 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Bi1	0.025844 (6)	0.242343 (7)	0.610971 (11)	0.02982 (6)
C1	0.12686 (18)	0.23152 (18)	0.8542 (3)	0.0326 (9)
C2	0.16677 (19)	0.24402 (17)	0.7868 (3)	0.0327 (9)
C3	0.23102 (19)	0.25200 (18)	0.8283 (4)	0.0402 (10)
H3A	0.2521	0.2502	0.9000	0.048*
C4	0.2625 (2)	0.2627 (2)	0.7598 (4)	0.0526 (13)
H4A	0.3055	0.2683	0.7856	0.063*
C5	0.2309 (2)	0.2652 (2)	0.6536 (4)	0.0457 (11)
H5A	0.2522	0.2712	0.6073	0.055*
C6	0.16695 (18)	0.25867 (17)	0.6173 (3)	0.0321 (9)
C7	0.1267 (2)	0.26348 (19)	0.5032 (4)	0.0379 (10)
C8	0.1129 (2)	0.0957 (2)	0.1789 (4)	0.0517 (12)
H8A	0.0951	0.0875	0.1079	0.062*
C9	0.0795 (2)	0.1308 (2)	0.2266 (4)	0.0578 (13)
H9A	0.0403	0.1458	0.1883	0.069*
C10	0.1061 (3)	0.1424 (2)	0.3317 (5)	0.0612 (14)
H10A	0.0845	0.1646	0.3663	0.073*
C11	0.1655 (2)	0.1207 (2)	0.3858 (4)	0.0525 (12)
H11A	0.1842	0.1287	0.4567	0.063*
C12	0.1965 (2)	0.08714 (19)	0.3336 (4)	0.0416 (10)
C13	0.2607 (2)	0.0635 (2)	0.3840 (3)	0.0413 (10)
C14	0.2988 (2)	0.0786 (2)	0.4839 (4)	0.0592 (13)
H14A	0.2846	0.1048	0.5242	0.071*
C15	0.3586 (2)	0.0543 (3)	0.5239 (4)	0.0638 (14)
H15A	0.3848	0.0646	0.5907	0.077*
C16	0.3783 (2)	0.0150 (2)	0.4636 (4)	0.0610 (14)
H16A	0.4180	-0.0019	0.4889	0.073*
C17	0.3379 (2)	0.0010 (2)	0.3640 (4)	0.0538 (12)
H17A	0.3510	-0.0259	0.3232	0.065*

supplementary materials

C18	0.1123 (2)	-0.0074 (2)	-0.0053 (4)	0.0475 (11)
H18A	0.1014	-0.0336	0.0390	0.057*
C19	0.0703 (2)	0.0024 (2)	-0.1038 (4)	0.0530 (12)
H19A	0.0315	-0.0163	-0.1256	0.064*
C20	0.0872 (2)	0.0408 (2)	-0.1695 (4)	0.0538 (12)
H20A	0.0595	0.0486	-0.2364	0.065*
C21	0.1451 (2)	0.0674 (2)	-0.1356 (4)	0.0461 (11)
H21A	0.1572	0.0926	-0.1799	0.055*
C22	0.18524 (18)	0.05626 (17)	-0.0343 (3)	0.0349 (9)
C23	0.24907 (18)	0.08145 (18)	0.0105 (3)	0.0367 (10)
C24	0.2784 (2)	0.1136 (2)	-0.0478 (4)	0.0468 (11)
H24A	0.2573	0.1218	-0.1181	0.056*
C25	0.3389 (2)	0.1332 (2)	0.0002 (4)	0.0553 (13)
H25A	0.3594	0.1540	-0.0378	0.066*
C26	0.3683 (2)	0.1215 (2)	0.1044 (4)	0.0535 (12)
H26A	0.4089	0.1346	0.1385	0.064*
C27	0.3365 (2)	0.0900 (2)	0.1577 (4)	0.0488 (11)
H27A	0.3565	0.0826	0.2286	0.059*
Cl1	0.03209 (6)	0.36683 (7)	0.61824 (12)	0.0693 (4)
Cl2	0.03436 (7)	0.12583 (6)	0.58147 (13)	0.0695 (4)
Cl3	0.22828 (6)	-0.08940 (5)	0.17806 (10)	0.0560 (3)
Cu1	0.22536 (2)	0.01472 (2)	0.17652 (4)	0.04341 (14)
N1	0.13659 (14)	0.24776 (13)	0.6839 (3)	0.0299 (7)
N2	0.16967 (16)	0.07324 (16)	0.2309 (3)	0.0413 (8)
N3	0.28081 (16)	0.02474 (16)	0.3248 (3)	0.0440 (9)
N4	0.16815 (15)	0.01917 (15)	0.0295 (3)	0.0386 (8)
N5	0.27864 (15)	0.06980 (15)	0.1128 (3)	0.0390 (8)
O1	0.06762 (13)	0.23135 (14)	0.8042 (2)	0.0413 (7)
O2	0.15260 (13)	0.22296 (15)	0.9484 (2)	0.0471 (7)
O3	0.06808 (14)	0.25476 (14)	0.4818 (2)	0.0455 (8)
O4	0.15195 (14)	0.27517 (16)	0.4391 (2)	0.0516 (8)
O5	-0.06953 (17)	0.2428 (2)	0.4468 (3)	0.0731 (12)
O1W	0.0000	0.5052 (4)	0.7500	0.173 (4)
H1	-0.073 (5)	0.257 (5)	0.389 (4)	0.208*
H2	-0.103 (3)	0.233 (5)	0.454 (9)	0.208*
O3W	-0.0652 (3)	0.2915 (3)	0.2650 (4)	0.151 (3)
O2W	0.4613 (3)	-0.0920 (3)	0.3689 (4)	0.1270 (19)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Bi1	0.02219 (9)	0.04836 (10)	0.01923 (9)	0.00043 (6)	0.00754 (6)	-0.00092 (6)
C1	0.028 (2)	0.044 (2)	0.025 (2)	0.0032 (17)	0.0069 (17)	-0.0018 (17)
C2	0.029 (2)	0.038 (2)	0.029 (2)	0.0015 (15)	0.0080 (18)	-0.0032 (17)
C3	0.027 (2)	0.057 (3)	0.032 (3)	-0.0022 (17)	0.0046 (18)	-0.0015 (19)
C4	0.022 (2)	0.083 (3)	0.047 (3)	-0.005 (2)	0.004 (2)	0.007 (2)
C5	0.034 (2)	0.064 (3)	0.045 (3)	-0.005 (2)	0.021 (2)	0.002 (2)
C6	0.028 (2)	0.042 (2)	0.028 (2)	-0.0020 (16)	0.0118 (18)	-0.0032 (17)

C7	0.037 (2)	0.049 (2)	0.032 (2)	-0.0020 (18)	0.018 (2)	-0.0033 (19)
C8	0.055 (3)	0.054 (3)	0.044 (3)	0.006 (2)	0.014 (2)	0.003 (2)
C9	0.053 (3)	0.059 (3)	0.063 (4)	0.001 (2)	0.022 (3)	0.002 (3)
C10	0.069 (4)	0.059 (3)	0.067 (4)	0.000 (3)	0.039 (3)	-0.008 (3)
C11	0.064 (3)	0.051 (3)	0.046 (3)	-0.012 (2)	0.023 (2)	-0.009 (2)
C12	0.047 (3)	0.040 (2)	0.039 (3)	-0.0089 (19)	0.017 (2)	0.002 (2)
C13	0.049 (3)	0.046 (2)	0.030 (2)	-0.012 (2)	0.015 (2)	0.0052 (19)
C14	0.063 (3)	0.079 (4)	0.035 (3)	-0.017 (3)	0.016 (2)	-0.009 (3)
C15	0.059 (3)	0.089 (4)	0.035 (3)	-0.017 (3)	0.006 (3)	-0.002 (3)
C16	0.048 (3)	0.073 (4)	0.051 (3)	-0.005 (2)	0.002 (2)	0.014 (3)
C17	0.053 (3)	0.060 (3)	0.046 (3)	0.004 (2)	0.013 (2)	0.008 (2)
C18	0.043 (3)	0.055 (3)	0.041 (3)	-0.008 (2)	0.011 (2)	0.005 (2)
C19	0.038 (2)	0.062 (3)	0.052 (3)	-0.003 (2)	0.007 (2)	-0.003 (2)
C20	0.046 (3)	0.067 (3)	0.037 (3)	0.007 (2)	-0.001 (2)	-0.001 (2)
C21	0.052 (3)	0.049 (3)	0.037 (3)	0.005 (2)	0.016 (2)	0.006 (2)
C22	0.040 (2)	0.035 (2)	0.032 (2)	0.0012 (17)	0.0138 (18)	-0.0058 (18)
C23	0.045 (3)	0.033 (2)	0.036 (3)	0.0007 (17)	0.020 (2)	-0.0029 (18)
C24	0.060 (3)	0.046 (3)	0.038 (3)	-0.008 (2)	0.021 (2)	-0.003 (2)
C25	0.067 (3)	0.048 (3)	0.061 (4)	-0.019 (2)	0.036 (3)	-0.009 (2)
C26	0.049 (3)	0.052 (3)	0.060 (4)	-0.015 (2)	0.019 (3)	-0.009 (2)
C27	0.047 (3)	0.054 (3)	0.041 (3)	-0.005 (2)	0.009 (2)	-0.003 (2)
Cl1	0.0662 (9)	0.0669 (8)	0.0728 (10)	-0.0064 (6)	0.0214 (7)	-0.0097 (7)
Cl2	0.0781 (9)	0.0537 (7)	0.0808 (10)	0.0080 (6)	0.0326 (8)	0.0003 (7)
Cl3	0.0835 (9)	0.0496 (6)	0.0356 (6)	0.0040 (6)	0.0213 (6)	0.0012 (5)
Cu1	0.0458 (3)	0.0514 (3)	0.0292 (3)	-0.0035 (2)	0.0081 (2)	0.0028 (2)
N1	0.0232 (16)	0.0408 (19)	0.0261 (19)	0.0002 (12)	0.0092 (14)	-0.0019 (14)
N2	0.045 (2)	0.046 (2)	0.034 (2)	-0.0004 (16)	0.0153 (17)	0.0033 (16)
N3	0.044 (2)	0.050 (2)	0.033 (2)	-0.0016 (17)	0.0084 (17)	0.0047 (17)
N4	0.0390 (19)	0.044 (2)	0.031 (2)	-0.0048 (15)	0.0089 (15)	-0.0021 (16)
N5	0.0378 (19)	0.046 (2)	0.031 (2)	-0.0052 (15)	0.0085 (16)	-0.0026 (16)
O1	0.0284 (15)	0.070 (2)	0.0261 (16)	0.0004 (13)	0.0101 (12)	0.0002 (14)
O2	0.0369 (16)	0.077 (2)	0.0258 (17)	0.0039 (15)	0.0083 (13)	0.0036 (15)
O3	0.0317 (16)	0.080 (2)	0.0268 (17)	-0.0045 (13)	0.0126 (13)	-0.0020 (14)
O4	0.0481 (18)	0.082 (2)	0.0329 (18)	-0.0098 (16)	0.0247 (15)	-0.0006 (17)
O5	0.036 (2)	0.143 (4)	0.032 (2)	0.0002 (19)	0.0021 (16)	0.001 (2)
O1W	0.187 (9)	0.111 (6)	0.191 (11)	0.000	0.026 (8)	0.000
O3W	0.221 (6)	0.136 (5)	0.053 (3)	-0.077 (5)	-0.009 (4)	0.012 (3)
O2W	0.149 (5)	0.119 (4)	0.110 (5)	0.006 (4)	0.040 (4)	-0.011 (3)

Geometric parameters (Å, °)

Bi1—O3	2.297 (3)	C14—H14A	0.9300
Bi1—N1	2.385 (3)	C15—C16	1.373 (8)
Bi1—O1	2.485 (3)	C15—H15A	0.9300
Bi1—O5	2.531 (4)	C16—C17	1.390 (7)
Bi1—Cl2	2.6174 (15)	C16—H16A	0.9300
Bi1—Cl1	2.7479 (16)	C17—N3	1.336 (5)
C1—O2	1.229 (5)	C17—H17A	0.9300
C1—O1	1.290 (5)	C18—N4	1.336 (5)

supplementary materials

C1—C2	1.527 (6)	C18—C19	1.372 (6)
C2—N1	1.335 (5)	C18—H18A	0.9300
C2—C3	1.393 (6)	C19—C20	1.380 (7)
C3—C4	1.382 (7)	C19—H19A	0.9300
C3—H3A	0.9300	C20—C21	1.375 (6)
C4—C5	1.379 (7)	C20—H20A	0.9300
C4—H4A	0.9300	C21—C22	1.392 (6)
C5—C6	1.382 (6)	C21—H21A	0.9300
C5—H5A	0.9300	C22—N4	1.347 (5)
C6—N1	1.341 (5)	C22—C23	1.483 (5)
C6—C7	1.515 (6)	C23—N5	1.349 (5)
C7—O4	1.229 (5)	C23—C24	1.396 (6)
C7—O3	1.285 (5)	C24—C25	1.381 (6)
C8—N2	1.341 (6)	C24—H24A	0.9300
C8—C9	1.395 (7)	C25—C26	1.371 (7)
C8—H8A	0.9300	C25—H25A	0.9300
C9—C10	1.374 (7)	C26—C27	1.381 (6)
C9—H9A	0.9300	C26—H26A	0.9300
C10—C11	1.390 (7)	C27—N5	1.329 (5)
C10—H10A	0.9300	C27—H27A	0.9300
C11—C12	1.380 (6)	Cl3—Cu1	2.2962 (14)
C11—H11A	0.9300	Cu1—N4	1.983 (3)
C12—N2	1.355 (6)	Cu1—N3	1.994 (4)
C12—C13	1.485 (6)	Cu1—N5	2.107 (3)
C13—N3	1.360 (6)	Cu1—N2	2.118 (4)
C13—C14	1.380 (6)	O5—H1	0.83 (2)
C14—C15	1.393 (7)	O5—H2	0.83 (2)
O3—Bi1—N1	69.30 (11)	N3—C17—C16	122.3 (5)
O3—Bi1—O1	135.54 (10)	N3—C17—H17A	118.9
N1—Bi1—O1	66.26 (10)	C16—C17—H17A	118.9
O3—Bi1—O5	77.55 (13)	N4—C18—C19	122.7 (4)
N1—Bi1—O5	146.83 (13)	N4—C18—H18A	118.6
O1—Bi1—O5	146.85 (12)	C19—C18—H18A	118.6
O3—Bi1—Cl2	85.76 (8)	C18—C19—C20	118.2 (4)
N1—Bi1—Cl2	89.34 (8)	C18—C19—H19A	120.9
O1—Bi1—Cl2	92.60 (8)	C20—C19—H19A	120.9
O5—Bi1—Cl2	87.41 (12)	C19—C20—C21	119.8 (4)
O3—Bi1—Cl1	83.01 (8)	C19—C20—H20A	120.1
N1—Bi1—Cl1	84.28 (8)	C21—C20—H20A	120.1
O1—Bi1—Cl1	93.61 (8)	C20—C21—C22	119.2 (4)
O5—Bi1—Cl1	92.67 (12)	C20—C21—H21A	120.4
Cl2—Bi1—Cl1	168.47 (5)	C22—C21—H21A	120.4
O2—C1—O1	126.1 (4)	N4—C22—C21	120.6 (4)
O2—C1—C2	119.0 (3)	N4—C22—C23	115.2 (4)
O1—C1—C2	114.9 (4)	C21—C22—C23	124.2 (4)
N1—C2—C3	120.7 (4)	N5—C23—C24	121.1 (4)
N1—C2—C1	116.3 (3)	N5—C23—C22	115.3 (3)
C3—C2—C1	123.0 (4)	C24—C23—C22	123.6 (4)
C4—C3—C2	118.1 (4)	C25—C24—C23	119.3 (4)

C4—C3—H3A	121.0	C25—C24—H24A	120.4
C2—C3—H3A	121.0	C23—C24—H24A	120.4
C5—C4—C3	120.5 (4)	C26—C25—C24	119.2 (4)
C5—C4—H4A	119.7	C26—C25—H25A	120.4
C3—C4—H4A	119.7	C24—C25—H25A	120.4
C6—C5—C4	118.7 (4)	C25—C26—C27	118.6 (4)
C6—C5—H5A	120.6	C25—C26—H26A	120.7
C4—C5—H5A	120.6	C27—C26—H26A	120.7
N1—C6—C5	120.5 (4)	N5—C27—C26	123.3 (5)
N1—C6—C7	115.7 (3)	N5—C27—H27A	118.4
C5—C6—C7	123.8 (4)	C26—C27—H27A	118.4
O4—C7—O3	125.5 (4)	N4—Cu1—N3	170.69 (14)
O4—C7—C6	118.4 (4)	N4—Cu1—N5	80.20 (13)
O3—C7—C6	116.1 (4)	N3—Cu1—N5	95.56 (14)
N2—C8—C9	123.0 (5)	N4—Cu1—N2	93.21 (14)
N2—C8—H8A	118.5	N3—Cu1—N2	80.07 (15)
C9—C8—H8A	118.5	N5—Cu1—N2	107.23 (14)
C10—C9—C8	118.2 (5)	N4—Cu1—Cl3	93.77 (10)
C10—C9—H9A	120.9	N3—Cu1—Cl3	95.46 (11)
C8—C9—H9A	120.9	N5—Cu1—Cl3	124.09 (10)
C9—C10—C11	119.2 (5)	N2—Cu1—Cl3	128.65 (10)
C9—C10—H10A	120.4	C2—N1—C6	121.4 (3)
C11—C10—H10A	120.4	C2—N1—Bi1	121.7 (3)
C12—C11—C10	119.7 (5)	C6—N1—Bi1	116.8 (3)
C12—C11—H11A	120.2	C8—N2—C12	118.3 (4)
C10—C11—H11A	120.2	C8—N2—Cu1	129.0 (3)
N2—C12—C11	121.4 (4)	C12—N2—Cu1	112.6 (3)
N2—C12—C13	115.1 (4)	C17—N3—C13	119.5 (4)
C11—C12—C13	123.5 (4)	C17—N3—Cu1	123.7 (3)
N3—C13—C14	120.6 (4)	C13—N3—Cu1	116.4 (3)
N3—C13—C12	115.4 (4)	C18—N4—C22	119.4 (4)
C14—C13—C12	124.0 (4)	C18—N4—Cu1	123.7 (3)
C13—C14—C15	119.7 (5)	C22—N4—Cu1	116.5 (3)
C13—C14—H14A	120.1	C27—N5—C23	118.5 (4)
C15—C14—H14A	120.1	C27—N5—Cu1	129.0 (3)
C16—C15—C14	119.2 (5)	C23—N5—Cu1	112.3 (3)
C16—C15—H15A	120.4	C1—O1—Bi1	120.7 (2)
C14—C15—H15A	120.4	C7—O3—Bi1	121.7 (3)
C15—C16—C17	118.6 (5)	Bi1—O5—H1	128 (8)
C15—C16—H16A	120.7	Bi1—O5—H2	117 (8)
C17—C16—H16A	120.7	H1—O5—H2	114 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O5—H1 \cdots O3W	0.83 (2)	1.92 (3)	2.733 (7)	169 (11)
O5—H2 \cdots O2 ⁱ	0.83 (2)	2.04 (6)	2.777 (5)	147 (10)

Symmetry codes: (i) $-x, y, -z+3/2$.

Fig. 1




Fig. 2

