

Diaquabis(dihydrogen 3-azaniumyl-1-hydroxypropylidene-1,1-diphosphonato- $\kappa^2 O,O'$)cobalt(II)

Natalia V. Tsaryk,* Anatolij V. Dudko, Alexandra N. Kozachkova and Vasily I. Pekhnyo

Institute of General and Inorganic Chemistry, NAS Ukraine, Kyiv, prosp. Palladina 32/34, 03680, Ukraine

Correspondence e-mail: complex@ionc.kiev.ua


Received 21 September 2011; accepted 27 October 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.029; wR factor = 0.080; data-to-parameter ratio = 16.6.

The asymmetric unit of title compound, $[Co(C_3H_{10}NO_7P_2)_2(H_2O)_2]$, contains one half-molecule of the complex. The Co^{II} atom is located on an inversion centre and displays a distorted octahedral coordination geometry defined by four O atoms of two 3-azaniumyl-1-hydroxypropylidene-1,1-bisphosphonato ligands in the equatorial plane and two water molecules located in axial positions. The ligand molecules, which exist in a zwitterionic state, form two six-membered chelate rings with chair conformations. In the crystal, molecules are interlinked by $O-H\cdots O$ and $N-H\cdots O$ hydrogen bonds, forming a three-dimensional supramolecular structure.

Related literature

For general background to organic diphosphonic acids and their applications, see: Matczak-Jon & Videnova-Adrabinska (2005). For applications of bisphosphonate metal complexes in medicine, see: Matkovskaya *et al.* (2001). For a related structure, see: Bon *et al.* (2010). For bond-length data, see: Allen *et al.* (2004).



Experimental

Crystal data

$[Co(C_3H_{10}NO_7P_2)_2(H_2O)_2]$

$M_r = 563.08$

Monoclinic, $P2_1/c$

$a = 7.3292$ (2) Å

$b = 10.8172$ (3) Å

$c = 12.6403$ (3) Å

$\beta = 120.801$ (1)°
 $V = 860.79$ (4) Å³
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 1.46$ mm⁻¹
 $T = 100$ K
 $0.50 \times 0.25 \times 0.15$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{min} = 0.528$, $T_{max} = 0.811$

6688 measured reflections
2613 independent reflections
2273 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.080$
 $S = 1.04$
2613 reflections
157 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.70$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.41$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1O \cdots O5 ⁱ	0.85 (3)	2.05 (3)	2.845 (2)	155 (2)
O3—H3O \cdots O7 ⁱ	0.76 (3)	1.72 (3)	2.4627 (19)	167 (3)
O6—H6O \cdots O4 ⁱⁱ	0.74 (3)	1.78 (3)	2.5124 (18)	175 (3)
O8—H81 \cdots O4 ⁱⁱⁱ	0.89 (3)	1.88 (3)	2.7277 (18)	160 (2)
O8—H82 \cdots O1 ^{iv}	0.77 (3)	2.25 (3)	2.8953 (19)	142 (3)
N1—H2N \cdots O6 ^{iv}	0.88 (3)	2.15 (3)	2.975 (2)	156 (2)
N1—H3N \cdots O2 ⁱⁱⁱ	0.91 (3)	2.30 (3)	3.096 (2)	146 (2)
N1—H3N \cdots O5 ^v	0.91 (3)	2.32 (3)	3.030 (2)	134 (2)
N1—H1N \cdots O4 ^{vi}	0.87 (2)	2.33 (2)	3.071 (2)	143 (2)

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z - \frac{1}{2}$; (ii) $x - 1, -y + \frac{3}{2}, z - \frac{1}{2}$; (iii) $-x + 1, -y + 1, -z$; (iv) $-x, y - \frac{1}{2}, -z - \frac{1}{2}$; (v) $x + 1, y, z$; (vi) $-x + 1, y - \frac{1}{2}, -z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2010); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors gratefully acknowledge the support of this work by the Ukrainian National Academy of Sciences.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZ2263).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (2004). *International Tables for Crystallography, Volume C: Mathematical, physical and chemical tables, third edition*. Edited by E. Prince, pp. 778–789. Dordrecht/Boston/London: Kluwer.
- Bon, V. V., Dudko, A. V., Kozachkova, A. N., Pekhnyo, V. I. & Tsaryk, N. V. (2010). *Acta Cryst. E66*, m537–m538.
- Brandenburg, K. & Putz, H. (2010). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2007). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Matczak-Jon, E. & Videnova-Adrabinska, V. (2005). *Coord. Chem. Rev.* **249**, 2458–2488.
- Matkovskaya, T. A., Popov, K. I. & Yuryeva, E. A. (2001). *Bisphosphonates. Properties, Structure and Application in Medicine*, p. 223. Moscow: Khimiya.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst. 43*, 920–925.

supplementary materials

Acta Cryst. (2011). E67, m1694 [doi:10.1107/S1600536811045120]

Diaquabis(dihydrogen 3-azaniumyl-1-hydroxypropylidene-1,1-diphosphonato- $\kappa^2 O,O'$)cobalt(II)

N. V. Tsaryk, A. V. Dudko, A. N. Kozachkova and V. I. Pekhnyo

Comment

In recent years the design and synthesis of novel metal-organic coordination compounds based on gem-diphosphonic acids has attracted much interest due to their structural diversity and possible applications in many areas (Matczak-Jon & Videnova-Adrabinska, 2005). Particular attention has been paid to 3-amino-1-hydroxypropane-1,1-diyl)bis(phosphonic acid) (pamidronic acid) due to its biological activity and as a result of its usage as a drug to prevent calcification and inhibit bone resorption, *etc.* (Matkovskaya *et al.*, 2001).

The molecular structure of title complex is shown in Fig. 1; as illustrated the molecule of the complex forms discrete monomeric units. The asymmetric unit contains one-half of the formula unit $[Co(C_3H_{10}NO_7P_2)_2(H_2O)_2]$, with the Co atom lying on an inversion center.

The Co^{II} ion is coordinated in a slightly distorted octahedral geometry which consists of six oxygen atoms, two from water molecules located in the axial positions and four from the two phosphonate groups of two different ligands, which exist in zwitterionic form, creating two six-membered $[O, O]$ chelate rings. The Co—O bond lengths and the O—Co—O angles have expected values (Allen *et al.*, 2004) and conform well to the previously reported related structure (Bon *et al.*, 2010). In the packing, O—H···O and N—H···O hydrogen-bonds exist between the water molecules, phosphonate, hydroxyl O atoms and nitrogen atoms of the amino group (Fig. 2, Table 1). Thus, the molecules are interlinked by these hydrogen bonds to create a three-dimensional structure which partially influences and stabilizes the configuration of the molecule.

Experimental

Light pink crystals of the title compound were obtained from a mixture of $Co(NO_3)_2 \cdot 6H_2O$ (0,25 mmol; 0,0728 g) and 3-aminohydroxypropylidene-1,1-diphosphonic acid (0,5 mmol; 0,1175 g) in 20 ml H_2O . The combined solution was allowed to slowly evaporate. After 30 days, suitable crystals for X-ray data collection were obtained.

Refinement

All non-H atoms were refined with anisotropic displacement parameters. H atoms bonded to O and N atoms were located in a difference Fourier map. Their positions were refined freely whereas displacement parameters were fixed to $U_{iso}(H) = 1.5U_{eq}(N,O)$. The H1n atom of the amino group was refined with a distance restraint ($N—H = 0.91 \text{ \AA}$). Other H atoms bonded to C were positioned geometrically and refined using a riding model with $C—H = 0.99 \text{ \AA}$ for CH_2 with $U_{iso}(H) = 1.2U_{eq}(C)$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Crystal packing of the title compound viewed down the a axis, showing the three-dimensional chain structure. Hydrogen bonds are shown as dashed lines.

Diaquabis(dihydrogen 3-azaniumyl-1-hydroxypropylidene-1,1-diphosphonato- $\kappa^2 O,O'$)cobalt(II)

Crystal data

[Co(C ₃ H ₁₀ NO ₇ P ₂) ₂ (H ₂ O) ₂]	$F(000) = 578$
$M_r = 563.08$	$D_x = 2.172 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3543 reflections
$a = 7.3292 (2) \text{ \AA}$	$\theta = 2.7\text{--}30.6^\circ$
$b = 10.8172 (3) \text{ \AA}$	$\mu = 1.46 \text{ mm}^{-1}$
$c = 12.6403 (3) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 120.801 (1)^\circ$	Block, pink
$V = 860.79 (4) \text{ \AA}^3$	$0.50 \times 0.25 \times 0.15 \text{ mm}$
$Z = 2$	

Data collection

Bruker APEXII CCD diffractometer	2613 independent reflections
Radiation source: fine-focus sealed tube graphite	2273 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.020$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2007)	$\theta_{\text{max}} = 30.6^\circ, \theta_{\text{min}} = 2.7^\circ$
$T_{\text{min}} = 0.528, T_{\text{max}} = 0.811$	$h = -10 \rightarrow 9$
6688 measured reflections	$k = -12 \rightarrow 15$
	$l = -18 \rightarrow 12$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.029$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.080$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0452P)^2 + 0.4224P]$ where $P = (F_o^2 + 2F_c^2)/3$
2613 reflections	$(\Delta/\sigma)_{\max} = 0.003$
157 parameters	$\Delta\rho_{\max} = 0.70 \text{ e \AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.41 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.0000	0.5000	0.0000	0.01045 (9)
P1	0.25936 (6)	0.70145 (4)	-0.05843 (4)	0.00902 (10)
P2	-0.12574 (6)	0.58120 (4)	-0.28208 (4)	0.00964 (10)
O1	0.1630 (2)	0.72770 (13)	-0.29229 (12)	0.0156 (3)
H1O	0.127 (4)	0.802 (3)	-0.289 (2)	0.023*
O2	0.25165 (18)	0.60515 (12)	0.02492 (11)	0.0124 (2)
O3	0.1139 (2)	0.81197 (12)	-0.07370 (12)	0.0145 (3)
H3O	0.154 (4)	0.874 (3)	-0.079 (2)	0.022*
O4	0.48210 (18)	0.74524 (13)	-0.01964 (11)	0.0137 (3)
O5	-0.1323 (2)	0.49001 (12)	-0.19349 (11)	0.0143 (3)
O6	-0.2632 (2)	0.69798 (12)	-0.29892 (12)	0.0151 (3)
H6O	-0.334 (4)	0.712 (2)	-0.365 (2)	0.023*
O7	-0.1968 (2)	0.52889 (12)	-0.40857 (11)	0.0137 (3)
O8	0.1663 (2)	0.33228 (13)	0.02254 (13)	0.0158 (3)
H81	0.270 (4)	0.320 (2)	0.007 (2)	0.024*
H82	0.082 (4)	0.283 (2)	-0.014 (2)	0.024*
N1	0.4212 (3)	0.38826 (17)	-0.30480 (17)	0.0192 (3)

supplementary materials

H2N	0.356 (4)	0.323 (3)	-0.297 (2)	0.029*
H3N	0.553 (5)	0.394 (2)	-0.237 (3)	0.029*
H1N	0.439 (4)	0.383 (3)	-0.3675 (19)	0.029*
C1	0.1504 (2)	0.63551 (16)	-0.21320 (15)	0.0108 (3)
C2	0.2914 (3)	0.52507 (17)	-0.20331 (16)	0.0140 (3)
H2A	0.4370	0.5391	-0.1330	0.017*
H2B	0.2357	0.4493	-0.1858	0.017*
C3	0.3018 (3)	0.50443 (17)	-0.31959 (18)	0.0164 (4)
H3A	0.1565	0.4980	-0.3922	0.020*
H3B	0.3743	0.5750	-0.3325	0.020*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.01100 (15)	0.00938 (17)	0.00988 (16)	-0.00047 (11)	0.00457 (12)	0.00097 (12)
P1	0.00920 (18)	0.0085 (2)	0.00885 (19)	-0.00167 (14)	0.00423 (15)	-0.00129 (15)
P2	0.01040 (19)	0.0090 (2)	0.00774 (18)	-0.00077 (14)	0.00333 (15)	-0.00018 (15)
O1	0.0225 (6)	0.0115 (6)	0.0157 (6)	0.0003 (5)	0.0118 (5)	0.0027 (5)
O2	0.0120 (5)	0.0124 (6)	0.0107 (5)	-0.0017 (4)	0.0042 (4)	0.0020 (5)
O3	0.0171 (6)	0.0076 (6)	0.0190 (6)	-0.0005 (5)	0.0094 (5)	-0.0013 (5)
O4	0.0110 (5)	0.0168 (7)	0.0133 (6)	-0.0049 (4)	0.0063 (5)	-0.0040 (5)
O5	0.0163 (6)	0.0141 (6)	0.0103 (6)	-0.0043 (5)	0.0052 (5)	0.0014 (5)
O6	0.0148 (6)	0.0147 (6)	0.0109 (6)	0.0051 (5)	0.0031 (5)	0.0004 (5)
O7	0.0162 (6)	0.0125 (6)	0.0092 (5)	-0.0015 (5)	0.0041 (5)	-0.0030 (5)
O8	0.0139 (6)	0.0134 (6)	0.0210 (6)	-0.0007 (5)	0.0096 (5)	-0.0009 (5)
N1	0.0186 (7)	0.0182 (8)	0.0237 (8)	-0.0020 (6)	0.0129 (7)	-0.0071 (7)
C1	0.0114 (7)	0.0099 (8)	0.0103 (7)	-0.0005 (6)	0.0050 (6)	0.0001 (6)
C2	0.0155 (7)	0.0125 (8)	0.0125 (7)	0.0038 (6)	0.0061 (6)	-0.0007 (7)
C3	0.0181 (8)	0.0156 (9)	0.0180 (9)	0.0017 (6)	0.0111 (7)	-0.0008 (7)

Geometric parameters (\AA , $^\circ$)

Co1—O2 ⁱ	2.0494 (12)	O1—H1O	0.85 (3)
Co1—O2	2.0494 (12)	O3—H3O	0.76 (3)
Co1—O8 ⁱ	2.1221 (14)	O6—H6O	0.74 (3)
Co1—O8	2.1221 (14)	O8—H81	0.89 (3)
Co1—O5 ⁱ	2.1225 (13)	O8—H82	0.77 (3)
Co1—O5	2.1225 (13)	N1—C3	1.487 (3)
P1—O2	1.5031 (13)	N1—H2N	0.88 (3)
P1—O4	1.5199 (12)	N1—H3N	0.91 (3)
P1—O3	1.5469 (14)	N1—H1N	0.870 (17)
P1—C1	1.8367 (17)	C1—C2	1.542 (2)
P2—O5	1.5115 (13)	C2—C3	1.527 (3)
P2—O7	1.5149 (13)	C2—H2A	0.9900
P2—O6	1.5609 (14)	C2—H2B	0.9900
P2—C1	1.8422 (16)	C3—H3A	0.9900
O1—C1	1.448 (2)	C3—H3B	0.9900
O2 ⁱ —Co1—O2	180.0	P2—O5—Co1	130.65 (7)

O2 ⁱ —Co1—O8 ⁱ	92.52 (5)	P2—O6—H6O	111 (2)
O2—Co1—O8 ⁱ	87.48 (5)	Co1—O8—H81	126.4 (17)
O2 ⁱ —Co1—O8	87.48 (5)	Co1—O8—H82	107 (2)
O2—Co1—O8	92.52 (5)	H81—O8—H82	107 (2)
O8 ⁱ —Co1—O8	180.0	C3—N1—H2N	111.7 (18)
O2 ⁱ —Co1—O5 ⁱ	92.85 (5)	C3—N1—H3N	109.5 (17)
O2—Co1—O5 ⁱ	87.15 (5)	H2N—N1—H3N	109 (2)
O8 ⁱ —Co1—O5 ⁱ	90.19 (5)	C3—N1—H1N	107.2 (19)
O8—Co1—O5 ⁱ	89.81 (5)	H2N—N1—H1N	113 (2)
O2 ⁱ —Co1—O5	87.15 (5)	H3N—N1—H1N	106 (2)
O2—Co1—O5	92.85 (5)	O1—C1—C2	108.19 (14)
O8 ⁱ —Co1—O5	89.81 (5)	O1—C1—P1	108.73 (11)
O8—Co1—O5	90.19 (5)	C2—C1—P1	107.85 (11)
O5 ⁱ —Co1—O5	180.00 (7)	O1—C1—P2	109.67 (11)
O2—P1—O4	114.17 (7)	C2—C1—P2	108.63 (12)
O2—P1—O3	110.62 (8)	P1—C1—P2	113.62 (9)
O4—P1—O3	110.93 (8)	C3—C2—C1	113.39 (14)
O2—P1—C1	109.00 (8)	C3—C2—H2A	108.9
O4—P1—C1	105.91 (8)	C1—C2—H2A	108.9
O3—P1—C1	105.73 (8)	C3—C2—H2B	108.9
O5—P2—O7	114.46 (8)	C1—C2—H2B	108.9
O5—P2—O6	111.46 (8)	H2A—C2—H2B	107.7
O7—P2—O6	108.02 (7)	N1—C3—C2	108.60 (15)
O5—P2—C1	107.58 (7)	N1—C3—H3A	110.0
O7—P2—C1	108.58 (8)	C2—C3—H3A	110.0
O6—P2—C1	106.40 (8)	N1—C3—H3B	110.0
C1—O1—H1O	119.0 (17)	C2—C3—H3B	110.0
P1—O2—Co1	129.11 (7)	H3A—C3—H3B	108.4
P1—O3—H3O	115 (2)		
O4—P1—O2—Co1	−170.27 (9)	O4—P1—C1—C2	61.76 (13)
O3—P1—O2—Co1	63.77 (11)	O3—P1—C1—C2	179.55 (12)
C1—P1—O2—Co1	−52.07 (12)	O2—P1—C1—P2	58.97 (11)
O8 ⁱ —Co1—O2—P1	−55.98 (10)	O4—P1—C1—P2	−177.76 (9)
O8—Co1—O2—P1	124.02 (10)	O3—P1—C1—P2	−59.97 (11)
O5 ⁱ —Co1—O2—P1	−146.30 (11)	O5—P2—C1—O1	−177.36 (11)
O5—Co1—O2—P1	33.70 (11)	O7—P2—C1—O1	58.25 (13)
O7—P2—O5—Co1	166.92 (9)	O6—P2—C1—O1	−57.80 (13)
O6—P2—O5—Co1	−70.11 (12)	O5—P2—C1—C2	64.58 (13)
C1—P2—O5—Co1	46.17 (13)	O7—P2—C1—C2	−59.81 (13)
O2 ⁱ —Co1—O5—P2	148.83 (11)	O6—P2—C1—C2	−175.86 (11)
O2—Co1—O5—P2	−31.17 (11)	O5—P2—C1—P1	−55.46 (11)
O8 ⁱ —Co1—O5—P2	56.30 (11)	O7—P2—C1—P1	−179.85 (8)
O8—Co1—O5—P2	−123.70 (11)	O6—P2—C1—P1	64.10 (11)
O2—P1—C1—O1	−178.61 (10)	O1—C1—C2—C3	−31.88 (19)
O4—P1—C1—O1	−55.34 (13)	P1—C1—C2—C3	−149.34 (13)
O3—P1—C1—O1	62.45 (12)	P2—C1—C2—C3	87.10 (16)

supplementary materials


O2—P1—C1—C2 -61.51 (13) C1—C2—C3—N1 -173.76 (14)
Symmetry codes: (i) $-x, -y+1, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
O1—H1O \cdots O5 ⁱⁱ	0.85 (3)	2.05 (3)	2.845 (2)
O3—H3O \cdots O7 ⁱⁱ	0.76 (3)	1.72 (3)	2.4627 (19)
O6—H6O \cdots O4 ⁱⁱⁱ	0.74 (3)	1.78 (3)	2.5124 (18)
O8—H81 \cdots O4 ^{iv}	0.89 (3)	1.88 (3)	2.7277 (18)
O8—H82 \cdots O1 ^v	0.77 (3)	2.25 (3)	2.8953 (19)
N1—H2N \cdots O6 ^v	0.88 (3)	2.15 (3)	2.975 (2)
N1—H3N \cdots O2 ^{iv}	0.91 (3)	2.30 (3)	3.096 (2)
N1—H3N \cdots O5 ^{vi}	0.91 (3)	2.32 (3)	3.030 (2)
N1—H1N \cdots O4 ^{vii}	0.87 (2)	2.33 (2)	3.071 (2)

Symmetry codes: (ii) $-x, y+1/2, -z-1/2$; (iii) $x-1, -y+3/2, z-1/2$; (iv) $-x+1, -y+1, -z$; (v) $-x, y-1/2, -z-1/2$; (vi) $x+1, y, z$; (vii) $-x+1, y-1/2, -z-1/2$.

Fig. 1



supplementary materials

Fig. 2

