

Bis(chloroacetato- $\kappa^2 O,O'$)bis(2-fluoro-benzyl- κC^1)tin(IV)

Aixia Deng,* Mouyong Teng, Qian Xie and Nana Yan

College of Materials Science and Engineering, Liaocheng University, Shandong 252059, People's Republic of China
Correspondence e-mail: dengaixia@lcu.edu.cn


Received 4 November 2011; accepted 28 November 2011

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.006$ Å;
R factor = 0.033; wR factor = 0.071; data-to-parameter ratio = 14.3.

In the title complex, $[Sn(C_2H_2ClO_2)_2(C_7H_6F)_2]$, the Sn^{IV} atom is located on a twofold rotation axis and forms a strongly distorted *trans*-octahedral geometry. The equatorial plane is defined by two chelating chloroacetate ligands with asymmetrical $Sn-O$ bond lengths, while the axial positions are occupied by the C atoms of two 2-fluorobenzyl groups. In the crystal, infinite chains in the [010] direction are formed through intermolecular $Sn \cdots O$ interactions [$Sn \cdots O$ separation = 3.682 (3) Å].

Related literature

For details of the synthesis, see: Zhang *et al.* (2007).



Experimental

Crystal data

$[Sn(C_2H_2ClO_2)_2(C_7H_6F)_2]$	$V = 1997.2$ (4) Å ³
$M_r = 523.90$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 17.3841$ (18) Å	$\mu = 1.59$ mm ⁻¹
$b = 5.0480$ (8) Å	$T = 298$ K
$c = 22.808$ (2) Å	$0.29 \times 0.15 \times 0.12$ mm
$\beta = 93.760$ (1)°	

Data collection

Bruker SMART 1000 CCD area-detector diffractometer	4738 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	1754 independent reflections
$R_{\text{int}} = 0.032$	1551 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.656$, $T_{\max} = 0.833$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$	123 parameters
$wR(F^2) = 0.071$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\max} = 0.71$ e Å ⁻³
1754 reflections	$\Delta\rho_{\min} = -0.26$ e Å ⁻³

Table 1
Selected bond lengths (Å).

$Sn1-O1$	2.109 (2)	$Sn1-O2$	2.537 (3)
$Sn1-C3$	2.121 (4)		

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the State Key Laboratory of Crystal Materials (SRT11055HX2), Liaocheng University, China, and the Liaocheng University Foundation (xo9013) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2395).

References

- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhang, J.-H., Ma, C.-L. & Zhang, R.-F. (2007). *Acta Cryst. E* **63**, m2161.

supplementary materials

Acta Cryst. (2011). E67, m1905 [doi:10.1107/S1600536811051002]

Bis(chloroacetato- κ^2O,O')bis(2-fluorobenzyl- κC^1)tin(IV)

A. Deng, M. Teng, Q. Xie and N. Yan

Comment

The title complex was obtained using a route similar to that used for the synthesis of a trinuclear tin complex (Zhang *et al.*, 2007). The title complex has 2-fold symmetry, with the Sn atom placed on the crystallographic symmetry axis (Fig. 1). Selected bond lengths and angles are given in table 1. The coordination geometry of tin can be described as a distorted *trans*-octahedron geometry, with two C atoms of 2-fluorobenzyl groups occupying the axial positions. The C3—Sn1—C3ⁱ bond angle of 133.6 (2)° (symmetry code *i*: 1 - *x*, *y*, 1/2 - *z*) reflects the distortion from octahedral geometry. The equatorial plane is defined by four O atoms of two symmetry-related chloroacetate ligands. The bond lengths in the equatorial plane are Sn1—O1 = 2.109 (2) and Sn1—O2 = 2.537 (3) Å, reflecting the asymmetrical coordination of acetate groups.

The complex forms infinite chains containing Sn₂O₂ rings, through intermolecular Sn···O contacts, characterized by separations Sn···O = 3.682 (3) Å (Fig. 2).

Experimental

Chloroacetic acid (2 mmol) was added to a sodium ethoxide solution (2 mmol, 20 ml of ethanol), and the mixture was stirred for 30 min. Then, 1 mmol of bis(2-fluorobenzyl)tin(IV)dichloride (Zhang *et al.*, 2007) was added to the mixture, continuing the reaction for 12 h at 318 K. After cooling down to room temperature, the reaction was filtered off. The solvent of the filtrate was gradually removed by evaporation under vacuum, until a solid product was obtained. The solid was recrystallized from ether-dichloromethane and colourless crystals suitable for X-ray diffraction were obtained (m.p. 464 K). Analysis calculated for C₁₈H₁₆Cl₂F₂O₄Sn: C 41.26, H 3.08%; found: C 41.29, H 3.06%.

Refinement

All H atoms were placed geometrically and treated as riding on their parent atoms with C—H bond lengths fixed to 0.93 Å for aromatic CH and 0.97 Å for methylene CH₂. Isotropic displacement parameters for H atoms were calculated as $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids.

supplementary materials




Fig. 2. The unit cell of the title compound.

Bis(chloroacetato- $\kappa^2 O,O'$)bis(2-fluorobenzyl- κC^1)tin(IV)

Crystal data

[Sn(C ₂ H ₂ ClO ₂) ₂ (C ₇ H ₆ F) ₂]	$F(000) = 1032$
$M_r = 523.90$	$D_x = 1.742 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	Melting point: 464 K
Hall symbol: -C 2yc	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 17.3841 (18) \text{ \AA}$	Cell parameters from 1705 reflections
$b = 5.0480 (8) \text{ \AA}$	$\theta = 2.9\text{--}25.0^\circ$
$c = 22.808 (2) \text{ \AA}$	$\mu = 1.59 \text{ mm}^{-1}$
$\beta = 93.760 (1)^\circ$	$T = 298 \text{ K}$
$V = 1997.2 (4) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.29 \times 0.15 \times 0.12 \text{ mm}$

Data collection

Bruker SMART 1000 CCD area-detector diffractometer	1754 independent reflections
Radiation source: fine-focus sealed tube graphite	1551 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.032$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 1.8^\circ$
$T_{\text{min}} = 0.656, T_{\text{max}} = 0.833$	$h = -20 \rightarrow 17$
4738 measured reflections	$k = -5 \rightarrow 6$
	$l = -27 \rightarrow 22$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.033$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.071$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.0342P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
1754 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
123 parameters	$\Delta\rho_{\text{max}} = 0.71 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$
0 constraints	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.5000	0.07163 (7)	0.2500	0.04380 (15)
Cl1	0.58874 (11)	0.2741 (3)	0.03745 (5)	0.1169 (6)
F1	0.29621 (15)	0.3089 (6)	0.24062 (11)	0.0831 (8)
O1	0.52467 (15)	0.3929 (5)	0.19515 (10)	0.0517 (7)
O2	0.55526 (17)	0.0310 (5)	0.15017 (12)	0.0602 (7)
C1	0.5493 (2)	0.2692 (9)	0.15046 (17)	0.0542 (10)
C2	0.5669 (4)	0.4449 (10)	0.0999 (2)	0.098 (2)
H2A	0.6101	0.5583	0.1120	0.117*
H2B	0.5227	0.5579	0.0903	0.117*
C3	0.3963 (2)	-0.0939 (7)	0.21189 (17)	0.0522 (10)
H3A	0.4080	-0.2616	0.1937	0.063*
H3B	0.3620	-0.1296	0.2428	0.063*
C4	0.3557 (2)	0.0799 (8)	0.16691 (16)	0.0465 (9)
C5	0.3052 (2)	0.2743 (9)	0.18211 (18)	0.0537 (10)
C6	0.2662 (3)	0.4379 (9)	0.1434 (2)	0.0703 (13)
H6	0.2318	0.5635	0.1561	0.084*
C7	0.2789 (3)	0.4122 (11)	0.0846 (2)	0.0820 (15)
H7	0.2537	0.5231	0.0570	0.098*
C8	0.3286 (3)	0.2242 (12)	0.0670 (2)	0.0863 (16)
H8	0.3370	0.2078	0.0273	0.104*
C9	0.3667 (3)	0.0579 (10)	0.10703 (19)	0.0683 (12)
H9	0.4000	-0.0703	0.0940	0.082*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.0453 (2)	0.0360 (2)	0.0496 (2)	0.000	-0.00011 (16)	0.000
Cl1	0.1927 (17)	0.1008 (12)	0.0620 (8)	0.0479 (12)	0.0451 (10)	0.0020 (8)
F1	0.0875 (19)	0.084 (2)	0.0804 (18)	0.0152 (16)	0.0270 (15)	0.0010 (15)
O1	0.0685 (18)	0.0384 (16)	0.0496 (15)	0.0031 (13)	0.0147 (13)	0.0003 (12)
O2	0.078 (2)	0.0402 (17)	0.0635 (17)	0.0078 (14)	0.0121 (15)	0.0013 (13)
C1	0.065 (3)	0.046 (3)	0.052 (2)	0.003 (2)	0.010 (2)	0.001 (2)
C2	0.171 (6)	0.058 (3)	0.071 (3)	0.017 (4)	0.057 (4)	0.005 (3)
C3	0.049 (2)	0.039 (2)	0.068 (3)	-0.0047 (19)	-0.0033 (19)	-0.004 (2)
C4	0.043 (2)	0.043 (2)	0.053 (2)	-0.0078 (19)	-0.0053 (17)	-0.0040 (19)
C5	0.047 (2)	0.057 (3)	0.057 (3)	-0.003 (2)	0.003 (2)	-0.003 (2)
C6	0.058 (3)	0.062 (3)	0.089 (4)	0.014 (2)	-0.008 (2)	-0.002 (3)
C7	0.085 (4)	0.072 (4)	0.084 (4)	0.003 (3)	-0.032 (3)	0.012 (3)
C8	0.105 (4)	0.098 (4)	0.052 (3)	0.002 (4)	-0.022 (3)	-0.003 (3)
C9	0.071 (3)	0.072 (3)	0.060 (3)	0.008 (3)	-0.008 (2)	-0.016 (2)

Geometric parameters (\AA , $^\circ$)


Sn1—O1 ⁱ	2.109 (2)	C3—C4	1.492 (5)
---------------------	-----------	-------	-----------

supplementary materials

Sn1—O1	2.109 (2)	C3—H3A	0.9700
Sn1—C3	2.121 (4)	C3—H3B	0.9700
Sn1—C3 ⁱ	2.121 (4)	C4—C5	1.375 (5)
Sn1—O2 ⁱ	2.537 (3)	C4—C9	1.396 (6)
Sn1—O2	2.537 (3)	C5—C6	1.359 (6)
Cl1—C2	1.728 (5)	C6—C7	1.380 (7)
F1—C5	1.365 (4)	C6—H6	0.9300
O1—C1	1.292 (4)	C7—C8	1.361 (7)
O2—C1	1.207 (4)	C7—H7	0.9300
C1—C2	1.503 (6)	C8—C9	1.377 (7)
C2—H2A	0.9700	C8—H8	0.9300
C2—H2B	0.9700	C9—H9	0.9300
O1 ⁱ —Sn1—O1	79.50 (13)	H2A—C2—H2B	107.7
O1 ⁱ —Sn1—C3	110.21 (13)	C4—C3—Sn1	113.7 (3)
O1—Sn1—C3	105.08 (12)	C4—C3—H3A	108.8
O1 ⁱ —Sn1—C3 ⁱ	105.08 (12)	Sn1—C3—H3A	108.8
O1—Sn1—C3 ⁱ	110.21 (13)	C4—C3—H3B	108.8
C3—Sn1—C3 ⁱ	133.6 (2)	Sn1—C3—H3B	108.8
O1 ⁱ —Sn1—O2 ⁱ	54.99 (9)	H3A—C3—H3B	107.7
O1—Sn1—O2 ⁱ	134.28 (9)	C5—C4—C9	115.7 (4)
C3—Sn1—O2 ⁱ	88.52 (13)	C5—C4—C3	121.8 (3)
C3 ⁱ —Sn1—O2 ⁱ	87.82 (13)	C9—C4—C3	122.5 (4)
O1 ⁱ —Sn1—O2	134.28 (9)	C6—C5—F1	118.2 (4)
O1—Sn1—O2	54.99 (9)	C6—C5—C4	124.7 (4)
C3—Sn1—O2	87.82 (13)	F1—C5—C4	117.0 (4)
C3 ⁱ —Sn1—O2	88.52 (13)	C5—C6—C7	118.0 (4)
O2 ⁱ —Sn1—O2	170.72 (12)	C5—C6—H6	121.0
C1—O1—Sn1	100.8 (2)	C7—C6—H6	121.0
C1—O2—Sn1	82.9 (2)	C8—C7—C6	119.8 (5)
O2—C1—O1	121.3 (4)	C8—C7—H7	120.1
O2—C1—C2	124.2 (4)	C6—C7—H7	120.1
O1—C1—C2	114.5 (4)	C7—C8—C9	121.1 (5)
C1—C2—Cl1	113.9 (3)	C7—C8—H8	119.5
C1—C2—H2A	108.8	C9—C8—H8	119.5
Cl1—C2—H2A	108.8	C8—C9—C4	120.6 (4)
C1—C2—H2B	108.8	C8—C9—H9	119.7
Cl1—C2—H2B	108.8	C4—C9—H9	119.7

Symmetry codes: (i) $-x+1, y, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

