

2-[(1*RS*,3*RS*,3*aRS*,6*aSR*)-5-Benzyl-4,6-dioxo-3-phenyloctahydropyrrolo[3,4-*c*]-pyrrol-1-yl]acetamideKonstantin V. Kudryavtsev,^{a*} Andrei V. Churakov^b and Ozdemir Dogan^c^aDepartment of Chemistry, M.V. Lomonosov Moscow State University, Leninskie Gory 1/3, Moscow 119991, Russian Federation, ^bInstitute of General and Inorganic Chemistry, Russian Academy of Sciences, Leninskii prosp. 31, Moscow 119991, Russian Federation, and ^cDepartment of Chemistry, Middle East Technical University, Ankara 06531, Turkey

Correspondence e-mail: kudr@org.chem.msu.ru


Received 23 October 2011; accepted 27 October 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.035; wR factor = 0.108; data-to-parameter ratio = 10.8.

In the title compound, $\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_3$, the relative stereochemistry of the four stereogenic C atoms has been determined. The dihedral angle between the phenyl rings is $77.63(7)^\circ$. In the crystal, ribbons spread along the a axis are formed by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. $\text{C}-\text{H}\cdots\pi$ interactions also occur.

Related literature

For general background to chemistry affording polycyclic pyrrolidine-based scaffolds, see: Kudryavtsev & Irkha (2005); Kudryavtsev (2008, 2011).



Experimental

Crystal data

 $\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_3$ $M_r = 363.41$

Triclinic, $P\bar{1}$
 $a = 9.101(5)$ Å
 $b = 9.270(5)$ Å
 $c = 12.945(5)$ Å
 $\alpha = 103.73(4)^\circ$
 $\beta = 92.30(4)^\circ$
 $\gamma = 113.94(4)^\circ$

$V = 958.1(8)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 293$ K
 $0.50 \times 0.40 \times 0.30$ mm

Data collection

Enraf-Nonius CAD-4
 diffractometer
 4399 measured reflections
 3561 independent reflections

2309 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.011$
 2 standard reflections every 120 min
 intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.108$
 $S = 1.02$
 3561 reflections

329 parameters
 All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.16$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C10–C15 phenyl ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N3}-\text{H32}\cdots\text{O1}^{\text{i}}$	0.91 (2)	2.194 (19)	3.003 (2)	147.3 (16)
$\text{N3}-\text{H31}\cdots\text{O3}^{\text{ii}}$	0.90 (2)	2.00 (2)	2.903 (2)	175.7 (18)
$\text{C18}-\text{H18}\cdots\text{Cg1}^{\text{iii}}$	0.93 (2)	2.71	3.627	166.5

Symmetry codes: (i) $-x, -y + 2, -z$; (ii) $-x + 1, -y + 2, -z$; (iii) $-x, -y + 2, -z + 1$.

Data collection: *CAD4* (Schagen *et al.*, 1988); cell refinement: *CAD4*; data reduction: *XCAD4* (Harms, 1997); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This study was partially supported by the Russian Foundation for Basic Research (project Nos. 11-03-00630 – a and 11-03-91375-ST_a) and State Contract No. 11.519.11.2032.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FF2039).

References

- Harms, K. (1997). *XCAD4*. University of Marburg, Germany.
 Kudryavtsev, K. V. (2008). *Russ. Chem. Bull.* **57**, 2364–2372.
 Kudryavtsev, K. V. (2011). *Heterocycles*, **83**, 323–330.
 Kudryavtsev, K. V. & Irkha, V. V. (2005). *Molecules*, **10**, 755–761.
 Schagen, J. D., Strauer, L., van Meurs, F. & Williams, G. (1988). *CAD4*. Enraf-Nonius, Delft, The Netherlands.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2011). E67, o3186 [doi:10.1107/S1600536811045181]

2-[(1*RS*,3*RS*,3*aRS*,6*aSR*)-5-Benzyl-4,6-dioxo-3-phenyloctahydropyrrolo[3,4-*c*]pyrrol-1-yl]acetamide

K. V. Kudryavtsev, A. V. Churakov and O. Dogan

Comment

The adjacent molecules are combined into ribbons along *a*-axis by hydrogen bonds between neighbouring amide groups (Fig. 2). These ribbons are linked by T-shaped C—H \cdots π interactions between phenyl substituents. Of interest, secondary amine hydrogen atom *NH* does not participate in hydrogen bonding.

Experimental

1.134 g (10.69 mmol) of benzaldehyde, 1.410 g (10.68 mmol) of *L*-asparagine and 2.000 g (10.68 mmol) of *N*-benzyl-maleimide were stirred at 140°C in 30 ml of DMF under an inert atmosphere during 4 h. After cooling to room temperature the reaction mixture was concentrated on rotary evaporator. The residue was chromatographed on silica gel 60 (particle size 0.040–0.063 mm) using CHCl₃—MeOH (40: 1) as eluent. 2-[(1*RS*,3*RS*,3*aRS*,6*aSR*)-5-Benzyl-4,6-dioxo-3-phenyloctahydropyrrolo[3,4-*c*]pyrrol-1-yl]acetamide. Yield 22%, colorless crystals, mp 148–149°C. ¹H NMR (DMSO-*d*₆, δ , J/Hz): 2.44 (dd, 1H, *J* = 15.0, 8.8); 2.58 (dd, 1H, *J* = 15.0, 3.5); 3.23 (dd, 1H, *J* = 8.8, 7.5); 3.42 (dt, 2H, *J* = 8.1, 3.5); 4.15 (d, 1H, *J* = 7.5); 4.57 (s, 2H); 6.93 (s, 1H); 7.25–7.31(m, 4H); 7.32–7.40 (m, 5H); 7.46 (br.s, 1H); 7.47–7.51 (m, 2H). Found (%): C, 69.34; H, 5.85; N, 11.78. C₂₁H₂₁N₃O₃. Calculated (%): C, 69.41; H, 5.82; N, 11.56. The crystals for X-ray analysis were obtained by slow evaporation of saturated chloroform solution at ambient temperature.

Refinement

All hydrogen atoms were located in a difference Fourier map and refined with isotropic thermal parameters.

Figures




Fig. 1. The molecular structure of the title compound, showing the numbering scheme adopted. Displacement ellipsoids are shown at the 50% probability level.




Fig. 2. Hydrogen-bonded ribbons along *a*-axis in the structure of the title compound.

2-[(1*RS*,3*RS*,3*aRS*,6*aSR*)-5-Benzyl-4,6-dioxo-3-phenyloctahydropyrrolo[3,4-*c*]pyrrol-1-yl]acetamide

Crystal data

$C_{21}H_{21}N_3O_3$	$Z = 2$
$M_r = 363.41$	$F(000) = 384$
Triclinic, $P\bar{1}$	$D_x = 1.260 \text{ Mg m}^{-3}$
Hall symbol: $-P\ 1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.101 (5) \text{ \AA}$	Cell parameters from 25 reflections
$b = 9.270 (5) \text{ \AA}$	$\theta = 12\text{--}13^\circ$
$c = 12.945 (5) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\alpha = 103.73 (4)^\circ$	$T = 293 \text{ K}$
$\beta = 92.30 (4)^\circ$	Block, colourless
$\gamma = 113.94 (4)^\circ$	$0.50 \times 0.40 \times 0.30 \text{ mm}$
$V = 958.1 (8) \text{ \AA}^3$	

Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.011$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 2.5^\circ$
graphite	$h = -10 \rightarrow 10$
ω scans	$k = -11 \rightarrow 11$
4399 measured reflections	$l = -2 \rightarrow 15$
3561 independent reflections	2 standard reflections every 120 min
2309 reflections with $I > 2\sigma(I)$	intensity decay: none

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.035$	All H-atom parameters refined
$wR(F^2) = 0.108$	$w = 1/[\sigma^2(F_o^2) + (0.0636P)^2 + 0.0755P]$
$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
3561 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
329 parameters	$\Delta\rho_{\text{max}} = 0.23 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.16 \text{ e \AA}^{-3}$
	Extinction correction: <i>SHELXTL</i> (Sheldrick, 2008),
	$F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.019 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	-0.24934 (15)	0.58343 (15)	0.22990 (10)	0.0479 (3)
N2	0.22129 (15)	0.92640 (17)	0.28729 (10)	0.0483 (3)
N3	0.32943 (19)	1.0559 (2)	-0.00733 (12)	0.0609 (4)
O1	-0.27577 (14)	0.69214 (15)	0.09260 (9)	0.0641 (3)
O2	-0.16268 (15)	0.53957 (17)	0.38310 (11)	0.0802 (4)
O3	0.36090 (17)	0.8890 (2)	0.08216 (11)	0.0822 (5)
C1	0.15214 (17)	0.83233 (18)	0.36351 (11)	0.0422 (3)
C2	-0.02806 (17)	0.80608 (18)	0.34669 (11)	0.0425 (3)
C3	-0.15032 (18)	0.6298 (2)	0.32723 (12)	0.0504 (4)
C4	-0.20472 (17)	0.70517 (18)	0.17794 (11)	0.0441 (3)
C5	-0.05359 (16)	0.84700 (18)	0.24194 (11)	0.0399 (3)
C6	0.10000 (17)	0.85783 (19)	0.19148 (11)	0.0419 (3)
C7	0.13948 (19)	0.9617 (2)	0.11336 (14)	0.0480 (4)
C8	0.28660 (18)	0.9660 (2)	0.06133 (12)	0.0504 (4)
C9	-0.3871 (2)	0.4207 (2)	0.18643 (16)	0.0583 (4)
C10	-0.54238 (19)	0.41544 (18)	0.22548 (13)	0.0510 (4)
C11	-0.6492 (2)	0.4469 (2)	0.16611 (15)	0.0647 (5)
C12	-0.7917 (3)	0.4421 (3)	0.2033 (2)	0.0843 (7)
C13	-0.8263 (3)	0.4063 (3)	0.2991 (2)	0.0888 (7)
C14	-0.7207 (3)	0.3752 (3)	0.3577 (2)	0.0867 (7)
C15	-0.5794 (3)	0.3789 (2)	0.32126 (16)	0.0691 (5)
C16	0.23862 (16)	0.91778 (18)	0.47794 (11)	0.0410 (3)
C17	0.34576 (19)	1.0818 (2)	0.50956 (13)	0.0510 (4)
C18	0.4222 (2)	1.1578 (2)	0.61503 (15)	0.0604 (4)
C19	0.3912 (2)	1.0709 (3)	0.69041 (14)	0.0607 (5)
C20	0.2857 (2)	0.9079 (3)	0.65957 (14)	0.0651 (5)
C21	0.2094 (2)	0.8303 (2)	0.55396 (13)	0.0559 (4)
H1	0.1496 (17)	0.7255 (19)	0.3441 (11)	0.043 (4)*
H6	0.0843 (17)	0.7458 (19)	0.1533 (11)	0.045 (4)*
H72	0.045 (2)	0.917 (2)	0.0574 (14)	0.060 (5)*
H5	-0.0619 (17)	0.9502 (18)	0.2500 (11)	0.041 (4)*

supplementary materials

H22	-0.0542 (19)	0.874 (2)	0.4050 (13)	0.054 (4)*
H17	0.369 (2)	1.143 (2)	0.4586 (14)	0.062 (5)*
H71	0.158 (2)	1.071 (2)	0.1478 (14)	0.063 (5)*
H21	0.135 (2)	0.718 (2)	0.5356 (14)	0.068 (5)*
H92	-0.356 (2)	0.339 (2)	0.2133 (14)	0.070 (5)*
H18	0.497 (2)	1.268 (2)	0.6354 (15)	0.075 (6)*
H91	-0.397 (2)	0.400 (2)	0.1100 (16)	0.072 (6)*
H2	0.308 (2)	0.920 (2)	0.2746 (13)	0.060 (5)*
H19	0.450 (2)	1.126 (2)	0.7648 (17)	0.083 (6)*
H32	0.275 (2)	1.114 (2)	-0.0222 (15)	0.074 (6)*
H11	-0.622 (2)	0.479 (2)	0.1011 (16)	0.076 (6)*
H15	-0.502 (3)	0.348 (3)	0.3604 (17)	0.099 (7)*
H20	0.262 (2)	0.843 (2)	0.7129 (17)	0.085 (6)*
H31	0.423 (3)	1.069 (2)	-0.0340 (16)	0.082 (6)*
H14	-0.748 (3)	0.347 (3)	0.4305 (19)	0.111 (8)*
H12	-0.858 (3)	0.462 (3)	0.1610 (19)	0.103 (8)*
H13	-0.926 (3)	0.398 (3)	0.323 (2)	0.111 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0463 (7)	0.0459 (7)	0.0468 (7)	0.0128 (6)	0.0111 (6)	0.0166 (6)
N2	0.0359 (7)	0.0718 (9)	0.0450 (7)	0.0238 (6)	0.0155 (5)	0.0272 (6)
N3	0.0601 (9)	0.0864 (11)	0.0618 (9)	0.0404 (8)	0.0319 (7)	0.0461 (8)
O1	0.0597 (7)	0.0739 (8)	0.0508 (7)	0.0174 (6)	0.0008 (5)	0.0249 (6)
O2	0.0680 (8)	0.0850 (9)	0.0834 (9)	0.0099 (7)	0.0072 (7)	0.0579 (8)
O3	0.0839 (9)	0.1354 (12)	0.0919 (10)	0.0783 (9)	0.0596 (8)	0.0819 (9)
C1	0.0432 (8)	0.0473 (9)	0.0417 (8)	0.0220 (7)	0.0146 (6)	0.0164 (7)
C2	0.0403 (8)	0.0511 (8)	0.0374 (8)	0.0194 (7)	0.0138 (6)	0.0138 (7)
C3	0.0455 (8)	0.0585 (9)	0.0492 (9)	0.0169 (7)	0.0150 (7)	0.0268 (8)
C4	0.0425 (8)	0.0514 (8)	0.0413 (8)	0.0204 (7)	0.0123 (6)	0.0166 (7)
C5	0.0391 (7)	0.0436 (8)	0.0435 (8)	0.0205 (6)	0.0150 (6)	0.0172 (6)
C6	0.0415 (8)	0.0505 (9)	0.0424 (8)	0.0240 (7)	0.0171 (6)	0.0191 (7)
C7	0.0434 (9)	0.0611 (10)	0.0488 (9)	0.0246 (8)	0.0156 (7)	0.0261 (8)
C8	0.0497 (9)	0.0702 (10)	0.0435 (8)	0.0296 (8)	0.0173 (7)	0.0288 (8)
C9	0.0572 (10)	0.0448 (9)	0.0604 (11)	0.0109 (8)	0.0126 (8)	0.0115 (8)
C10	0.0520 (9)	0.0370 (8)	0.0525 (9)	0.0090 (7)	0.0098 (7)	0.0101 (7)
C11	0.0676 (11)	0.0618 (11)	0.0554 (10)	0.0225 (9)	0.0022 (9)	0.0105 (9)
C12	0.0716 (14)	0.0691 (13)	0.1083 (18)	0.0337 (11)	-0.0010 (13)	0.0137 (12)
C13	0.0777 (15)	0.0622 (12)	0.130 (2)	0.0300 (11)	0.0492 (15)	0.0267 (13)
C14	0.0970 (17)	0.0768 (14)	0.0990 (17)	0.0351 (13)	0.0525 (14)	0.0445 (13)
C15	0.0751 (12)	0.0611 (11)	0.0751 (13)	0.0222 (9)	0.0264 (10)	0.0363 (10)
C16	0.0377 (7)	0.0511 (8)	0.0428 (8)	0.0245 (6)	0.0125 (6)	0.0170 (6)
C17	0.0521 (9)	0.0527 (9)	0.0531 (9)	0.0243 (8)	0.0122 (7)	0.0192 (8)
C18	0.0535 (10)	0.0590 (11)	0.0627 (11)	0.0251 (9)	0.0055 (8)	0.0048 (9)
C19	0.0530 (10)	0.0871 (14)	0.0468 (9)	0.0407 (10)	0.0049 (8)	0.0080 (9)
C20	0.0699 (12)	0.0902 (14)	0.0494 (10)	0.0406 (11)	0.0139 (9)	0.0321 (10)
C21	0.0582 (10)	0.0600 (11)	0.0508 (9)	0.0211 (9)	0.0101 (8)	0.0250 (8)

Geometric parameters (Å, °)

N1—C4	1.379 (2)	C9—C10	1.506 (2)
N1—C3	1.386 (2)	C9—H92	1.030 (18)
N1—C9	1.473 (2)	C9—H91	0.955 (19)
N2—C6	1.452 (2)	C10—C11	1.377 (3)
N2—C1	1.4639 (19)	C10—C15	1.381 (2)
N2—H2	0.832 (18)	C11—C12	1.389 (3)
N3—C8	1.324 (2)	C11—H11	0.97 (2)
N3—H32	0.91 (2)	C12—C13	1.374 (4)
N3—H31	0.90 (2)	C12—H12	0.90 (2)
O1—C4	1.2120 (18)	C13—C14	1.361 (4)
O2—C3	1.2053 (19)	C13—H13	0.95 (3)
O3—C8	1.2281 (19)	C14—C15	1.378 (3)
C1—C16	1.511 (2)	C14—H14	1.05 (2)
C1—C2	1.555 (2)	C15—H15	1.01 (2)
C1—H1	0.953 (15)	C16—C17	1.378 (2)
C2—C3	1.509 (2)	C16—C21	1.386 (2)
C2—C5	1.525 (2)	C17—C18	1.384 (3)
C2—H22	0.966 (16)	C17—H17	0.943 (18)
C4—C5	1.494 (2)	C18—C19	1.375 (3)
C5—C6	1.543 (2)	C18—H18	0.93 (2)
C5—H5	0.972 (14)	C19—C20	1.368 (3)
C6—C7	1.513 (2)	C19—H19	0.99 (2)
C6—H6	0.987 (15)	C20—C21	1.386 (3)
C7—C8	1.513 (2)	C20—H20	1.00 (2)
C7—H72	0.978 (18)	C21—H21	0.944 (19)
C7—H71	0.944 (19)		
C4—N1—C3	112.94 (13)	O3—C8—N3	122.40 (15)
C4—N1—C9	123.65 (14)	O3—C8—C7	121.10 (13)
C3—N1—C9	123.41 (14)	N3—C8—C7	116.50 (14)
C6—N2—C1	107.05 (12)	N1—C9—C10	112.20 (14)
C6—N2—H2	112.4 (12)	N1—C9—H92	106.3 (10)
C1—N2—H2	109.2 (12)	C10—C9—H92	109.0 (10)
C8—N3—H32	122.6 (12)	N1—C9—H91	105.6 (11)
C8—N3—H31	118.2 (12)	C10—C9—H91	111.2 (11)
H32—N3—H31	118.8 (17)	H92—C9—H91	112.5 (15)
N2—C1—C16	113.54 (13)	C11—C10—C15	119.19 (18)
N2—C1—C2	101.61 (11)	C11—C10—C9	120.46 (16)
C16—C1—C2	114.38 (12)	C15—C10—C9	120.35 (17)
N2—C1—H1	112.9 (9)	C10—C11—C12	119.8 (2)
C16—C1—H1	108.1 (8)	C10—C11—H11	120.2 (12)
C2—C1—H1	106.1 (9)	C12—C11—H11	119.9 (12)
C3—C2—C5	104.30 (13)	C13—C12—C11	120.3 (2)
C3—C2—C1	114.00 (13)	C13—C12—H12	123.5 (16)
C5—C2—C1	106.24 (11)	C11—C12—H12	116.2 (16)
C3—C2—H22	108.6 (9)	C14—C13—C12	119.9 (2)
C5—C2—H22	110.1 (9)	C14—C13—H13	120.0 (15)

supplementary materials

C1—C2—H22	113.2 (9)	C12—C13—H13	120.1 (15)
O2—C3—N1	123.89 (15)	C13—C14—C15	120.3 (2)
O2—C3—C2	127.97 (15)	C13—C14—H14	119.8 (13)
N1—C3—C2	108.13 (13)	C15—C14—H14	119.9 (13)
O1—C4—N1	124.11 (15)	C14—C15—C10	120.5 (2)
O1—C4—C5	127.43 (14)	C14—C15—H15	122.1 (12)
N1—C4—C5	108.40 (13)	C10—C15—H15	117.3 (12)
C4—C5—C2	105.30 (12)	C17—C16—C21	118.60 (15)
C4—C5—C6	112.59 (12)	C17—C16—C1	121.77 (14)
C2—C5—C6	103.62 (11)	C21—C16—C1	119.62 (14)
C4—C5—H5	111.1 (8)	C16—C17—C18	120.74 (17)
C2—C5—H5	115.4 (8)	C16—C17—H17	119.8 (11)
C6—C5—H5	108.6 (8)	C18—C17—H17	119.4 (11)
N2—C6—C7	113.43 (13)	C19—C18—C17	120.39 (18)
N2—C6—C5	100.79 (11)	C19—C18—H18	119.3 (12)
C7—C6—C5	113.69 (12)	C17—C18—H18	120.3 (12)
N2—C6—H6	111.2 (8)	C20—C19—C18	119.24 (17)
C7—C6—H6	108.8 (8)	C20—C19—H19	121.2 (12)
C5—C6—H6	108.7 (8)	C18—C19—H19	119.5 (12)
C6—C7—C8	113.10 (13)	C19—C20—C21	120.81 (18)
C6—C7—H72	108.4 (9)	C19—C20—H20	120.4 (12)
C8—C7—H72	109.2 (10)	C21—C20—H20	118.8 (12)
C6—C7—H71	111.0 (10)	C20—C21—C16	120.22 (18)
C8—C7—H71	108.4 (11)	C20—C21—H21	118.8 (11)
H72—C7—H71	106.6 (14)	C16—C21—H21	121.0 (11)
C6—N2—C1—C16	-162.41 (12)	C4—C5—C6—C7	90.44 (16)
C6—N2—C1—C2	-39.10 (14)	C2—C5—C6—C7	-156.31 (13)
N2—C1—C2—C3	129.33 (13)	N2—C6—C7—C8	67.80 (18)
C16—C1—C2—C3	-107.94 (15)	C5—C6—C7—C8	-177.81 (13)
N2—C1—C2—C5	15.04 (14)	C6—C7—C8—O3	0.9 (2)
C16—C1—C2—C5	137.77 (13)	C6—C7—C8—N3	-179.38 (15)
C4—N1—C3—O2	-176.19 (15)	C4—N1—C9—C10	-92.44 (19)
C9—N1—C3—O2	3.4 (2)	C3—N1—C9—C10	87.97 (19)
C4—N1—C3—C2	2.69 (17)	N1—C9—C10—C11	92.8 (2)
C9—N1—C3—C2	-177.68 (13)	N1—C9—C10—C15	-87.3 (2)
C5—C2—C3—O2	171.11 (16)	C15—C10—C11—C12	0.2 (3)
C1—C2—C3—O2	55.7 (2)	C9—C10—C11—C12	-179.83 (16)
C5—C2—C3—N1	-7.71 (15)	C10—C11—C12—C13	0.1 (3)
C1—C2—C3—N1	-123.14 (13)	C11—C12—C13—C14	-0.1 (3)
C3—N1—C4—O1	-178.84 (14)	C12—C13—C14—C15	-0.2 (4)
C9—N1—C4—O1	1.5 (2)	C13—C14—C15—C10	0.5 (3)
C3—N1—C4—C5	3.74 (16)	C11—C10—C15—C14	-0.5 (3)
C9—N1—C4—C5	-175.89 (13)	C9—C10—C15—C14	179.51 (17)
O1—C4—C5—C2	174.29 (15)	N2—C1—C16—C17	13.95 (19)
N1—C4—C5—C2	-8.41 (15)	C2—C1—C16—C17	-102.06 (16)
O1—C4—C5—C6	-73.49 (19)	N2—C1—C16—C21	-166.52 (13)
N1—C4—C5—C6	103.82 (14)	C2—C1—C16—C21	77.47 (17)
C3—C2—C5—C4	9.55 (14)	C21—C16—C17—C18	-0.2 (2)
C1—C2—C5—C4	130.31 (12)	C1—C16—C17—C18	179.33 (13)

C3—C2—C5—C6	-108.87 (13)	C16—C17—C18—C19	-0.6 (2)
C1—C2—C5—C6	11.89 (14)	C17—C18—C19—C20	0.9 (3)
C1—N2—C6—C7	168.92 (12)	C18—C19—C20—C21	-0.4 (3)
C1—N2—C6—C5	47.03 (14)	C19—C20—C21—C16	-0.4 (3)
C4—C5—C6—N2	-147.85 (12)	C17—C16—C21—C20	0.7 (2)
C2—C5—C6—N2	-34.60 (14)	C1—C16—C21—C20	-178.86 (14)

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C10—C15 phenyl ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N3—H32 \cdots O1 ⁱ	0.91 (2)	2.194 (19)	3.003 (2)	147.3 (16)
N3—H31 \cdots O3 ⁱⁱ	0.90 (2)	2.00 (2)	2.903 (2)	175.7 (18)
C18—H18 \cdots Cg1 ⁱⁱⁱ	0.93 (2)	2.71	3.627	166.5

Symmetry codes: (i) $-x, -y+2, -z$; (ii) $-x+1, -y+2, -z$; (iii) $-x, -y+2, -z+1$.

Fig. 1




Fig. 2

