

(E)-1-(4-Benzhydrylpiperazin-1-yl)-3-(2-ethoxyphenyl)prop-2-en-1-one

 Yan Zhong,^a XiaoPing Zhang^b and Bin Wu^{c*}

^aSchool of Chemistry and Chemical Engineering, Southeast University, Sipailou No. 2 Nanjing, Nanjing 210096, People's Republic of China, ^bCentre of Laboratory Animals, Nanjing Medical University, Hanzhong Road No. 140 Nanjing, Nanjing 210029, People's Republic of China, and ^cSchool of Pharmacy, Nanjing Medical University, Hanzhong Road No. 140 Nanjing, Nanjing 210029, People's Republic of China

Correspondence e-mail: wubin@njmu.edu.cn


Received 31 October 2011; accepted 1 November 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.077; wR factor = 0.155; data-to-parameter ratio = 16.8.

In the title molecule, $\text{C}_{28}\text{H}_{30}\text{N}_2\text{O}_2$, the piperazine ring adopts a chair conformation and the $\text{C}=\text{C}$ bond exhibits an *E* conformation. The dihedral angle between the terminal phenyl rings is 71.4 (2). In the crystal, molecules are linked by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming [010] chains.

Related literature

For properties of cinnamic acid derivatives, see: Shi *et al.* (2005); Qian *et al.* (2010). For the synthesis, see: Wu *et al.* (2008). For related structures, see: Mouillé *et al.* (1975); Zhong *et al.* (2011).



Experimental

Crystal data

$\text{C}_{28}\text{H}_{30}\text{N}_2\text{O}_2$
 $M_r = 426.54$
 Monoclinic, $P2_1/n$

$a = 11.858$ (2) Å
 $b = 12.786$ (3) Å
 $c = 16.044$ (3) Å

$\beta = 94.63$ (3)°
 $V = 2424.6$ (8) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.07$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.20 \times 0.10$ mm

Data collection

Enraf–Nonius CAD-4 diffractometer
 4673 measured reflections
 4446 independent reflections
 2022 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.063$
 3 standard reflections every 200 reflections
 intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.077$
 $wR(F^2) = 0.155$
 $S = 1.00$
 4446 reflections

265 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.13$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.22$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C11}-\text{H11A}\cdots\text{O1}^i$	0.93	2.54	3.343 (6)	146

 Symmetry code: (i) $x, y + 1, z$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1989); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXL97* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank Professor Hua-Qin Wang of the Analysis Centre, Nanjing University, for the diffraction measurements. This work was supported by the Natural Science Foundation of the Education Department of Jiangsu Province (No. 05KJB350084) and the Natural Science Foundation of Jiangsu Province (No. BK2010538).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6487).

References

- Enraf–Nonius (1989). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
 Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
 Mouillé, Y., Cotrait, M., Hospital, M. & Marsau, P. (1975). *Acta Cryst.* **B31**, 1495–1496.
 Qian, Y., Zhang, H.-J., Zhang, H., Xu, J. & Zhu, H.-L. (2010). *Bioorg. Med. Chem.* **18**, 4991–4996.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Shi, Y., Chen, Q.-X., Wang, Q., Song, K.-K. & Qiu, L. (2005). *Food Chem.* **92**, 707–712.
 Wu, B., Zhou, L. & Cai, H.-H. (2008). *Chin. Chem. Lett.* **19**, 1163–1166.
 Zhong, Y. & Wu, B. (2011). *Acta Cryst.* **E67**, o2992.

supplementary materials

Acta Cryst. (2011). E67, o3200 [doi:10.1107/S1600536811045922]

(*E*)-1-(4-Benzhydrylpiperazin-1-yl)-3-(2-ethoxyphenyl)prop-2-en-1-one

Y. Zhong, X. P. Zhang and B. Wu

Comment

Many compounds containing a cinnamoyl moiety have been reported to possess antimicrobial, anticancer and neuroprotective activities (Shi *et al.*, 2005; Qian *et al.*, 2010). We report herein the crystal structure of the title compound. The molecule of the title compound exists an *E* configuration with respect to the C19=C20 ethene bond [1.291 (4)] and the torsion angle C18—C19—C20—C21 = 177.2 (3). The piperazine ring adopts a chair conformation. In the crystal, molecules are linked by C—H \cdots O interactions.

Experimental

The synthesis follows the method of Wu *et al.* (2008). The title compound was prepared by stirring a mixture of (*E*)-3-(2-ethoxyphenyl)acrylic acid (0.769 g; 4 mmol), dimethyl sulfoxide (2 ml) and dichloromethane (30 ml) for 6 h at room temperature. The solvent was removed under reduced pressure. The residue was dissolved in acetone (15 ml) and reacted with 1-benzhydrylpiperazine (1.514 g; 6 mmol) in the presence of triethylamine (5 ml) for 12 h at room temperature. The resultant mixture was cooled. The solid, (*E*)-1-(4-benzhydrylpiperazin-1-yl)-3-(2-ethoxyphenyl)prop-2-en-1-one obtained was filtered and was recrystallized from ethanol. Pale-yellow blocks were grown in ethyl acetate and hexane by a slow evaporation at room temperature.

Refinement

All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were positioned geometrically with C—H distances ranging from 0.93 Å to 0.98 Å and refined as riding on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}$ of the carrier atom.

Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids for non-H atoms drawn at the 70% probability level.




Fig. 2. Packing diagram of the title compound.

(E)-1-(4-Benzhydrylpiperazin-1-yl)-3-(2-ethoxyphenyl)prop-2-en-1-one

Crystal data

$C_{28}H_{30}N_2O_2$

$M_r = 426.54$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 11.858 (2) \text{ \AA}$

$b = 12.786 (3) \text{ \AA}$

$c = 16.044 (3) \text{ \AA}$

$\beta = 94.63 (3)^\circ$

$V = 2424.6 (8) \text{ \AA}^3$

$Z = 4$

$F(000) = 912$

$D_x = 1.169 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 25 reflections

$\theta = 9\text{--}12^\circ$

$\mu = 0.07 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, pale-yellow

$0.30 \times 0.20 \times 0.10 \text{ mm}$

Data collection

Enraf-Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube
graphite

$\omega/2\theta$ scans

4673 measured reflections

4446 independent reflections

2022 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.063$

$\theta_{\text{max}} = 25.4^\circ$, $\theta_{\text{min}} = 2.0^\circ$

$h = 0 \rightarrow 14$

$k = 0 \rightarrow 15$

$l = -19 \rightarrow 19$

3 standard reflections every 200 reflections

intensity decay: 1%

Refinement

Refinement on F^2

Least-squares matrix: full

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

$$R[F^2 > 2\sigma(F^2)] = 0.077$$

$$wR(F^2) = 0.155$$

$$S = 1.00$$

4446 reflections

265 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.030P)^2 + 1.550P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.13 \text{ e } \text{Å}^{-3}$$

$$\Delta\rho_{\min} = -0.22 \text{ e } \text{Å}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.2436 (2)	0.32431 (17)	0.49193 (13)	0.0823 (8)
N1	0.2112 (2)	0.69070 (19)	0.57757 (17)	0.0641 (7)
C1	0.0062 (5)	0.7832 (3)	0.6573 (3)	0.1159 (15)
H1A	-0.0014	0.7739	0.5996	0.139*
C2	-0.0941 (4)	0.7913 (3)	0.6932 (3)	0.103
H2A	-0.1671	0.7873	0.6674	0.124*
C3	-0.0616 (5)	0.8068 (3)	0.7764 (3)	0.1158 (15)
H3A	-0.1233	0.8218	0.8067	0.139*
C4	0.0365 (5)	0.8061 (4)	0.8263 (3)	0.1107 (15)
H4A	0.0396	0.8135	0.8841	0.133*
C5	0.1251 (5)	0.7940 (3)	0.7854 (4)	0.1213 (16)
H5A	0.1962	0.7901	0.8142	0.146*
C6	0.1140 (4)	0.7863 (3)	0.6906 (4)	0.1098 (17)
O2	0.3846 (2)	0.12578 (18)	0.28272 (15)	0.0852 (8)
N2	0.2698 (3)	0.4974 (2)	0.50411 (17)	0.0790 (9)
C7	0.2064 (3)	0.7757 (3)	0.6338 (3)	0.095
H7A	0.2752	0.7684	0.6715	0.114*
C8	0.2259 (4)	0.8738 (3)	0.5848 (3)	0.1061 (14)
C9	0.3187 (3)	0.9292 (3)	0.6198 (2)	0.084
H9A	0.3618	0.9034	0.6663	0.100*
C10	0.3444 (4)	1.0185 (3)	0.5861 (3)	0.097
H10A	0.4150	1.0458	0.6040	0.117*
C11	0.2883 (5)	1.0710 (4)	0.5344 (3)	0.1141 (15)
H11A	0.3101	1.1387	0.5221	0.137*

supplementary materials

C12	0.1862 (5)	1.0252 (5)	0.4921 (3)	0.1197 (17)
H12A	0.1442	1.0597	0.4490	0.144*
C13	0.1550 (4)	0.9251 (5)	0.5206 (3)	0.1267 (17)
H13A	0.0891	0.8930	0.4979	0.152*
C14	0.2828 (3)	0.6824 (3)	0.5164 (3)	0.1075 (15)
H14A	0.3513	0.7181	0.5377	0.129*
H14B	0.2498	0.7249	0.4707	0.129*
C15	0.3189 (4)	0.5913 (3)	0.4794 (3)	0.1085 (15)
H15A	0.4002	0.5859	0.4912	0.130*
H15B	0.3041	0.5981	0.4193	0.130*
C16	0.1949 (4)	0.4975 (3)	0.5695 (3)	0.1160 (16)
H16A	0.1277	0.4580	0.5508	0.139*
H16B	0.2315	0.4614	0.6175	0.139*
C17	0.1613 (4)	0.5974 (3)	0.5953 (3)	0.1131 (16)
H17A	0.0823	0.6043	0.5749	0.136*
H17B	0.1627	0.5941	0.6557	0.136*
C18	0.2861 (3)	0.4023 (3)	0.4683 (2)	0.0757 (11)
C19	0.3575 (3)	0.4008 (2)	0.39672 (19)	0.0670 (9)
H19A	0.3894	0.4628	0.3797	0.080*
C20	0.3765 (3)	0.3152 (2)	0.35741 (18)	0.0720 (10)
H20A	0.3453	0.2550	0.3786	0.086*
C21	0.4405 (3)	0.3001 (3)	0.28366 (19)	0.0637 (9)
C22	0.4967 (3)	0.3810 (3)	0.2482 (2)	0.0733 (10)
H22A	0.4923	0.4476	0.2711	0.088*
C23	0.5596 (3)	0.3673 (3)	0.1799 (2)	0.0918 (13)
H23A	0.5981	0.4232	0.1584	0.110*
C24	0.5641 (3)	0.2701 (3)	0.1446 (2)	0.0873 (12)
H24A	0.6081	0.2590	0.1000	0.105*
C25	0.5004 (3)	0.1846 (3)	0.1769 (2)	0.0829 (11)
H25A	0.4989	0.1199	0.1504	0.100*
C26	0.4417 (3)	0.1990 (3)	0.2469 (2)	0.0721 (10)
C27	0.3835 (3)	0.0218 (3)	0.2493 (2)	0.0919 (12)
H27A	0.4600	-0.0052	0.2505	0.110*
H27B	0.3516	0.0223	0.1917	0.110*
C28	0.3142 (3)	-0.0455 (3)	0.3005 (3)	0.1021 (13)
H28A	0.3140	-0.1158	0.2794	0.153*
H28B	0.2381	-0.0194	0.2977	0.153*
H28C	0.3457	-0.0449	0.3575	0.153*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.129 (2)	0.0544 (14)	0.0689 (15)	-0.0102 (14)	0.0430 (14)	-0.0047 (12)
N1	0.0595 (18)	0.0530 (16)	0.0809 (19)	0.0014 (14)	0.0131 (15)	-0.0082 (15)
C1	0.119 (4)	0.121 (4)	0.111 (4)	-0.025 (3)	0.028 (3)	0.009 (3)
C2	0.103	0.103	0.103	0.000	0.008	0.000
C3	0.128 (4)	0.102 (3)	0.120 (4)	0.010 (3)	0.024 (3)	-0.005 (3)
C4	0.114 (4)	0.110 (4)	0.110 (4)	0.029 (3)	0.024 (3)	-0.014 (3)

C5	0.112 (4)	0.097 (3)	0.153 (5)	0.016 (3)	0.003 (4)	-0.006 (3)
C6	0.102 (4)	0.077 (3)	0.162 (5)	0.004 (3)	0.080 (4)	-0.032 (3)
O2	0.104 (2)	0.0662 (15)	0.0907 (18)	0.0147 (14)	0.0390 (15)	-0.0212 (14)
N2	0.113 (2)	0.0495 (16)	0.081 (2)	-0.0052 (16)	0.0478 (18)	-0.0007 (15)
C7	0.095	0.095	0.095	0.000	0.008	0.000
C8	0.107 (4)	0.090 (3)	0.125 (4)	-0.007 (3)	0.032 (3)	-0.010 (3)
C9	0.084	0.084	0.084	0.000	0.007	0.000
C10	0.097	0.097	0.097	0.000	0.008	0.000
C11	0.137 (4)	0.088 (3)	0.120 (4)	0.009 (3)	0.029 (3)	0.013 (3)
C12	0.124 (5)	0.146 (5)	0.090 (3)	0.046 (4)	0.011 (3)	-0.005 (4)
C13	0.113 (4)	0.154 (5)	0.114 (4)	-0.005 (4)	0.019 (3)	-0.032 (4)
C14	0.104 (3)	0.066 (2)	0.162 (4)	-0.023 (2)	0.068 (3)	-0.001 (3)
C15	0.153 (4)	0.074 (3)	0.108 (3)	-0.016 (3)	0.067 (3)	-0.019 (3)
C16	0.171 (4)	0.079 (3)	0.110 (3)	-0.009 (3)	0.085 (3)	-0.013 (2)
C17	0.136 (4)	0.072 (3)	0.143 (4)	-0.006 (3)	0.088 (3)	-0.010 (3)
C18	0.098 (3)	0.059 (2)	0.076 (2)	0.006 (2)	0.047 (2)	-0.0040 (19)
C19	0.084 (2)	0.0472 (18)	0.075 (2)	0.0158 (17)	0.0369 (19)	0.0063 (17)
C20	0.114 (3)	0.0499 (19)	0.057 (2)	0.0050 (19)	0.039 (2)	0.0019 (16)
C21	0.064 (2)	0.068 (2)	0.061 (2)	0.0222 (18)	0.0102 (17)	0.0044 (18)
C22	0.094 (3)	0.063 (2)	0.066 (2)	0.023 (2)	0.024 (2)	0.0038 (18)
C23	0.137 (4)	0.080 (3)	0.065 (2)	0.000 (3)	0.046 (2)	0.007 (2)
C24	0.090 (3)	0.112 (3)	0.065 (2)	0.012 (3)	0.034 (2)	0.005 (2)
C25	0.070 (2)	0.101 (3)	0.082 (3)	0.024 (2)	0.030 (2)	-0.011 (2)
C26	0.074 (3)	0.075 (2)	0.069 (2)	0.016 (2)	0.013 (2)	0.003 (2)
C27	0.097 (3)	0.093 (3)	0.087 (3)	0.018 (2)	0.014 (2)	-0.037 (2)
C28	0.097 (3)	0.083 (3)	0.129 (4)	0.014 (2)	0.027 (3)	-0.017 (3)

Geometric parameters (Å, °)

O1—C18	1.194 (4)	C13—H13A	0.9300
N1—C14	1.352 (4)	C14—C15	1.391 (4)
N1—C17	1.372 (4)	C14—H14A	0.9700
N1—C7	1.416 (4)	C14—H14B	0.9700
C1—C6	1.347 (6)	C15—H15A	0.9700
C1—C2	1.367 (6)	C15—H15B	0.9700
C1—H1A	0.9300	C16—C17	1.410 (5)
C2—C3	1.373 (6)	C16—H16A	0.9700
C2—H2A	0.9300	C16—H16B	0.9700
C3—C4	1.357 (6)	C17—H17A	0.9700
C3—H3A	0.9300	C17—H17B	0.9700
C4—C5	1.292 (6)	C18—C19	1.480 (4)
C4—H4A	0.9300	C19—C20	1.291 (4)
C5—C6	1.519 (6)	C19—H19A	0.9300
C5—H5A	0.9300	C20—C21	1.470 (4)
C6—C7	1.487 (5)	C20—H20A	0.9300
O2—C26	1.315 (4)	C21—C22	1.378 (4)
O2—C27	1.434 (4)	C21—C26	1.421 (4)
N2—C18	1.366 (4)	C22—C23	1.385 (4)
N2—C15	1.405 (4)	C22—H22A	0.9300

supplementary materials

N2—C16	1.428 (4)	C23—C24	1.368 (5)
C7—C8	1.508 (6)	C23—H23A	0.9300
C7—H7A	0.9800	C24—C25	1.449 (5)
C8—C9	1.389 (5)	C24—H24A	0.9300
C8—C13	1.435 (6)	C25—C26	1.380 (4)
C9—C10	1.310 (5)	C25—H25A	0.9300
C9—H9A	0.9300	C27—C28	1.482 (5)
C10—C11	1.222 (5)	C27—H27A	0.9700
C10—H10A	0.9300	C27—H27B	0.9700
C11—C12	1.461 (6)	C28—H28A	0.9600
C11—H11A	0.9300	C28—H28B	0.9600
C12—C13	1.419 (6)	C28—H28C	0.9600
C12—H12A	0.9300		
C14—N1—C17	112.8 (3)	C14—C15—H15A	108.2
C14—N1—C7	125.6 (3)	N2—C15—H15A	108.2
C17—N1—C7	119.7 (3)	C14—C15—H15B	108.2
C6—C1—C2	131.4 (5)	N2—C15—H15B	108.2
C6—C1—H1A	114.3	H15A—C15—H15B	107.3
C2—C1—H1A	114.3	C17—C16—N2	115.1 (3)
C1—C2—C3	103.6 (4)	C17—C16—H16A	108.5
C1—C2—H2A	128.2	N2—C16—H16A	108.5
C3—C2—H2A	128.2	C17—C16—H16B	108.5
C4—C3—C2	137.0 (5)	N2—C16—H16B	108.5
C4—C3—H3A	111.5	H16A—C16—H16B	107.5
C2—C3—H3A	111.5	N1—C17—C16	126.2 (3)
C5—C4—C3	113.3 (5)	N1—C17—H17A	105.8
C5—C4—H4A	123.4	C16—C17—H17A	105.8
C3—C4—H4A	123.4	N1—C17—H17B	105.8
C4—C5—C6	120.5 (5)	C16—C17—H17B	105.8
C4—C5—H5A	119.7	H17A—C17—H17B	106.2
C6—C5—H5A	119.7	O1—C18—N2	122.2 (3)
C1—C6—C7	118.5 (5)	O1—C18—C19	121.2 (3)
C1—C6—C5	113.7 (4)	N2—C18—C19	116.6 (3)
C7—C6—C5	127.7 (5)	C20—C19—C18	121.6 (3)
C26—O2—C27	119.1 (3)	C20—C19—H19A	119.2
C18—N2—C15	124.4 (3)	C18—C19—H19A	119.2
C18—N2—C16	115.3 (3)	C19—C20—C21	128.8 (3)
C15—N2—C16	120.2 (3)	C19—C20—H20A	115.6
N1—C7—C6	122.0 (4)	C21—C20—H20A	115.6
N1—C7—C8	106.9 (3)	C22—C21—C26	119.4 (3)
C6—C7—C8	113.3 (4)	C22—C21—C20	122.1 (3)
N1—C7—H7A	104.3	C26—C21—C20	118.5 (3)
C6—C7—H7A	104.3	C21—C22—C23	122.7 (3)
C8—C7—H7A	104.3	C21—C22—H22A	118.7
C9—C8—C13	117.2 (4)	C23—C22—H22A	118.7
C9—C8—C7	111.3 (5)	C24—C23—C22	119.0 (3)
C13—C8—C7	130.4 (5)	C24—C23—H23A	120.5
C10—C9—C8	118.7 (4)	C22—C23—H23A	120.5
C10—C9—H9A	120.6	C23—C24—C25	119.9 (3)

C8—C9—H9A	120.6	C23—C24—H24A	120.1
C11—C10—C9	128.8 (5)	C25—C24—H24A	120.1
C11—C10—H10A	115.6	C26—C25—C24	120.1 (3)
C9—C10—H10A	115.6	C26—C25—H25A	119.9
C10—C11—C12	118.8 (5)	C24—C25—H25A	119.9
C10—C11—H11A	120.6	O2—C26—C25	124.9 (3)
C12—C11—H11A	120.6	O2—C26—C21	116.3 (3)
C13—C12—C11	116.0 (5)	C25—C26—C21	118.8 (3)
C13—C12—H12A	122.0	O2—C27—C28	108.6 (3)
C11—C12—H12A	122.0	O2—C27—H27A	110.0
C12—C13—C8	119.4 (5)	C28—C27—H27A	110.0
C12—C13—H13A	120.3	O2—C27—H27B	110.0
C8—C13—H13A	120.3	C28—C27—H27B	110.0
N1—C14—C15	127.5 (3)	H27A—C27—H27B	108.3
N1—C14—H14A	105.4	C27—C28—H28A	109.5
C15—C14—H14A	105.4	C27—C28—H28B	109.5
N1—C14—H14B	105.4	H28A—C28—H28B	109.5
C15—C14—H14B	105.4	C27—C28—H28C	109.5
H14A—C14—H14B	106.0	H28A—C28—H28C	109.5
C14—C15—N2	116.4 (3)	H28B—C28—H28C	109.5
C6—C1—C2—C3	1.4 (7)	C18—N2—C15—C14	172.9 (4)
C1—C2—C3—C4	-7.1 (8)	C16—N2—C15—C14	-5.5 (6)
C2—C3—C4—C5	5.4 (9)	C18—N2—C16—C17	-168.4 (4)
C3—C4—C5—C6	2.2 (7)	C15—N2—C16—C17	10.1 (6)
C2—C1—C6—C7	-179.6 (4)	C14—N1—C17—C16	13.8 (7)
C2—C1—C6—C5	4.0 (8)	C7—N1—C17—C16	-151.5 (5)
C4—C5—C6—C1	-5.9 (7)	N2—C16—C17—N1	-15.2 (8)
C4—C5—C6—C7	178.0 (4)	C15—N2—C18—O1	179.2 (4)
C14—N1—C7—C6	170.6 (4)	C16—N2—C18—O1	-2.4 (6)
C17—N1—C7—C6	-26.1 (6)	C15—N2—C18—C19	-2.2 (6)
C14—N1—C7—C8	38.0 (5)	C16—N2—C18—C19	176.2 (3)
C17—N1—C7—C8	-158.7 (4)	O1—C18—C19—C20	-0.1 (6)
C1—C6—C7—N1	-53.6 (6)	N2—C18—C19—C20	-178.7 (4)
C5—C6—C7—N1	122.3 (5)	C18—C19—C20—C21	177.2 (3)
C1—C6—C7—C8	76.4 (5)	C19—C20—C21—C22	3.7 (6)
C5—C6—C7—C8	-107.7 (5)	C19—C20—C21—C26	-174.7 (4)
N1—C7—C8—C9	-122.0 (4)	C26—C21—C22—C23	-3.2 (5)
C6—C7—C8—C9	100.8 (5)	C20—C21—C22—C23	178.5 (3)
N1—C7—C8—C13	70.4 (6)	C21—C22—C23—C24	1.7 (6)
C6—C7—C8—C13	-66.7 (6)	C22—C23—C24—C25	2.3 (6)
C13—C8—C9—C10	-8.2 (6)	C23—C24—C25—C26	-4.7 (6)
C7—C8—C9—C10	-177.6 (3)	C27—O2—C26—C25	1.5 (5)
C8—C9—C10—C11	12.5 (7)	C27—O2—C26—C21	-179.3 (3)
C9—C10—C11—C12	-10.9 (7)	C24—C25—C26—O2	-177.6 (3)
C10—C11—C12—C13	5.5 (7)	C24—C25—C26—C21	3.2 (5)
C11—C12—C13—C8	-2.8 (6)	C22—C21—C26—O2	-178.7 (3)
C9—C8—C13—C12	4.3 (6)	C20—C21—C26—O2	-0.2 (5)
C7—C8—C13—C12	171.3 (4)	C22—C21—C26—C25	0.6 (5)
C17—N1—C14—C15	-8.2 (7)	C20—C21—C26—C25	179.1 (3)

supplementary materials

C7—N1—C14—C15	156.1 (5)	C26—O2—C27—C28	179.8 (3)
N1—C14—C15—N2	4.7 (8)		

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
C11—H11A···O1 ⁱ	0.93	2.54	3.343 (6)	146

Symmetry codes: (i) $x, y+1, z$.

Fig. 1




Fig. 2

