

2,6-Dibromo-4-(2-hydroxyethyl)phenol

Ding-qiang Lu,^{a*} Hong Chai,^a Xiu-quan Ling,^a Jia Chen^a and Jia-li Wang^b

^aState Key Laboratory of Materials-Oriented Chemical Engineering, School of Pharmaceutical Sciences, Nanjing University of Technology, Xinmofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China, and ^bState Key Laboratory of Materials-Oriented Chemical Engineering, College of Life Science and Pharmaceutical Engineering, Nanjing University of Technology, Xinmofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China
Correspondence e-mail: wangzhe_guilai@126.com


Received 1 November 2011; accepted 4 November 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.014\text{ \AA}$; R factor = 0.067; wR factor = 0.161; data-to-parameter ratio = 15.7.

The title compound, $\text{C}_8\text{H}_8\text{Br}_2\text{O}_2$, crystallized with two independent molecules (*A* and *B*) in the asymmetric unit. They differ in the conformation of the 2-hydroxyethyl chain with the $\text{C}-\text{C}-\text{C}-\text{O}$ torsion angle being $-68.0(12)^\circ$ in molecule *A* and $172.2(9)^\circ$ in molecule *B*. In the crystal, the *A* molecules are linked *via* pairs of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming inversion dimers, while the *B* molecules are linked *via* an $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond, forming a polymeric chain propagating in [010]. In addition, there are $\text{O}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{Br}$ hydrogen bonds, and $\text{Br}\cdots\text{Br}$ [$3.599(2)\text{ \AA}$] and $\pi-\pi$ interactions [centroid-centroid distances = $3.581(6)$ and $3.931(6)\text{ \AA}$], leading to the formation of a two-dimensional network parallel to (001).

Related literature

For background and further synthetic details, see: Guerard *et al.* (2009); Bovicelli *et al.* (2007). For standard bond-length data, see: Allen *et al.* (1987). For a related structure, see: Zhu *et al.* (2011)

**Experimental***Crystal data*

$\text{C}_8\text{H}_8\text{Br}_2\text{O}_2$
 $M_r = 295.94$
Triclinic, $P\bar{1}$
 $a = 8.5740(17)\text{ \AA}$
 $b = 9.845(2)\text{ \AA}$
 $c = 11.392(2)\text{ \AA}$
 $\alpha = 86.08(3)^\circ$
 $\beta = 75.79(3)^\circ$

$\gamma = 87.39(3)^\circ$

$V = 929.6(3)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 8.68\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.20 \times 0.10 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.276$, $T_{\max} = 0.478$
3664 measured reflections

3416 independent reflections

1874 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.087$

3 standard reflections every 200
reflections
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.067$
 $wR(F^2) = 0.161$
 $S = 1.00$
3416 reflections
217 parameters

1 restraint
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.63\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.72\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots\text{A}$	$D-\text{H}$	$\text{H}\cdots\text{A}$	$D\cdots\text{A}$	$D-\text{H}\cdots\text{A}$
$\text{O}1-\text{H}1\text{O}\cdots\text{O}2^{\text{i}}$	0.82	1.93	2.665 (11)	149
$\text{O}4-\text{H}4\text{O}\cdots\text{O}3^{\text{ii}}$	0.85	2.09	2.837 (9)	146
$\text{O}2-\text{H}2\text{O}\cdots\text{O}4^{\text{iii}}$	0.82	2.09	2.896 (10)	168
$\text{O}3-\text{H}3\text{O}\cdots\text{Br}2^{\text{iv}}$	0.85	2.55	3.291 (7)	147

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x, y + 1, z$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x + 1, y - 1, z - 1$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

This research work was supported financially by the Department of Science and Technology of Jiangsu Province (BE200830457) and the '863' project (2007 A 02Z211) from the Ministry of Science and Technology, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2342).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bovicelli, P., Antonioletti, R., Mancini, S., Causio, S., Borioni, G., Ammendola, S. & Barontini, M. (2007). *Synth. Commun.* **37**, 4245–4252.
- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Guerard, K. C., Sabot, C., Racicot, L. & Canesi, S. (2009). *J. Org. Chem.* **74**, 2039–2045.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.

- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zhu, J., Yang, W., Wang, L., Wang, K. & Hu, Y. (2011). *Acta Cryst. E* **67**, o1109.

supplementary materials

Acta Cryst. (2011). E67, o3263-o3264 [doi:10.1107/S1600536811046538]

2,6-Dibromo-4-(2-hydroxyethyl)phenol

D. Lu, H. Chai, X. Ling, J. Chen and J. Wang

Comment

The title compound is used as a key intermediate in drug synthesis, and has been synthesized following a procedure described previously by (Bovicelli *et al.*, 2007). We report herein on its synthesis and crystal structure.

The title compound crystallized with two independent molecules (A & B) in the asymmetric unit (Fig. 1). They differ significantly in conformation, as may be seen from the torsion angles in the 2-hydroxyethyl chain. For molecule A the torsion angles C1—C6—C7—C8 and C6—C7—C8—O2 are -56.6 (14) and -68.0 (13) $^{\circ}$, respectively, while the corresponding torsion angles C13—C14—C15—C16 and C14—C15—C16—O4 in molecule B are -82.6 (12) and 172.2 (9) $^{\circ}$, respectively. The bond lengths (Allen *et al.*, 1987) and bond angles are otherwise within normal ranges.

In the crystal, the A molecules are linked *via* O—H \cdots O hydrogen bonds to form inversion dimers, while the B molecules are linked *via* an O—H \cdots O hydrogen bond to form a polymer chain propagating in [010], see Fig. 2. In the crystal, there are other O—H \cdots O and O—H \cdots Br hydrogen bonds (Table 1), and weak $\pi\cdots\pi$ stacking interactions involving the aromatic rings with their inversion related rings. The centroid-centroid distances are 3.931 (6) Å for Cg1 \cdots Cg1ⁱ [Cg1 is the centroid of ring (C1—C6); symmetry code (i) $-x + 1, -y + 1, -z + 2$] and 3.581 (6) Å for Cg2 \cdots Cg2ⁱⁱ [Cg2 is the centroid of ring (C9—C14); symmetry code (ii) $-x + 2, -y, -z + 1$]. There is also a short Br1 \cdots Br4ⁱⁱⁱ interaction present [3.599 (2) Å; symmetry code: (iii) $-x + 1, -y, -z + 2$]. These interactions result in the formation of two-dimensional networks lying parallel to the *ab* plane (Fig. 3).

Experimental

The title compound was synthesized using a slightly modified version of the procedure reported on by (Bovicelli *et al.*, 2007), using twice the quantity of NaBr. To a solution of 4-hydroxyphenethyl alcohol (217.4 mmol, 30 g) and NaBr (434.8 mmol, 44.34 g) in acetone (600 ml), a solution of oxone (200 g) in water (1 L) was added dropwise at 263 K within 3 h. The progress of the reaction was monitored by thin-layer chromatography (TLC, hexane/ethyl acetate 3:2), and when the reaction was over (complete consumption of the substrate), AcOEt (500 ml) was added to the mixture. The organic layer was separated, and the aqueous phase was extracted with two 300 mL portions of AcOEt (300 ml). The combined organic solutions were washed with water (300 ml), dried over anhydrous Na₂SO₄ (white powder, 100 g), and evaporated. The product obtained in almost quantitative yield (59.7 g), appeared to be spectroscopically pure: white solid. Crystals of the title compound, suitable for X-ray diffraction analysis, were obtained by slow evaporation of an acetone solution at room temperature.

Refinement

H atoms were positioned geometrically and constrained to ride on their parent atoms: O—H = 0.82 or 0.85 Å, C—H = 0.93, 0.98 and 0.97 Å for aromatic, methine and methylene H atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$, where $k = 1.5$ for methyl H atoms, and $k = 1.2$ for all other H atoms.

supplementary materials

Figures




Fig. 1. The molecular structure of the two independent molecules (A and B) of the title compound, with the atom numbering scheme and displacement ellipsoids drawn at 30% probability levels.




Fig. 2. A view along the a axis of the hydrogen bonded inversion dimers of A molecules and the polymer chain of B molecules of the title compound [hydrogen bonds are shown as dashed cyan lines; see Table 1 for details; symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x, y + 1, z$; (iii) $x, y - 1, z$].




Fig. 3. A perspective view along the b axis of the crystal packing of the title compound, showing the O—H···O and O—H···Br hydrogen bonds as dashed lines (see Table 1 for details).

2,6-Dibromo-4-(2-hydroxyethyl)phenol

Crystal data

$C_8H_8Br_2O_2$	$Z = 4$
$M_r = 295.94$	$F(000) = 568$
Triclinic, $P\bar{1}$	$D_x = 2.115 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 8.5740 (17) \text{ \AA}$	Cell parameters from 25 reflections
$b = 9.845 (2) \text{ \AA}$	$\theta = 10\text{--}13^\circ$
$c = 11.392 (2) \text{ \AA}$	$\mu = 8.68 \text{ mm}^{-1}$
$\alpha = 86.08 (3)^\circ$	$T = 293 \text{ K}$
$\beta = 75.79 (3)^\circ$	Block, colourless
$\gamma = 87.39 (3)^\circ$	$0.20 \times 0.10 \times 0.10 \text{ mm}$
$V = 929.6 (3) \text{ \AA}^3$	

Data collection

Enraf–Nonius CAD-4 diffractometer	1874 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.087$
graphite	$\theta_{\max} = 25.4^\circ, \theta_{\min} = 1.9^\circ$
$\omega/2\theta$ scans	$h = 0 \rightarrow 10$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$k = -11 \rightarrow 11$
$T_{\min} = 0.276, T_{\max} = 0.478$	$l = -13 \rightarrow 13$
3664 measured reflections	3 standard reflections every 200 reflections

3416 independent reflections intensity decay: 1%

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.067$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.161$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.072P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
3416 reflections	$(\Delta/\sigma)_{\max} < 0.001$
217 parameters	$\Delta\rho_{\max} = 0.63 \text{ e \AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.72 \text{ e \AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.68665 (14)	0.22580 (14)	1.16796 (12)	0.0661 (5)
Br2	0.15646 (15)	0.59523 (13)	1.18207 (12)	0.0703 (5)
O1	0.4403 (9)	0.4472 (8)	1.2658 (7)	0.070 (3)
O2	0.4818 (9)	0.3052 (8)	0.6910 (7)	0.066 (3)
C1	0.4707 (12)	0.2251 (11)	1.0171 (9)	0.050 (4)
C2	0.5035 (11)	0.2892 (9)	1.1125 (9)	0.043 (3)
C3	0.4101 (13)	0.3990 (10)	1.1676 (9)	0.049 (4)
C4	0.2875 (12)	0.4483 (10)	1.1133 (9)	0.050 (3)
C5	0.2490 (12)	0.3861 (10)	1.0200 (9)	0.050 (4)
C6	0.3404 (13)	0.2735 (11)	0.9689 (10)	0.053 (4)
C7	0.2967 (12)	0.1977 (12)	0.8703 (9)	0.056 (4)
C8	0.4333 (12)	0.1821 (9)	0.7602 (9)	0.049 (3)
Br3	1.12544 (15)	-0.02904 (13)	0.15184 (12)	0.0742 (5)
Br4	0.64211 (13)	-0.12778 (11)	0.58132 (10)	0.0558 (4)
O3	0.8985 (8)	-0.1939 (6)	0.3477 (6)	0.049 (2)
O4	0.7207 (8)	0.5766 (7)	0.4611 (6)	0.056 (3)
C9	0.9903 (11)	0.1713 (10)	0.3173 (9)	0.046 (4)
C10	0.9906 (11)	0.0338 (10)	0.2919 (9)	0.045 (3)

supplementary materials

C11	0.8926 (11)	-0.0623 (9)	0.3728 (9)	0.039 (3)
C12	0.7878 (10)	-0.0124 (9)	0.4787 (9)	0.036 (3)
C13	0.7884 (11)	0.1234 (10)	0.5006 (10)	0.048 (3)
C14	0.8884 (11)	0.2154 (10)	0.4232 (10)	0.047 (3)
C15	0.8832 (12)	0.3638 (11)	0.4564 (10)	0.052 (4)
C16	0.7456 (13)	0.4450 (10)	0.4198 (11)	0.059 (4)
H1	0.53400	0.15080	0.98510	0.0600*
H1O	0.46340	0.52760	1.25190	0.1050*
H2O	0.41630	0.32850	0.65120	0.0990*
H5	0.16160	0.41900	0.99050	0.0600*
H7A	0.26060	0.10790	0.90340	0.0670*
H7B	0.20760	0.24600	0.84580	0.0670*
H8A	0.52520	0.14080	0.78600	0.0580*
H8B	0.40200	0.11990	0.70820	0.0580*
H3O	0.96250	-0.21940	0.28290	0.0590*
H4O	0.80520	0.62110	0.43080	0.0840*
H9	1.05730	0.23190	0.26410	0.0560*
H13	0.71880	0.15470	0.57020	0.0580*
H15A	0.87040	0.36660	0.54320	0.0620*
H15B	0.98440	0.40530	0.41600	0.0620*
H16A	0.64730	0.39530	0.45080	0.0710*
H16C	0.76670	0.45120	0.33200	0.0710*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0559 (7)	0.0740 (9)	0.0707 (8)	0.0040 (6)	-0.0190 (6)	-0.0112 (6)
Br2	0.0754 (9)	0.0556 (8)	0.0731 (9)	0.0157 (6)	-0.0043 (6)	-0.0185 (6)
O1	0.079 (6)	0.067 (6)	0.062 (5)	-0.009 (4)	-0.010 (4)	-0.010 (4)
O2	0.074 (5)	0.066 (5)	0.056 (5)	-0.011 (4)	-0.010 (4)	-0.015 (4)
C1	0.043 (6)	0.055 (7)	0.041 (6)	-0.001 (5)	0.013 (5)	-0.012 (5)
C2	0.045 (6)	0.034 (5)	0.050 (6)	-0.003 (4)	-0.009 (5)	-0.016 (4)
C3	0.062 (7)	0.038 (6)	0.047 (6)	-0.007 (5)	-0.007 (5)	-0.016 (5)
C4	0.049 (6)	0.045 (6)	0.046 (6)	0.006 (5)	0.003 (5)	0.011 (5)
C5	0.042 (6)	0.051 (7)	0.052 (7)	0.008 (5)	-0.003 (5)	-0.010 (5)
C6	0.049 (6)	0.054 (7)	0.057 (7)	-0.003 (5)	-0.010 (5)	-0.018 (5)
C7	0.052 (6)	0.065 (8)	0.049 (7)	-0.004 (6)	-0.003 (5)	-0.023 (5)
C8	0.062 (7)	0.032 (5)	0.051 (6)	-0.002 (5)	-0.009 (5)	-0.012 (4)
Br3	0.0710 (8)	0.0624 (8)	0.0702 (9)	0.0004 (6)	0.0217 (6)	-0.0141 (6)
Br4	0.0524 (7)	0.0416 (6)	0.0638 (7)	-0.0090 (5)	0.0075 (5)	-0.0094 (5)
O3	0.055 (4)	0.037 (4)	0.046 (4)	0.004 (3)	0.008 (3)	-0.016 (3)
O4	0.056 (5)	0.044 (4)	0.065 (5)	0.000 (4)	-0.004 (4)	-0.017 (4)
C9	0.040 (6)	0.035 (6)	0.056 (7)	0.000 (5)	0.000 (5)	0.011 (5)
C10	0.038 (5)	0.048 (6)	0.046 (6)	-0.007 (5)	0.001 (4)	-0.012 (5)
C11	0.043 (5)	0.024 (5)	0.051 (6)	0.002 (4)	-0.013 (4)	-0.017 (4)
C12	0.033 (5)	0.024 (5)	0.051 (6)	-0.001 (4)	-0.007 (4)	-0.010 (4)
C13	0.030 (5)	0.045 (6)	0.063 (7)	0.010 (5)	0.002 (5)	-0.012 (5)
C14	0.039 (5)	0.045 (5)	0.061 (7)	0.003 (5)	-0.016 (5)	-0.015 (5)

C15	0.043 (6)	0.052 (6)	0.062 (7)	0.005 (5)	-0.012 (5)	-0.015 (5)
C16	0.061 (7)	0.038 (6)	0.071 (8)	0.007 (5)	-0.002 (6)	-0.016 (5)

Geometric parameters (\AA , $^\circ$)

Br1—C2	1.897 (10)	C1—H1	0.9300
Br2—C4	1.882 (10)	C5—H5	0.9300
Br3—C10	1.852 (10)	C7—H7B	0.9700
Br4—C12	1.857 (9)	C7—H7A	0.9700
O1—C3	1.330 (13)	C8—H8B	0.9700
O2—C8	1.420 (12)	C8—H8A	0.9700
O1—H1O	0.8200	C9—C10	1.403 (14)
O2—H2O	0.8200	C9—C14	1.388 (15)
O3—C11	1.342 (11)	C10—C11	1.423 (14)
O4—C16	1.398 (12)	C11—C12	1.419 (14)
O3—H3O	0.8500	C12—C13	1.377 (13)
O4—H4O	0.8500	C13—C14	1.387 (15)
C1—C2	1.385 (14)	C14—C15	1.530 (15)
C1—C6	1.412 (15)	C15—C16	1.522 (16)
C2—C3	1.404 (14)	C9—H9	0.9300
C3—C4	1.398 (15)	C13—H13	0.9300
C4—C5	1.376 (14)	C15—H15A	0.9700
C5—C6	1.403 (15)	C15—H15B	0.9700
C6—C7	1.518 (15)	C16—H16A	0.9700
C7—C8	1.502 (14)	C16—H16C	0.9700
C3—O1—H1O	109.00	C7—C8—H8B	109.00
C8—O2—H2O	109.00	O2—C8—H8A	109.00
C11—O3—H3O	119.00	C10—C9—C14	118.9 (9)
C16—O4—H4O	108.00	C9—C10—C11	122.0 (9)
C2—C1—C6	119.4 (10)	Br3—C10—C11	117.8 (7)
C1—C2—C3	123.3 (9)	Br3—C10—C9	120.2 (7)
Br1—C2—C1	117.5 (7)	O3—C11—C12	122.1 (8)
Br1—C2—C3	119.2 (7)	O3—C11—C10	120.7 (9)
O1—C3—C2	119.7 (10)	C10—C11—C12	117.2 (8)
C2—C3—C4	115.5 (9)	Br4—C12—C13	120.5 (8)
O1—C3—C4	124.8 (9)	Br4—C12—C11	119.9 (7)
Br2—C4—C3	118.1 (7)	C11—C12—C13	119.6 (9)
Br2—C4—C5	118.8 (8)	C12—C13—C14	122.8 (10)
C3—C4—C5	122.9 (9)	C9—C14—C15	121.4 (9)
C4—C5—C6	120.6 (10)	C9—C14—C13	119.5 (9)
C1—C6—C5	118.1 (10)	C13—C14—C15	119.1 (9)
C5—C6—C7	122.1 (10)	C14—C15—C16	111.6 (9)
C1—C6—C7	119.6 (10)	O4—C16—C15	114.7 (9)
C6—C7—C8	113.7 (9)	C10—C9—H9	121.00
O2—C8—C7	115.0 (8)	C14—C9—H9	121.00
C2—C1—H1	120.00	C12—C13—H13	119.00
C6—C1—H1	120.00	C14—C13—H13	119.00
C6—C5—H5	120.00	C14—C15—H15A	109.00
C4—C5—H5	120.00	C14—C15—H15B	109.00

supplementary materials


C6—C7—H7A	109.00	C16—C15—H15A	109.00
C8—C7—H7B	109.00	C16—C15—H15B	109.00
C6—C7—H7B	109.00	H15A—C15—H15B	108.00
C8—C7—H7A	109.00	O4—C16—H16A	109.00
H7A—C7—H7B	108.00	O4—C16—H16C	109.00
O2—C8—H8B	109.00	C15—C16—H16A	109.00
C7—C8—H8A	109.00	C15—C16—H16C	109.00
H8A—C8—H8B	108.00	H16A—C16—H16C	108.00
C6—C1—C2—Br1	-176.8 (8)	C14—C9—C10—Br3	-179.8 (8)
C6—C1—C2—C3	2.0 (16)	C14—C9—C10—C11	-1.6 (15)
C2—C1—C6—C5	0.2 (15)	C10—C9—C14—C13	-0.7 (15)
C2—C1—C6—C7	-175.8 (9)	C10—C9—C14—C15	179.6 (9)
Br1—C2—C3—O1	-7.3 (13)	Br3—C10—C11—O3	0.0 (13)
Br1—C2—C3—C4	173.7 (7)	Br3—C10—C11—C12	-178.9 (7)
C1—C2—C3—O1	173.9 (10)	C9—C10—C11—O3	-178.2 (9)
C1—C2—C3—C4	-5.1 (15)	C9—C10—C11—C12	2.9 (14)
O1—C3—C4—Br2	0.9 (14)	O3—C11—C12—Br4	-4.8 (13)
O1—C3—C4—C5	-172.6 (10)	O3—C11—C12—C13	179.2 (9)
C2—C3—C4—Br2	179.8 (7)	C10—C11—C12—Br4	174.2 (7)
C2—C3—C4—C5	6.3 (15)	C10—C11—C12—C13	-1.9 (14)
Br2—C4—C5—C6	-177.9 (8)	Br4—C12—C13—C14	-176.3 (8)
C3—C4—C5—C6	-4.5 (16)	C11—C12—C13—C14	-0.2 (15)
C4—C5—C6—C1	0.9 (16)	C12—C13—C14—C9	1.6 (16)
C4—C5—C6—C7	176.8 (10)	C12—C13—C14—C15	-178.7 (9)
C1—C6—C7—C8	-56.6 (13)	C9—C14—C15—C16	97.1 (12)
C5—C6—C7—C8	127.6 (11)	C13—C14—C15—C16	-82.6 (12)
C6—C7—C8—O2	-68.0 (12)	C14—C15—C16—O4	172.2 (9)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1—H1O \cdots O2 ⁱ	0.82	1.93	2.665 (11)	149
O4—H4O \cdots O3 ⁱⁱ	0.85	2.09	2.837 (9)	146
O2—H2O \cdots O4 ⁱⁱⁱ	0.82	2.09	2.896 (10)	168
O3—H3O \cdots Br2 ^{iv}	0.85	2.55	3.291 (7)	147

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $x, y+1, z$; (iii) $-x+1, -y+1, -z+1$; (iv) $x+1, y-1, z-1$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

