

4-Chloroanilinium 4-methylbenzene-sulfonate

Jerry P. Jasinski,^{a*} James A. Golen,^a A. S. Praveen,^b
H. S. Yathirajan^b and B. Narayana^c

^aDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, ^bDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and ^cDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri, 574 199, India
Correspondence e-mail: jjasinski@keene.edu


Received 29 October 2011; accepted 8 November 2011

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.032; wR factor = 0.088; data-to-parameter ratio = 18.9.

In the crystal structure of the title salt, $\text{C}_6\text{H}_7\text{ClN}^+\cdot\text{C}_7\text{H}_7\text{O}_3\text{S}^-$, the cations and anions are linked *via* $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds into double chains in [101]. Weak intermolecular $\text{C}-\text{H}\cdots\pi$ -ring interactions link these chains into layers parallel to the *ac* plane.

Related literature

For background literature concerning molecular-ionic compounds, see: Czupinski *et al.* (2002); Katrusiak & Szafranski (2006). For related structures, see: Chanawanno *et al.* (2009); Chantrapromma *et al.* (2010); Collier *et al.* (2006); Fun *et al.* (2010); Kobkeathawin *et al.* (2009); Li *et al.* (2005); Lin, (2010); Rahmouni *et al.* (2010); Smith *et al.* (2009); Tabatabaee & Noozari, (2011); Wu *et al.* (2009); Zhang & Liu (2010).



Experimental

Crystal data

$\text{C}_6\text{H}_7\text{ClN}^+\cdot\text{C}_7\text{H}_7\text{O}_3\text{S}^-$
 $M_r = 299.76$
Triclinic, $P\bar{1}$
 $a = 5.7253$ (5) Å

$b = 7.5160$ (6) Å
 $c = 15.7642$ (13) Å
 $\alpha = 95.166$ (6)°
 $\beta = 96.148$ (7)°

$\gamma = 92.353$ (7)°
 $V = 670.83$ (10) Å³
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 0.44$ mm⁻¹
 $T = 173$ K
 $0.40 \times 0.20 \times 0.12$ mm

Data collection

Oxford Diffraction Xcalibur Eos
Gemini diffractometer
Absorption correction: multi-scan
(*CrysAlis RED*; Oxford
Diffraction, 2010)
 $T_{\min} = 0.843$, $T_{\max} = 0.949$

5280 measured reflections
3439 independent reflections
3144 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.088$
 $S = 1.08$
3439 reflections
182 parameters
6 restraints

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\text{max}} = 0.42$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.45$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg2 is the centroid of the C8–C13 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1NC}\cdots\text{O1}^{\text{i}}$	0.92 (1)	2.02 (1)	2.8579 (16)	151 (2)
$\text{N1}-\text{H1NC}\cdots\text{O1}$	0.92 (1)	2.42 (2)	3.0814 (16)	129 (1)
$\text{N1}-\text{H1NB}\cdots\text{O3}^{\text{ii}}$	0.92 (1)	1.88 (1)	2.7940 (15)	175 (2)
$\text{N1}-\text{H1NA}\cdots\text{O2}^{\text{iii}}$	0.93 (1)	1.98 (1)	2.8764 (16)	163 (2)
$\text{C2}-\text{H2A}\cdots\text{Cg2}^{\text{i}}$	0.95	2.91	3.5340 (16)	124

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $x - 1, y, z$; (iii) $-x + 1, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

ASP and HSY thank the UoM for research facilities. JPI acknowledges the NSF-MRI program (grant No. CHE1039027) for funds to purchase the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5699).

References

- Chanawanno, K., Chantrapromma, S. & Fun, H.-K. (2009). *Anal. Sci.* **25**, 127–128.
Chantrapromma, S., Chanawanno, K. & Fun, H.-K. (2010). *Acta Cryst.* **E66**, o1975–o1976.
Collier, E. A., Davey, R. J., Black, S. N. & Roberts, R. J. (2006). *Acta Cryst.* **B62**, 498–505.
Czupinski, O., Bator, G., Ciunik, Z., Jakubas, R., Medycki, W. & Swiergiel, J. (2002). *J. Phys. Condens. Matter*, **14**, 8497–8512.
Fun, H.-K., Kobkeathawin, T. & Chantrapromma, S. (2010). *Acta Cryst.* **E66**, o1053–o1054.
Katrusiak, A. & Szafranski, M. (2006). *J. Am. Chem. Soc.* **128**, 15775–15785.
Kobkeathawin, T., Suwunwong, T., Chantrapromma, S. & Fun, H.-K. (2009). *Acta Cryst.* **E65**, o76–o77.
Li, X.-M., Lu, L.-P., Feng, S.-S., Zhang, H.-M., Qin, S.-D. & Zhu, M.-L. (2005). *Acta Cryst.* **E61**, o811–o813.
Lin, J. R. (2010). *Acta Cryst.* **E66**, o1557.

- Oxford Diffraction (2010). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Rahmouni, H., Smirani, W., Rzaigui, M. & S. Al-Deyab, S. (2010). *Acta Cryst.* **E66**, o993.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Smith, G., Wermuth, U. D. & White, J. M. (2009). *Acta Cryst.* **E65**, o2111.
- Tabatabaee, M. & Noozari, N. (2011). *Acta Cryst.* **E67**, o1457.
- Wu, T.-Q., Xia, L., Hu, A.-X. & Ye, J. (2009). *Acta Cryst.* **E65**, o368.
- Zhang, Y. & Liu, X. (2010). *Acta Cryst.* **E66**, o790.

supplementary materials

Acta Cryst. (2011). E67, o3288–o3289 [doi:10.1107/S160053681104712X]

4-Chloroanilinium 4-methylbenzenesulfonate

J. P. Jasinski, J. A. Golen, A. S. Praveen, H. S. Yathirajan and B. Narayana

Comment

A variety of pharmaceutical drugs are prepared as salts of benzenesulfonic acid and are known as besylates. Recently much attention has been devoted to simple molecular–ionic crystals containing organic cations and anions due to the tunability of their special structural features and their interesting physical properties (Czupinski *et al.*, 2002; Katrusiak & Szafranski, 2006). In the title compound, the proton of the sulfonic group of sulfonic acid has been transferred to the N atom of the 4-chloroaniline molecule, leading to the formation of the molecular complex, (I).

Crystal structures of some benzenesulfonate derivatives, viz., 2,4,6-triamino-1,3,5-triazin-1-ium 4-methylbenzenesulfonate monohydrate (Li *et al.*, 2005), ephedrine besylate (Collier *et al.*, 2006), 2-ethyl-6-methylanilinium 4-methylbenzenesulfonate (Wu *et al.*, 2009), 2-[(E)-2-(4-ethoxyphenyl)ethenyl]-1-methylpyridinium 4-methylbenzenesulfonate monohydrate (Chanawanno *et al.*, 2009), (E)-2-[4-(dimethylamino)styryl]-1-methylquinolinium 4-methylbenzenesulfonate monohydrate (Kobkeatthawin *et al.*, 2009), 4-chloroanilinium 2-carboxy-4,5-dichlorobenzoate (Smith *et al.*, 2009), 4-chloroanilinium (4-chlorophenyl)guanidinium dichloride hemihydrates (Zhang & Liu, 2010), 4-chloroanilinium hydrogen oxalate hemihydrates (Rahmouni *et al.*, 2010), 4-(cyanomethyl)anilinium 4-methylbenzene sulfonate monohydrate (Lin, 2010), 1-methyl-2-[(E)-2-(2-thienyl)ethenyl] quinolinium 4-bromobenzenesulfonate (Fun *et al.*, 2010), (E)-2-[4-(dimethylamino)styryl]-1-methylpyridinium 4-methylbenzenesulfonate monohydrate (Chantrapromma *et al.*, 2010), 2-aminopyrimidin-1-ium 4-methylbenzenesulfonate (Tabatabaee & Noozari, 2011), have been reported. In view of the importance of benzenesulphonic acid, we report herein the crystal structure of the title compound (I).

In the crystal structure of the title salt, $C_6H_7ClN^+ \cdot C_7H_7O_3S^-$, (Fig. 1) the cations and anions are linked *via* N—H \cdots O hydrogen bonds into doubled chains in [101] (Fig. 2). Weak intermolecular C—H \cdots Cg2 π -ring interactions (table 1) link further these chains into layers parallel to the *ac* plane. [Cg2 = C8—C13 centroid]

Experimental

4-methylbenzenesulfonic acid monohydrate (1g, 5.25 mmol) was added to a stirred solution of 4- chloroaniline (0.67 g, 5.25 mmol) in methanol (10 mL). The resulting suspension was dissolved in chloroform (10 ml) and stirred at 323 K for 10 minutes and cooled to room temperature to afford the title compound (I). Single crystals were grown from a mixture of chloroform and methanol by the slow evaporation method (m.p.: 524–532 K).

Refinement

H1NA, H1NB and H1NC were located by a Fourier map and refined isotropically (for N1, $dfix = 0.94 (2)\text{\AA}$; $dang = 1.50 (2)\text{\AA}$). All of the remaining H atoms were placed in their calculated positions and then refined using the riding model with Atom—H lengths of 0.95 (CH) or 0.98 \AA (CH₃). Isotropic displacement parameters for these atoms were set to 1.18–1.21 (CH) or 1.51 (CH₃) times U_{eq} of the parent atom.

Figures




Fig. 1. Molecular structure of the title compound showing the atom labeling scheme and 30% probability displacement ellipsoids.




Fig. 2. Packing diagram of the title compound viewed along the *b* axis. Dashed lines indicate N—H...O hydrogen bonds forming infinite 1-D chains along the *c* axis.

4-Chloroanilinium 4-methylbenzenesulfonate

Crystal data

$C_6H_7ClN^+ \cdot C_7H_7O_3S^-$

$M_r = 299.76$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 5.7253$ (5) Å

$b = 7.5160$ (6) Å

$c = 15.7642$ (13) Å

$\alpha = 95.166$ (6)°

$\beta = 96.148$ (7)°

$\gamma = 92.353$ (7)°

$V = 670.83$ (10) Å³

$Z = 2$

$F(000) = 312$

$D_x = 1.484$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2538 reflections

$\theta = 3.6$ – 29.9 °

$\mu = 0.44$ mm⁻¹

$T = 173$ K

Block, colorless

$0.40 \times 0.20 \times 0.12$ mm

Data collection

Oxford Diffraction Xcalibur Eos Gemini diffractometer

Radiation source: Enhance (Mo) X-ray Source graphite

Detector resolution: 16.1500 pixels mm⁻¹

ω scans

Absorption correction: multi-scan (*Crys.Alis RED*; Oxford Diffraction, 2010)

$T_{\min} = 0.843$, $T_{\max} = 0.949$

5280 measured reflections

3439 independent reflections

3144 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 30.0$ °, $\theta_{\min} = 3.6$ °

$h = -7 \rightarrow 7$

$k = -9 \rightarrow 9$

$l = -22 \rightarrow 21$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.032$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.088$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.0406P)^2 + 0.2844P]$
3439 reflections	where $P = (F_o^2 + 2F_c^2)/3$
182 parameters	$(\Delta/\sigma)_{\max} = 0.001$
6 restraints	$\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.45 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.64768 (6)	0.72752 (4)	0.391167 (19)	0.01562 (9)
Cl1	-0.20941 (8)	0.85567 (6)	0.86957 (3)	0.03622 (12)
O1	0.56763 (19)	0.89688 (13)	0.42702 (6)	0.0227 (2)
O2	0.52819 (19)	0.57587 (13)	0.42253 (6)	0.0231 (2)
O3	0.90320 (18)	0.71964 (16)	0.40103 (6)	0.0281 (2)
N1	0.2327 (2)	0.74505 (16)	0.54631 (7)	0.0179 (2)
H1NC	0.329 (3)	0.8434 (17)	0.5411 (11)	0.022*
H1NB	0.123 (2)	0.730 (2)	0.4992 (10)	0.022*
H1NA	0.323 (3)	0.6460 (18)	0.5467 (11)	0.022*
C1	0.5595 (2)	0.71390 (16)	0.27970 (8)	0.0150 (2)
C2	0.7106 (2)	0.7789 (2)	0.22507 (9)	0.0219 (3)
H2A	0.8638	0.8264	0.2470	0.026*
C3	0.6356 (3)	0.7737 (2)	0.13818 (9)	0.0277 (3)
H3A	0.7385	0.8182	0.1007	0.033*
C4	0.4120 (3)	0.7044 (2)	0.10515 (9)	0.0255 (3)
C5	0.2646 (3)	0.6388 (2)	0.16103 (10)	0.0267 (3)
H5A	0.1123	0.5896	0.1391	0.032*

supplementary materials

C6	0.3357 (2)	0.6439 (2)	0.24820 (9)	0.0223 (3)
H6A	0.2326	0.6001	0.2858	0.027*
C7	0.3278 (4)	0.7036 (3)	0.01100 (10)	0.0412 (4)
H7A	0.2057	0.6079	-0.0056	0.062*
H7B	0.4601	0.6835	-0.0227	0.062*
H7C	0.2628	0.8191	0.0002	0.062*
C8	-0.0816 (3)	0.82188 (18)	0.77464 (9)	0.0210 (3)
C9	-0.1916 (2)	0.88215 (18)	0.70096 (9)	0.0211 (3)
H9A	-0.3353	0.9407	0.7021	0.025*
C10	-0.0887 (2)	0.85572 (18)	0.62520 (9)	0.0191 (3)
H10A	-0.1614	0.8961	0.5739	0.023*
C11	0.1203 (2)	0.77012 (17)	0.62529 (8)	0.0161 (2)
C12	0.2313 (2)	0.71189 (17)	0.69946 (8)	0.0184 (3)
H12A	0.3764	0.6553	0.6985	0.022*
C13	0.1287 (3)	0.73699 (19)	0.77499 (9)	0.0218 (3)
H13A	0.2014	0.6966	0.8263	0.026*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.01884 (16)	0.01791 (16)	0.01046 (14)	0.00399 (11)	0.00187 (11)	0.00159 (11)
Cl1	0.0423 (2)	0.0452 (2)	0.02487 (19)	0.01334 (18)	0.01570 (17)	0.00412 (16)
O1	0.0323 (5)	0.0182 (5)	0.0174 (5)	0.0038 (4)	0.0042 (4)	-0.0023 (4)
O2	0.0340 (6)	0.0194 (5)	0.0183 (5)	0.0056 (4)	0.0089 (4)	0.0066 (4)
O3	0.0191 (5)	0.0480 (7)	0.0165 (5)	0.0067 (4)	-0.0019 (4)	0.0021 (4)
N1	0.0186 (5)	0.0198 (5)	0.0155 (5)	0.0024 (4)	0.0016 (4)	0.0022 (4)
C1	0.0190 (6)	0.0143 (5)	0.0120 (5)	0.0029 (4)	0.0017 (4)	0.0012 (4)
C2	0.0200 (6)	0.0294 (7)	0.0162 (6)	-0.0022 (5)	0.0019 (5)	0.0035 (5)
C3	0.0291 (8)	0.0391 (8)	0.0164 (6)	0.0010 (6)	0.0059 (6)	0.0079 (6)
C4	0.0311 (8)	0.0308 (8)	0.0138 (6)	0.0089 (6)	-0.0019 (5)	-0.0004 (5)
C5	0.0219 (7)	0.0341 (8)	0.0215 (7)	-0.0002 (6)	-0.0046 (5)	-0.0021 (6)
C6	0.0194 (6)	0.0282 (7)	0.0193 (6)	-0.0018 (5)	0.0023 (5)	0.0030 (5)
C7	0.0472 (11)	0.0604 (12)	0.0147 (7)	0.0126 (9)	-0.0049 (7)	0.0015 (7)
C8	0.0243 (7)	0.0208 (6)	0.0184 (6)	0.0018 (5)	0.0060 (5)	0.0002 (5)
C9	0.0181 (6)	0.0201 (6)	0.0254 (7)	0.0040 (5)	0.0031 (5)	0.0013 (5)
C10	0.0191 (6)	0.0195 (6)	0.0183 (6)	0.0026 (5)	-0.0017 (5)	0.0032 (5)
C11	0.0179 (6)	0.0147 (6)	0.0155 (6)	0.0005 (4)	0.0016 (5)	0.0010 (4)
C12	0.0185 (6)	0.0182 (6)	0.0185 (6)	0.0041 (5)	-0.0005 (5)	0.0022 (5)
C13	0.0262 (7)	0.0230 (7)	0.0163 (6)	0.0042 (5)	0.0003 (5)	0.0043 (5)

Geometric parameters (\AA , $^\circ$)

S1—O2	1.4565 (10)	C5—C6	1.387 (2)
S1—O1	1.4572 (10)	C5—H5A	0.9500
S1—O3	1.4587 (11)	C6—H6A	0.9500
S1—C1	1.7682 (13)	C7—H7A	0.9800
Cl1—C8	1.7381 (14)	C7—H7B	0.9800
N1—C11	1.4626 (17)	C7—H7C	0.9800
N1—H1NC	0.920 (12)	C8—C9	1.383 (2)

N1—H1NB	0.916 (12)	C8—C13	1.385 (2)
N1—H1NA	0.925 (12)	C9—C10	1.3894 (19)
C1—C6	1.3880 (19)	C9—H9A	0.9500
C1—C2	1.3892 (18)	C10—C11	1.3817 (18)
C2—C3	1.3875 (19)	C10—H10A	0.9500
C2—H2A	0.9500	C11—C12	1.3850 (18)
C3—C4	1.391 (2)	C12—C13	1.3845 (19)
C3—H3A	0.9500	C12—H12A	0.9500
C4—C5	1.391 (2)	C13—H13A	0.9500
C4—C7	1.510 (2)		
O2—S1—O1	111.46 (6)	C5—C6—C1	119.09 (13)
O2—S1—O3	113.20 (7)	C5—C6—H6A	120.5
O1—S1—O3	113.02 (7)	C1—C6—H6A	120.5
O2—S1—C1	106.02 (6)	C4—C7—H7A	109.5
O1—S1—C1	106.15 (6)	C4—C7—H7B	109.5
O3—S1—C1	106.33 (6)	H7A—C7—H7B	109.5
C11—N1—H1NC	110.2 (11)	C4—C7—H7C	109.5
C11—N1—H1NB	111.0 (11)	H7A—C7—H7C	109.5
H1NC—N1—H1NB	108.1 (13)	H7B—C7—H7C	109.5
C11—N1—H1NA	110.6 (11)	C9—C8—C13	121.80 (13)
H1NC—N1—H1NA	108.1 (13)	C9—C8—C11	119.03 (11)
H1NB—N1—H1NA	108.8 (13)	C13—C8—C11	119.16 (11)
C6—C1—C2	120.72 (12)	C8—C9—C10	118.97 (13)
C6—C1—S1	119.31 (10)	C8—C9—H9A	120.5
C2—C1—S1	119.93 (10)	C10—C9—H9A	120.5
C3—C2—C1	119.31 (13)	C11—C10—C9	119.32 (12)
C3—C2—H2A	120.3	C11—C10—H10A	120.3
C1—C2—H2A	120.3	C9—C10—H10A	120.3
C2—C3—C4	121.00 (14)	C10—C11—C12	121.49 (12)
C2—C3—H3A	119.5	C10—C11—N1	119.76 (12)
C4—C3—H3A	119.5	C12—C11—N1	118.71 (12)
C5—C4—C3	118.61 (13)	C13—C12—C11	119.38 (12)
C5—C4—C7	120.37 (15)	C13—C12—H12A	120.3
C3—C4—C7	121.01 (15)	C11—C12—H12A	120.3
C6—C5—C4	121.26 (14)	C12—C13—C8	119.02 (13)
C6—C5—H5A	119.4	C12—C13—H13A	120.5
C4—C5—H5A	119.4	C8—C13—H13A	120.5
O2—S1—C1—C6	32.26 (12)	C4—C5—C6—C1	-0.8 (2)
O1—S1—C1—C6	-86.39 (12)	C2—C1—C6—C5	0.3 (2)
O3—S1—C1—C6	153.01 (11)	S1—C1—C6—C5	177.95 (11)
O2—S1—C1—C2	-150.07 (11)	C13—C8—C9—C10	0.4 (2)
O1—S1—C1—C2	91.28 (12)	C11—C8—C9—C10	179.74 (11)
O3—S1—C1—C2	-29.32 (13)	C8—C9—C10—C11	0.0 (2)
C6—C1—C2—C3	0.1 (2)	C9—C10—C11—C12	-0.8 (2)
S1—C1—C2—C3	-177.49 (11)	C9—C10—C11—N1	-178.68 (12)
C1—C2—C3—C4	-0.1 (2)	C10—C11—C12—C13	1.2 (2)
C2—C3—C4—C5	-0.4 (2)	N1—C11—C12—C13	179.09 (12)
C2—C3—C4—C7	178.22 (15)	C11—C12—C13—C8	-0.8 (2)

supplementary materials

C3—C4—C5—C6	0.9 (2)	C9—C8—C13—C12	0.0 (2)
C7—C4—C5—C6	-177.77 (15)	C11—C8—C13—C12	-179.35 (11)

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the C8–C13 ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1NC \cdots O1 ⁱ	0.92 (1)	2.02 (1)	2.8579 (16)	151.(2)
N1—H1NC \cdots O1	0.92 (1)	2.42 (2)	3.0814 (16)	129.(1)
N1—H1NB \cdots O3 ⁱⁱ	0.92 (1)	1.88 (1)	2.7940 (15)	175.(2)
N1—H1NA \cdots O2 ⁱⁱⁱ	0.93 (1)	1.98 (1)	2.8764 (15)	163.(2)
C2—H2A \cdots Cg2 ⁱ	0.95	2.91	3.5340 (16)	124

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $x-1, y, z$; (iii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

