

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(4-Chloro-2-methylphenyl)maleamic acid

K. Shakuntala,^a Viktor Vrábek,^b B. Thimme Gowda^{a*} and Jozef Kožíšek^b

^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and ^bInstitute of Physical Chemistry and Chemical Physics, Slovak University of Technology, Radlinského 9, SK-812 37 Bratislava, Slovak Republic
Correspondence e-mail: gowdabt@yahoo.com


Received 8 November 2011; accepted 10 November 2011

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.033; wR factor = 0.089; data-to-parameter ratio = 11.8.

In the molecular structure of the title compound, $\text{C}_{11}\text{H}_{10}\text{ClNO}_3$, the conformation of the N—H bond in the amide segment is *syn* to the *ortho*-methyl group in the phenyl ring. The C=O and O—H bonds of the acid group are in the relatively rare *anti* position with respect to each other. This is an obvious consequence of the hydrogen bond donated to the amide carbonyl group. The central oxobutenoic acid core $\text{C}(\text{=O})-\text{C}=\text{C}-\text{C}-\text{OH}$ is twisted by 31.65 (6)° out of the plane of the 4-chloro-2-methylphenyl ring. An intramolecular O—H...O hydrogen bond occurs. In the crystal, N—H...O hydrogen bonds link the molecules into infinite chains running along the *a* axis.

Related literature

For studies on the effects of substituents on the structures and other aspects of *N*-(aryl)-amides, see: Gowda *et al.* (2000, 2010); Prasad *et al.* (2002), on *N*-(aryl)-methanesulfonamides, see: Jayalakshmi & Gowda (2004), on *N*-(aryl)-arylsulfonamides, see: Shetty & Gowda (2005) and on *N*-chloroarylsulfonamides, see: Gowda & Kumar (2003). For modes of interlinking carboxylic acids by hydrogen bonds, see: Leiserowitz (1976)



Experimental

Crystal data

$\text{C}_{11}\text{H}_{10}\text{ClNO}_3$
 $M_r = 239.65$
Orthorhombic, *Pbca*

$a = 12.1310$ (11) Å
 $b = 7.3990$ (7) Å
 $c = 25.466$ (2) Å

$V = 2285.7$ (3) Å³
 $Z = 8$
Mo $K\alpha$ radiation

$\mu = 0.33$ mm⁻¹
 $T = 295$ K
 $0.45 \times 0.35 \times 0.25$ mm

Data collection

Oxford Diffraction Xcalibur diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2009)
 $T_{\min} = 0.865$, $T_{\max} = 0.918$

17740 measured reflections
1819 independent reflections
1642 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.089$
 $S = 1.02$
1819 reflections
154 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.19$ e Å⁻³
 $\Delta\rho_{\min} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O3}^{\text{i}}$	0.86 (1)	2.10 (1)	2.9556 (19)	174 (2)
$\text{O2}-\text{H2A}\cdots\text{O1}$	0.92 (1)	1.57 (1)	2.4797 (17)	171 (3)

Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, -z$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2002); software used to prepare material for publication: *SHELXL97*, *PLATON* (Spek, 2009) and *WinGX* (Farrugia, 1999).

VV and JK thank the VEGA Grant Agency of the Slovak Ministry of Education (1/0679/11) and the Research and Development Agency of Slovakia (APVV-0202-10) for support and the Structural Funds, Interreg IIIA, for financial support in purchasing the diffractometer. VZR thanks the University Grants Commission, Government of India, New Delhi, for the award of an RFSMS research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2317).

References

- Brandenburg, K. (2002). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Gowda, B. T. & Kumar, B. H. A. (2003). *Oxid. Commun. A*, **26**, 403–425.
Gowda, B. T., Paulus, H. & Fuess, H. (2000). *Z. Naturforsch. Teil A*, **55**, 791–800.
Gowda, B. T., Tokarčík, M., Shakuntala, K., Kožíšek, J. & Fuess, H. (2010). *Acta Cryst. E* **66**, o1554.
Jayalakshmi, K. L. & Gowda, B. T. (2004). *Z. Naturforsch. Teil A*, **59**, 491–500.
Leiserowitz, L. (1976). *Acta Cryst.* **B32**, 775–802.
Oxford Diffraction (2009). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
Prasad, S. M., Sinha, R. B. P., Mandal, D. K. & Rani, A. (2002). *Acta Cryst. E* **58**, o1296–o1297.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Shetty, M. & Gowda, B. T. (2005). *Z. Naturforsch. Teil A*, **60**, 113–120.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2011). E67, o3317 [doi:10.1107/S1600536811047817]

***N*-(4-Chloro-2-methylphenyl)maleamic acid**

K. Shakuntala, V. Vrabel, B. T. Gowda and J. Kozisek

Comment

The amide moiety is the constituent of many biologically significant compounds. As part of our studies on the substituent effects on the structures and other aspects of *N*-(aryl)-amides (Gowda *et al.*, 2000, 2010), *N*-(aryl)-methanesulfonamides (Jayalakshmi & Gowda, 2004), *N*-(aryl)-arylsulfonamides (Shetty & Gowda, 2005) and *N*-chloroarylsulfoamides (Gowda & Kumar, 2003), in the present work, the crystal structure of *N*-(4-chloro-2-methylphenyl)-maleamic acid (I) has been determined (Fig.1).

The conformations of the N—H and the C=O bonds in the amide segment are *anti* to each other. But the conformation of the N—H bond is *syn* to the *ortho*-methyl group in the phenyl ring. In the maleamic acid moiety, the amide C=O bond is *anti* to the adjacent C—H bond, while the carboxyl C=O bond is *syn* to the adjacent C—H bond. The observed rare *anti* conformation of the C=O and O—H bonds of the acid group is similar to that observed in *N*-(2-methylphenyl)-maleamic acid (Gowda *et al.*, 2010). This is an obvious consequence of the hydrogen bond donated to the amide carbonyl group. The central oxobutenoic acid core C(=O)—C=C—C—OH is twisted by 31.65 (6)° out of the plane of the 4-chloro-2-methylphenyl ring. The C2—C3 bond length of 1.333 (2)Å clearly indicates the double bond character.

The various modes of interlinking carboxylic acids by hydrogen bonds is described elsewhere (Leiserowitz, 1976).

In (I), both the intramolecular O—H⋯O and intermolecular N—H⋯O hydrogen bonds have been observed. The packing of molecules linked by N—H⋯O hydrogen bonds into infinite chains running along the *a*-axis is shown in Fig. 2.

Experimental

The solution of maleic anhydride (0.025 mol) in toluene (25 ml) was treated dropwise with the solution of 4-chloro-2-methylaniline (0.025 mol) also in toluene (20 ml) with constant stirring. The resulting mixture was stirred for about 30 min. and set aside for an additional 30 min. at room temperature for the completion of reaction. The mixture was then treated with dilute hydrochloric acid to remove the unreacted 4-chloro-2-methylaniline. The resultant solid *N*-(4-chloro-2-methylphenyl)-maleamic acid was filtered under suction and washed thoroughly with water to remove the unreacted maleic anhydride and maleic acid. It was recrystallized to constant melting point from ethanol. The purity of the compound was checked by elemental analysis and characterized by its infrared spectra.

The plate like colorless single crystals of the title compound used in X-ray diffraction studies were grown in an ethanol solution by slow evaporation (0.5 g in about 30 ml of ethanol) at room temperature.

Refinement

All hydrogen atoms were placed in calculated positions with C—H distances of 0.93Å (C-aromatic) and 0.96Å (C-methyl), and constrained to ride on their parent atoms. Amide and O—H atoms were seen in difference map and were refined with

supplementary materials

the N—H and O—H distances restrained to 0.86 (1) Å and 0.92 (1) Å, respectively. The $U_{\text{iso}}(\text{H})$ values were set at 1.2 U_{eq} (C-aromatic, N) or 1.5 U_{eq} (C-methyl).

Figures




Fig. 1. Molecular structure of the title compound showing the atom labeling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Packing view of the title compound. Molecular chains along a -axis are generated by N—H \cdots O hydrogen bonds which are shown as dashed lines. H atoms not involved in H-bonding have been omitted.

N-(4-Chloro-2-methylphenyl)maleamic acid

Crystal data

$\text{C}_{11}\text{H}_{10}\text{ClNO}_3$

$M_r = 239.65$

Orthorhombic, $Pbca$

$a = 12.1310$ (11) Å

$b = 7.3990$ (7) Å

$c = 25.466$ (2) Å

$V = 2285.7$ (3) Å³

$Z = 8$

$F(000) = 992$

$D_x = 1.393$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1819 reflections

$\theta = 3.6$ – 24.4°

$\mu = 0.33$ mm⁻¹

$T = 295$ K

Plate, colourless

$0.45 \times 0.35 \times 0.25$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer

Radiation source: fine-focus sealed tube
graphite

Detector resolution: 0 pixels mm⁻¹

ω scans with κ offsets

1819 independent reflections

1642 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\text{max}} = 24.4^\circ$, $\theta_{\text{min}} = 3.6^\circ$

$h = -14 \rightarrow 13$

Absorption correction: multi-scan
(*CrysAlis PRO*; Oxford Diffraction, 2009) $k = -8 \rightarrow 8$
 $T_{\min} = 0.865$, $T_{\max} = 0.918$ $l = -29 \rightarrow 28$
17740 measured reflections

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.033$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.089$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0445P)^2 + 0.9459P]$
1819 reflections	where $P = (F_o^2 + 2F_c^2)/3$
154 parameters	$(\Delta/\sigma)_{\max} < 0.001$
2 restraints	$\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.29051 (13)	0.3610 (2)	0.08033 (7)	0.0428 (4)
C2	0.31769 (14)	0.3019 (3)	0.02624 (7)	0.0485 (5)
H2	0.3925	0.2917	0.0188	0.058*
C3	0.25011 (15)	0.2612 (3)	-0.01329 (6)	0.0505 (5)
H3	0.2860	0.2317	-0.0444	0.061*
C4	0.12845 (15)	0.2543 (3)	-0.01670 (7)	0.0496 (5)
C5	0.37645 (12)	0.4444 (2)	0.16488 (6)	0.0368 (4)
C6	0.46315 (13)	0.5525 (2)	0.18298 (6)	0.0383 (4)
C7	0.46319 (14)	0.6020 (2)	0.23578 (6)	0.0431 (4)
H7	0.5205	0.6723	0.2489	0.052*
C8	0.37918 (14)	0.5481 (2)	0.26889 (6)	0.0410 (4)
C9	0.29334 (14)	0.4439 (2)	0.25077 (7)	0.0449 (4)
H9	0.2366	0.4097	0.2732	0.054*

supplementary materials

C10	0.29265 (14)	0.3907 (2)	0.19860 (7)	0.0429 (4)
H10	0.2357	0.3185	0.1861	0.051*
C11	0.55428 (15)	0.6166 (3)	0.14712 (7)	0.0517 (5)
H11A	0.5937	0.5141	0.1336	0.078*
H11B	0.6039	0.6927	0.1664	0.078*
H11C	0.5231	0.6837	0.1185	0.078*
N1	0.37789 (11)	0.38698 (19)	0.11129 (5)	0.0406 (4)
H1	0.4407 (8)	0.369 (2)	0.0965 (6)	0.046 (5)*
O1	0.19506 (10)	0.3884 (2)	0.09562 (5)	0.0656 (4)
O2	0.06813 (10)	0.2998 (3)	0.02376 (5)	0.0746 (5)
H2A	0.1092 (19)	0.340 (3)	0.0519 (7)	0.098 (8)*
O3	0.08638 (11)	0.2036 (2)	-0.05732 (5)	0.0700 (4)
Cl1	0.38181 (4)	0.61481 (7)	0.334722 (17)	0.0585 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0340 (9)	0.0542 (10)	0.0401 (9)	-0.0017 (8)	-0.0029 (7)	-0.0048 (8)
C2	0.0339 (9)	0.0681 (12)	0.0436 (10)	-0.0021 (8)	0.0016 (7)	-0.0083 (9)
C3	0.0436 (9)	0.0720 (12)	0.0359 (9)	-0.0057 (9)	0.0018 (8)	-0.0065 (9)
C4	0.0438 (10)	0.0691 (12)	0.0358 (10)	-0.0091 (9)	-0.0067 (8)	0.0037 (9)
C5	0.0355 (8)	0.0380 (8)	0.0370 (9)	0.0011 (7)	-0.0049 (7)	-0.0031 (7)
C6	0.0355 (8)	0.0360 (8)	0.0434 (9)	-0.0009 (7)	-0.0029 (7)	-0.0012 (7)
C7	0.0441 (10)	0.0389 (9)	0.0463 (10)	-0.0041 (8)	-0.0090 (8)	-0.0061 (7)
C8	0.0509 (10)	0.0366 (8)	0.0356 (9)	0.0025 (8)	-0.0047 (7)	-0.0026 (7)
C9	0.0473 (10)	0.0460 (9)	0.0415 (9)	-0.0040 (8)	0.0030 (7)	0.0015 (8)
C10	0.0410 (9)	0.0452 (9)	0.0425 (9)	-0.0094 (8)	-0.0028 (7)	-0.0027 (7)
C11	0.0440 (10)	0.0592 (11)	0.0519 (11)	-0.0133 (9)	0.0002 (8)	-0.0048 (9)
N1	0.0321 (8)	0.0517 (9)	0.0380 (8)	-0.0033 (6)	-0.0008 (6)	-0.0070 (6)
O1	0.0350 (7)	0.1178 (13)	0.0441 (7)	0.0058 (7)	-0.0029 (5)	-0.0196 (7)
O2	0.0374 (7)	0.1425 (15)	0.0439 (8)	-0.0093 (8)	-0.0041 (6)	-0.0140 (9)
O3	0.0527 (8)	0.1129 (12)	0.0444 (7)	-0.0126 (8)	-0.0140 (6)	-0.0092 (8)
Cl1	0.0767 (4)	0.0599 (3)	0.0388 (3)	-0.0056 (2)	-0.0028 (2)	-0.00955 (19)

Geometric parameters (\AA , $^\circ$)

C1—O1	1.238 (2)	C6—C11	1.510 (2)
C1—O1	1.238 (2)	C7—C8	1.382 (2)
C1—N1	1.335 (2)	C7—H7	0.9300
C1—C2	1.482 (2)	C8—C9	1.375 (2)
C2—C3	1.333 (2)	C8—C11	1.7478 (16)
C2—H2	0.9300	C9—C10	1.386 (2)
C3—C4	1.479 (3)	C9—H9	0.9300
C3—H3	0.9300	C10—H10	0.9300
C4—O3	1.213 (2)	C11—H11A	0.9600
C4—O2	1.308 (2)	C11—H11B	0.9600
C5—C10	1.389 (2)	C11—H11C	0.9600
C5—C6	1.400 (2)	N1—H1	0.861 (5)
C5—N1	1.429 (2)	O1—O1	0.000 (5)

C6—C7	1.394 (2)	O2—H2A	0.920 (5)
O1—C1—O1	0.00 (12)	C8—C7—H7	119.6
O1—C1—N1	122.22 (16)	C6—C7—H7	119.6
O1—C1—N1	122.22 (16)	C9—C8—C7	121.03 (15)
O1—C1—C2	123.26 (15)	C9—C8—C11	119.60 (13)
O1—C1—C2	123.26 (15)	C7—C8—C11	119.37 (13)
N1—C1—C2	114.50 (14)	C8—C9—C10	119.05 (16)
C3—C2—C1	129.17 (16)	C8—C9—H9	120.5
C3—C2—H2	115.4	C10—C9—H9	120.5
C1—C2—H2	115.4	C9—C10—C5	120.47 (15)
C2—C3—C4	131.75 (17)	C9—C10—H10	119.8
C2—C3—H3	114.1	C5—C10—H10	119.8
C4—C3—H3	114.1	C6—C11—H11A	109.5
O3—C4—O2	121.08 (16)	C6—C11—H11B	109.5
O3—C4—C3	118.71 (17)	H11A—C11—H11B	109.5
O2—C4—C3	120.21 (15)	C6—C11—H11C	109.5
C10—C5—C6	120.67 (15)	H11A—C11—H11C	109.5
C10—C5—N1	120.95 (14)	H11B—C11—H11C	109.5
C6—C5—N1	118.36 (14)	C1—N1—C5	126.65 (14)
C7—C6—C5	117.92 (15)	C1—N1—H1	114.9 (12)
C7—C6—C11	120.05 (15)	C5—N1—H1	118.4 (12)
C5—C6—C11	122.03 (15)	O1—O1—C1	0(10)
C8—C7—C6	120.85 (15)	C4—O2—H2A	113.1 (17)
O1—C1—C2—C3	-3.2 (3)	C6—C7—C8—C11	-179.45 (13)
O1—C1—C2—C3	-3.2 (3)	C7—C8—C9—C10	0.9 (3)
N1—C1—C2—C3	178.4 (2)	C11—C8—C9—C10	-179.48 (13)
C1—C2—C3—C4	-2.4 (4)	C8—C9—C10—C5	-1.1 (3)
C2—C3—C4—O3	-176.3 (2)	C6—C5—C10—C9	0.2 (3)
C2—C3—C4—O2	2.7 (4)	N1—C5—C10—C9	178.60 (15)
C10—C5—C6—C7	0.8 (2)	O1—C1—N1—C5	1.3 (3)
N1—C5—C6—C7	-177.59 (14)	O1—C1—N1—C5	1.3 (3)
C10—C5—C6—C11	-178.68 (16)	C2—C1—N1—C5	179.73 (16)
N1—C5—C6—C11	2.9 (2)	C10—C5—N1—C1	34.9 (3)
C5—C6—C7—C8	-1.0 (2)	C6—C5—N1—C1	-146.67 (17)
C11—C6—C7—C8	178.52 (16)	N1—C1—O1—O1	0.00 (10)
C6—C7—C8—C9	0.1 (3)	C2—C1—O1—O1	0.00 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O3 ⁱ	0.86 (1)	2.10 (1)	2.9556 (19)	174.(2)
O2—H2A...O1	0.92 (1)	1.57 (1)	2.4797 (17)	171 (3)

Symmetry codes: (i) $x+1/2, -y+1/2, -z$.

Fig. 1




Fig. 2

