

1-Benzoyl-3-(4-n-butylphenyl)thiourea

M. Khawar Rauf,^a Masahiro Ebihara,^b Amin Badshah^{a*} and Imtiaz-ud-Din^a

^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan, and ^bDepartment of Chemistry, Faculty of Engineering, Gifu University Yanagido, Gifu 501-1193, Japan

Correspondence e-mail: aminbadshah@yahoo.com


Received 21 November 2011; accepted 1 December 2011

Key indicators: single-crystal X-ray study; $T = 123\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.043; wR factor = 0.103; data-to-parameter ratio = 18.2.

The dihedral angle between the benzoyl and phenyl groups in the title compound, $C_{18}H_{20}N_2OS$, is $30.57(4)^\circ$. The crystal packing is characterized by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. In the crystals, pairs of $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds link the molecules into inversion dimers.

Related literature

For background to our work on the structural chemistry of N,N' -disubstituted thioureas and for related structures, see: Khawar Rauf *et al.* (2009a,b). For bond-length data, see: Allen *et al.* (1987). For a description of the Cambridge Structural Database, see: Allen (2002).



Experimental

Crystal data

$C_{18}H_{20}N_2OS$	$\gamma = 92.700(8)^\circ$
$M_r = 312.42$	$V = 807.9(8)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 4.648(3)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 13.274(8)\text{ \AA}$	$\mu = 0.20\text{ mm}^{-1}$
$c = 13.690(8)\text{ \AA}$	$T = 123\text{ K}$
$\alpha = 106.765(7)^\circ$	$0.40 \times 0.10 \times 0.10\text{ mm}$
$\beta = 90.013(6)^\circ$	

Data collection

Rigaku/MSC Mercury CCD diffractometer	3632 independent reflections
6380 measured reflections	3242 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$	200 parameters
$wR(F^2) = 0.103$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\text{max}} = 0.33\text{ e \AA}^{-3}$
3632 reflections	$\Delta\rho_{\text{min}} = -0.27\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O1	0.88	1.88	2.630 (2)	142
N2—H2 \cdots S1 ⁱ	0.88	2.76	3.550 (2)	151

Symmetry code: (i) $-x + 1, -y, -z + 1$.

Data collection: *CrystalClear* (Molecular Structure Corporation & Rigaku, 2001); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976) and *TEXSAN* (Molecular Structure Corporation & Rigaku, 2004); software used to prepare material for publication: *Yadokari-XG 2009* (Kabuto *et al.*, 2009).

MKR is grateful to the Quaid-i-Azam University, Islamabad, for financial support for a post-doctoral fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5142).

References

- Allen, F. H. (2002). *Acta Cryst. B* **58**, 380–388.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2* pp. S1–19.
- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Kabuto, C., Akine, S., Nemoto, T. & Kwon, E. (2009). *J. Crystallogr. Soc. Jpn.* **51**, 218–224.
- Khawar Rauf, M., Bolte, M. & Badshah, A. (2009a). *Acta Cryst. E* **65**, o177.
- Khawar Rauf, M., Bolte, M. & Badshah, A. (2009b). *Acta Cryst. E* **65**, o240.
- Molecular Structure Corporation & Rigaku (2001). *CrystalClear*. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
- Molecular Structure Corporation & Rigaku (2004). *TEXSAN*. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2012). E68, o120 [doi:10.1107/S1600536811051774]

1-Benzoyl-3-(4-*n*-butylphenyl)thiourea

M. K. Rauf, M. Ebihara, A. Badshah and Imtiaz-ud-Din

Comment

The background to this study has been set out in our previous work for the structural chemistry of *N,N'*-disubstituted thiourea (Khawar Rauf *et al.*, 2009a, 2009b). Herein, as a continuation of these crystallographic studies, the structure of the title compound (I) is described, Fig. 1. Compared to *N*-benzoyl-*N'*-phenylthioureas [Cambridge Structural Database (*Mogul* Version 1.7; Allen, 2002) and (Allen *et al.*, 1987)], the *n*-butyl substitution at C(6) on phenyl ring, implies no significant effect on these bond lengths, and show the molecule to exist in the thione form with typical thiourea C—S and C—O bonds, as well as shortened C—N bond lengths. The dihedral angles to the N(1) C(1)S(1) N(2) C(2)O(1) plane are 23.66 (8) $^{\circ}$ for the ring formed by C(13) to C(18) and 8.16 (9) $^{\circ}$ for the ring formed by C(3) to C(8). An intramolecular N—H···O H-bond is present (Table 1), forming a six-membered ring commonly observed in this class of compounds (Khawar Rauf *et al.*, 2009a, 2009b). In the crystal packing of (I), intermolecular N—H···S H-bonds link the molecules into centrosymmetric dimers (Fig.2).

Experimental

Freshly prepared benzoylisothiocyanate (1.63 g, 10 mmol) was dissolved in acetone (30 ml) and stirred for 30 minutes. Afterwards neat 4-*n*-butylaniline (1.49 g, 10 mmol) was added and the resulting mixture was stirred for 2 h. The reaction mixture was then poured into acidified water and stirred well. The solid product was separated and washed with deionized water and purified by recrystallization from methanol/ 1,1-dichloromethane (1:1 v/v) to give fine crystals of the title compound (I), with an overall yield of 95%.

Refinement

Some strong reflections were saturated in the experimental condition and excluded from the refinement. Hydrogen atoms were included in calculated positions and refined as riding on their parent atom with N—H = 0.88 Å and $U_{\text{iso}}(\text{H}) = 1.2U(\text{N}_{\text{eq}})$, C_{aromatic}—H = 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2U(\text{C}_{\text{eq}})$ or C—H = 0.98–0.99 Å and $U_{\text{iso}}(\text{H}) = 1.5U(\text{C}_{\text{eq}})$, for butyl C atoms.

supplementary materials

Figures




Fig. 1. ORTEP of (I). Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds shown as dashed lines.




Fig. 2. Packing diagram of (I). Hydrogen bonds shown as dashed lines.

1-Benzoyl-3-(4-*n*-butylphenyl)thiourea

Crystal data

C ₁₈ H ₂₀ N ₂ OS	Z = 2
M _r = 312.42	F(000) = 332
Triclinic, P [̄] 1	D _x = 1.284 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71070 Å
a = 4.648 (3) Å	Cell parameters from 2558 reflections
b = 13.274 (8) Å	θ = 3.1–27.5°
c = 13.690 (8) Å	μ = 0.20 mm ⁻¹
α = 106.765 (7)°	T = 123 K
β = 90.013 (6)°	Needle like, colorless

$\gamma = 92.700(8)^\circ$ $0.40 \times 0.10 \times 0.10 \text{ mm}$
 $V = 807.9(8) \text{ \AA}^3$

Data collection

Rigaku/MSC Mercury CCD diffractometer	3242 reflections with $I > 2\sigma(I)$
Radiation source: Rotating Anode	$R_{\text{int}} = 0.043$
Graphite Monochromator	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.1^\circ$
Detector resolution: 14.6199 pixels mm^{-1}	$h = -6 \rightarrow 3$
dtintegrate.ref scans	$k = -17 \rightarrow 14$
6380 measured reflections	$l = -13 \rightarrow 17$
3632 independent reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.043$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.103$	H-atom parameters constrained
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0399P)^2 + 0.4308P]$ where $P = (F_o^2 + 2F_c^2)/3$
3632 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
200 parameters	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.27 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4183 (3)	0.19623 (11)	0.53711 (10)	0.0172 (3)
S1	0.33263 (9)	0.11726 (3)	0.60881 (3)	0.02424 (12)
N1	0.3191 (3)	0.29075 (10)	0.54395 (9)	0.0180 (3)
H1	0.3862	0.3194	0.4975	0.022*
N2	0.6153 (3)	0.16024 (10)	0.45859 (9)	0.0178 (3)

supplementary materials

H2	0.6907	0.0999	0.4556	0.021*
C2	0.7060 (3)	0.20705 (12)	0.38572 (11)	0.0190 (3)
O1	0.6324 (3)	0.29449 (9)	0.38503 (9)	0.0271 (3)
C3	0.1241 (3)	0.35335 (11)	0.61301 (11)	0.0165 (3)
C4	-0.0402 (3)	0.32010 (12)	0.68386 (12)	0.0212 (3)
H4	-0.0285	0.2506	0.6890	0.025*
C5	-0.2219 (3)	0.39021 (13)	0.74715 (12)	0.0217 (3)
H5	-0.3322	0.3674	0.7959	0.026*
C6	-0.2477 (3)	0.49186 (12)	0.74157 (11)	0.0190 (3)
C7	-0.0851 (3)	0.52320 (12)	0.66869 (11)	0.0210 (3)
H7	-0.1008	0.5922	0.6624	0.025*
C8	0.0986 (3)	0.45522 (12)	0.60552 (11)	0.0201 (3)
H8	0.2083	0.4781	0.5566	0.024*
C9	-0.4331 (3)	0.56761 (13)	0.81559 (11)	0.0227 (3)
H9B	-0.6086	0.5291	0.8295	0.027*
H9A	-0.4937	0.6221	0.7845	0.027*
C10	-0.2717 (3)	0.62093 (13)	0.91620 (11)	0.0218 (3)
H10B	-0.2076	0.5659	0.9459	0.026*
H10A	-0.0976	0.6598	0.9017	0.026*
C11	-0.4509 (4)	0.69711 (14)	0.99458 (12)	0.0264 (3)
H11B	-0.5213	0.7509	0.9644	0.032*
H11A	-0.6208	0.6580	1.0117	0.032*
C12	-0.2805 (4)	0.75183 (15)	1.09179 (13)	0.0329 (4)
H12C	-0.2089	0.6989	1.1217	0.049*
H12A	-0.4055	0.7983	1.1406	0.049*
H12B	-0.1174	0.7935	1.0757	0.049*
C13	0.8938 (3)	0.14395 (12)	0.30440 (11)	0.0185 (3)
C14	0.9027 (3)	0.16959 (13)	0.21251 (12)	0.0247 (3)
H14	0.7984	0.2268	0.2049	0.030*
C15	1.0633 (4)	0.11182 (14)	0.13232 (12)	0.0281 (4)
H15	1.0667	0.1287	0.0695	0.034*
C16	1.2191 (4)	0.02936 (14)	0.14376 (12)	0.0275 (4)
H16	1.3290	-0.0102	0.0887	0.033*
C17	1.2152 (3)	0.00445 (13)	0.23512 (12)	0.0245 (3)
H17	1.3228	-0.0520	0.2428	0.029*
C18	1.0537 (3)	0.06196 (12)	0.31573 (11)	0.0190 (3)
H18	1.0526	0.0452	0.3787	0.023*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0203 (7)	0.0183 (7)	0.0122 (6)	0.0008 (5)	0.0007 (5)	0.0033 (5)
S1	0.0384 (2)	0.01744 (19)	0.0190 (2)	0.00574 (15)	0.01041 (15)	0.00803 (14)
N1	0.0224 (6)	0.0170 (6)	0.0155 (6)	0.0029 (5)	0.0043 (5)	0.0061 (5)
N2	0.0225 (6)	0.0157 (6)	0.0157 (6)	0.0036 (5)	0.0035 (5)	0.0050 (5)
C2	0.0219 (7)	0.0185 (7)	0.0172 (7)	0.0005 (5)	0.0018 (5)	0.0062 (6)
O1	0.0365 (6)	0.0215 (6)	0.0274 (6)	0.0072 (5)	0.0122 (5)	0.0130 (5)
C3	0.0179 (7)	0.0171 (7)	0.0136 (6)	0.0014 (5)	0.0001 (5)	0.0028 (5)

C4	0.0213 (7)	0.0205 (7)	0.0227 (7)	0.0013 (6)	0.0026 (6)	0.0078 (6)
C5	0.0209 (7)	0.0258 (8)	0.0189 (7)	0.0008 (6)	0.0038 (6)	0.0073 (6)
C6	0.0165 (7)	0.0227 (8)	0.0157 (7)	0.0019 (5)	-0.0034 (5)	0.0019 (6)
C7	0.0256 (8)	0.0178 (7)	0.0196 (7)	0.0039 (6)	-0.0004 (6)	0.0052 (6)
C8	0.0242 (7)	0.0214 (7)	0.0163 (7)	0.0016 (6)	0.0021 (5)	0.0078 (6)
C9	0.0202 (7)	0.0267 (8)	0.0188 (7)	0.0056 (6)	-0.0002 (6)	0.0020 (6)
C10	0.0198 (7)	0.0264 (8)	0.0172 (7)	0.0041 (6)	0.0012 (6)	0.0029 (6)
C11	0.0261 (8)	0.0318 (9)	0.0186 (8)	0.0060 (7)	0.0029 (6)	0.0023 (7)
C12	0.0399 (10)	0.0363 (10)	0.0180 (8)	0.0041 (8)	0.0022 (7)	0.0004 (7)
C13	0.0203 (7)	0.0189 (7)	0.0163 (7)	-0.0016 (5)	0.0020 (5)	0.0055 (6)
C14	0.0300 (8)	0.0260 (8)	0.0214 (8)	0.0048 (6)	0.0040 (6)	0.0116 (6)
C15	0.0361 (9)	0.0335 (9)	0.0161 (7)	0.0021 (7)	0.0052 (6)	0.0091 (7)
C16	0.0299 (8)	0.0286 (9)	0.0208 (8)	0.0022 (7)	0.0077 (6)	0.0020 (7)
C17	0.0241 (8)	0.0237 (8)	0.0250 (8)	0.0026 (6)	0.0032 (6)	0.0058 (6)
C18	0.0203 (7)	0.0203 (7)	0.0165 (7)	-0.0014 (6)	0.0005 (5)	0.0058 (6)

Geometric parameters (Å, °)

C1—N1	1.335 (2)	C9—H9A	0.9900
C1—N2	1.4028 (19)	C10—C11	1.523 (2)
C1—S1	1.6659 (16)	C10—H10B	0.9900
N1—C3	1.4228 (18)	C10—H10A	0.9900
N1—H1	0.8800	C11—C12	1.521 (2)
N2—C2	1.3755 (19)	C11—H11B	0.9900
N2—H2	0.8800	C11—H11A	0.9900
C2—O1	1.2282 (19)	C12—H12C	0.9800
C2—C13	1.495 (2)	C12—H12A	0.9800
C3—C4	1.391 (2)	C12—H12B	0.9800
C3—C8	1.396 (2)	C13—C18	1.391 (2)
C4—C5	1.393 (2)	C13—C14	1.395 (2)
C4—H4	0.9500	C14—C15	1.386 (2)
C5—C6	1.384 (2)	C14—H14	0.9500
C5—H5	0.9500	C15—C16	1.386 (2)
C6—C7	1.397 (2)	C15—H15	0.9500
C6—C9	1.508 (2)	C16—C17	1.383 (2)
C7—C8	1.383 (2)	C16—H16	0.9500
C7—H7	0.9500	C17—C18	1.390 (2)
C8—H8	0.9500	C17—H17	0.9500
C9—C10	1.532 (2)	C18—H18	0.9500
C9—H9B	0.9900		
N1—C1—N2	114.78 (12)	C11—C10—C9	113.76 (13)
N1—C1—S1	127.89 (11)	C11—C10—H10B	108.8
N2—C1—S1	117.32 (11)	C9—C10—H10B	108.8
C1—N1—C3	131.41 (13)	C11—C10—H10A	108.8
C1—N1—H1	114.3	C9—C10—H10A	108.8
C3—N1—H1	114.3	H10B—C10—H10A	107.7
C2—N2—C1	128.45 (13)	C12—C11—C10	112.35 (14)
C2—N2—H2	115.8	C12—C11—H11B	109.1
C1—N2—H2	115.8	C10—C11—H11B	109.1

supplementary materials

O1—C2—N2	122.56 (14)	C12—C11—H11A	109.1
O1—C2—C13	121.20 (13)	C10—C11—H11A	109.1
N2—C2—C13	116.21 (13)	H11B—C11—H11A	107.9
C4—C3—C8	119.34 (13)	C11—C12—H12C	109.5
C4—C3—N1	125.23 (14)	C11—C12—H12A	109.5
C8—C3—N1	115.43 (13)	H12C—C12—H12A	109.5
C3—C4—C5	119.06 (14)	C11—C12—H12B	109.5
C3—C4—H4	120.5	H12C—C12—H12B	109.5
C5—C4—H4	120.5	H12A—C12—H12B	109.5
C6—C5—C4	122.41 (14)	C18—C13—C14	119.47 (14)
C6—C5—H5	118.8	C18—C13—C2	123.58 (13)
C4—C5—H5	118.8	C14—C13—C2	116.94 (14)
C5—C6—C7	117.70 (13)	C15—C14—C13	120.13 (15)
C5—C6—C9	120.87 (14)	C15—C14—H14	119.9
C7—C6—C9	121.34 (14)	C13—C14—H14	119.9
C8—C7—C6	120.92 (14)	C14—C15—C16	120.04 (15)
C8—C7—H7	119.5	C14—C15—H15	120.0
C6—C7—H7	119.5	C16—C15—H15	120.0
C7—C8—C3	120.55 (14)	C17—C16—C15	120.22 (15)
C7—C8—H8	119.7	C17—C16—H16	119.9
C3—C8—H8	119.7	C15—C16—H16	119.9
C6—C9—C10	111.53 (12)	C16—C17—C18	119.98 (15)
C6—C9—H9B	109.3	C16—C17—H17	120.0
C10—C9—H9B	109.3	C18—C17—H17	120.0
C6—C9—H9A	109.3	C17—C18—C13	120.14 (14)
C10—C9—H9A	109.3	C17—C18—H18	119.9
H9B—C9—H9A	108.0	C13—C18—H18	119.9
N2—C1—N1—C3	−178.89 (13)	N1—C3—C8—C7	179.89 (13)
S1—C1—N1—C3	2.0 (2)	C5—C6—C9—C10	83.10 (18)
N1—C1—N2—C2	−3.7 (2)	C7—C6—C9—C10	−93.57 (17)
S1—C1—N2—C2	175.53 (12)	C6—C9—C10—C11	−179.06 (14)
C1—N2—C2—O1	4.8 (2)	C9—C10—C11—C12	−177.71 (15)
C1—N2—C2—C13	−173.38 (13)	O1—C2—C13—C18	159.84 (15)
C1—N1—C3—C4	−9.2 (2)	N2—C2—C13—C18	−22.0 (2)
C1—N1—C3—C8	171.86 (14)	O1—C2—C13—C14	−20.9 (2)
C8—C3—C4—C5	−1.3 (2)	N2—C2—C13—C14	157.27 (14)
N1—C3—C4—C5	179.74 (14)	C18—C13—C14—C15	1.8 (2)
C3—C4—C5—C6	0.7 (2)	C2—C13—C14—C15	−177.51 (15)
C4—C5—C6—C7	0.5 (2)	C13—C14—C15—C16	−1.0 (3)
C4—C5—C6—C9	−176.28 (14)	C14—C15—C16—C17	−0.1 (3)
C5—C6—C7—C8	−1.0 (2)	C15—C16—C17—C18	0.2 (3)
C9—C6—C7—C8	175.77 (14)	C16—C17—C18—C13	0.6 (2)
C6—C7—C8—C3	0.3 (2)	C14—C13—C18—C17	−1.6 (2)
C4—C3—C8—C7	0.9 (2)	C2—C13—C18—C17	177.65 (14)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O1	0.88	1.88	2.630 (2)	142.


N2—H2···S1ⁱ

0.88

2.76

3.550 (2)

151.

Symmetry codes: (i) $-x+1, -y, -z+1$.**Fig. 1**

supplementary materials

Fig. 2

