

2,4,5-Tris(pyridin-4-yl)-1H-imidazole monohydrate

Shen-Tang Wang, Guang-Bo Che, Chun-Bo Liu,* Xing Wang and Ling Liu

School of Chemistry and Chemical Engineering, Jiangsu University, Zhenjiang 212013, People's Republic of China

Correspondence e-mail: guangbocheujs@yahoo.com.cn


Received 4 November 2011; accepted 9 December 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.058; wR factor = 0.125; data-to-parameter ratio = 13.2.

In the crystal structure of the title compound, $\text{C}_{18}\text{H}_{13}\text{N}_5 \cdot \text{H}_2\text{O}$, adjacent molecules are linked by $\text{O}-\text{H} \cdots \text{N}$ and $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds, generating a chain propagating along [001].

Related literature

For the use of 2,4,5-tri(4-pyridyl)imidazole in the construction of metal-organic coordination polymers, see: Wang *et al.* (2009); Liang *et al.* (2009). For related structures, see: Jiang & Hou (2011); Li (2011); Li & Xia (2011). For the preparation, see: Proskurnina *et al.* (2002).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{13}\text{N}_5 \cdot \text{H}_2\text{O}$

$M_r = 317.35$

Triclinic, $P\bar{1}$
 $a = 8.1510$ (16) Å
 $b = 9.5210$ (19) Å
 $c = 11.506$ (2) Å
 $\alpha = 103.80$ (3)°
 $\beta = 105.64$ (3)°
 $\gamma = 101.03$ (3)°

$V = 803.3$ (4) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 293$ K
 $0.35 \times 0.25 \times 0.2$ mm

Data collection

Bruker SMART diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2002)
 $T_{\min} = 0.970$, $T_{\max} = 1.000$

7510 measured reflections
 2912 independent reflections
 1792 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.058$
 $wR(F^2) = 0.125$
 $S = 1.02$
 2876 reflections

218 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O1}-\text{H1A} \cdots \text{N5}^i$	0.85	1.99	2.843 (3)	177
$\text{N1}-\text{H1} \cdots \text{O1}$	0.89	1.85	2.741 (3)	176
$\text{O1}-\text{H1B} \cdots \text{N4}^{ii}$	0.85	1.94	2.787 (3)	172

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $x, y, z + 1$.

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008) and DIAMOND (Brandenburg, 1998); software used to prepare material for publication: SHELXTL.

The authors thank Jiangsu University for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZJ2036).

References

- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2002). *SADABS, SAINT and SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Jiang, Y.-K. & Hou, G.-G. (2011). *Acta Cryst.* **E67**, o3073.
 Li, H.-D. (2011). *Acta Cryst.* **E67**, o3156.
 Li, C.-R. & Xia, Z.-Q. (2011). *Acta Cryst.* **E67**, o2481.
 Liang, X.-Q., Zhou, X.-H., Chen, C., Xiao, H.-P., Li, Y.-Z., Zuo, J.-L. & You, X.-Z. (2009). *Cryst. Growth Des.* **9**, 1041–1053.
 Proskurnina, M. V., Lozinskaya, N. A., Tkachenko, S. E. & Zefirov, N. S. (2002). *Russ. J. Org. Chem.* **38**, 1149–1153.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wang, L., Gu, W., Deng, J.-X., Liu, M.-L., Xu, N. & Liu, X. (2009). *Z. Anorg. Allg. Chem.* **635**, 1124–1128.

supplementary materials

Acta Cryst. (2012). E68, o130 [doi:10.1107/S1600536811053013]

2,4,5-Tris(pyridin-4-yl)-1*H*-imidazole monohydrate

S.-T. Wang, G.-B. Che, C.-B. Liu, X. Wang and L. Liu

Comment

In the 2,4,5-tris(4-pyridyl)imidazole three pyridyl groups (pyridyl ring B (C4-C8 and N3), C (C9-C13 and N4), D (C14-C18 and N5)) are directly connected with the imidazole ring A (C1-C3, N1 and N2). The dihedral angles between the mean planes of pyridyl ring B and imidazole ring A, pyridyl ring C and imidazole ring A, and pyridyl ring D and imidazole ring A are 9.1 (7)°, 21.5 (5)°, 45.5 (1)°, respectively.

We report herein on the crystal structure of the title compound (Fig. 1). In the crystal lattice the molecules are linked by O—H...N and N—H...O hydrogen bonds (Jiang *et al.* 2011; Li *et al.* 2011; Li 2011) interactions to generate a one-dimensional double chain structure (Fig. 2).

Experimental

The 2,4,5-tris(4-pyridyl)imidazole was prepared by the method reported in the literature (Proskurnina *et al.* 2002). A mixture of 2,4,5-tris(4-pyridyl)imidazole (0.030 g, 0.1 mmol), 2 drops of 1 mol/L HCl and water (10 mL) was placed in a 25 mL Teflon-lined autoclave and heated for 3 d at 433 K under autogenous pressure. Upon cooling and opening the bomb, colourless block-shaped crystals were obtained, then washed with water and dried in air.

Refinement

All H atoms on C atoms were positioned geometrically and refined as riding atoms, with (C—H = 0.93 Å) and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$. The hydrogen atoms of water molecules were located in a difference Fourier map, and were refined with suitable O—H distance restraint; $U_{\text{iso}} = 1.5 U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. All H atoms are presented as a small spheres of arbitrary radius.




Fig. 2. The one-dimensional supermolecular structure linked by the hydrogen bonds.

2,4,5-Tris(pyridin-4-yl)-1H-imidazole monohydrate

Crystal data

$C_{18}H_{13}N_5 \cdot H_2O$	$Z = 2$
$M_r = 317.35$	$F(000) = 332$
Triclinic, $P\bar{1}$	$D_x = 1.312 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 8.1510 (16) \text{ \AA}$	Cell parameters from 3107 reflections
$b = 9.5210 (19) \text{ \AA}$	$\theta = 3.0\text{--}25.2^\circ$
$c = 11.506 (2) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\alpha = 103.80 (3)^\circ$	$T = 293 \text{ K}$
$\beta = 105.64 (3)^\circ$	Prism, colourless
$\gamma = 101.03 (3)^\circ$	$0.35 \times 0.25 \times 0.2 \text{ mm}$
$V = 803.3 (4) \text{ \AA}^3$	

Data collection

Bruker SMART diffractometer	2912 independent reflections
Radiation source: fine-focus sealed tube graphite	1792 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.040$
Absorption correction: multi-scan (SADABS; Bruker, 2002)	$\theta_{\text{max}} = 25.2^\circ$, $\theta_{\text{min}} = 3.0^\circ$
$T_{\text{min}} = 0.970$, $T_{\text{max}} = 1.000$	$h = -9 \rightarrow 9$
7510 measured reflections	$k = -11 \rightarrow 11$
	$l = -13 \rightarrow 13$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.058$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.125$	H-atom parameters constrained
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0171P)^2 + 0.605P]$
2876 reflections	where $P = (F_o^2 + 2F_c^2)/3$
218 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.19 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.17 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.2548 (3)	0.5006 (2)	0.26415 (19)	0.0544 (6)
H1	0.2800	0.5437	0.3469	0.082*
N2	0.1414 (3)	0.3416 (2)	0.06806 (19)	0.0542 (6)
N3	-0.0531 (4)	0.0012 (3)	0.3185 (2)	0.0738 (7)
N4	0.2559 (4)	0.5283 (3)	-0.2901 (2)	0.0749 (7)
N5	0.5288 (4)	1.0560 (3)	0.3379 (3)	0.0764 (8)
C1	0.1603 (3)	0.3582 (3)	0.1895 (2)	0.0518 (6)
C2	0.2265 (3)	0.4798 (3)	0.0649 (2)	0.0523 (6)
C3	0.2960 (3)	0.5811 (3)	0.1858 (2)	0.0530 (7)
C4	0.0870 (3)	0.2404 (3)	0.2365 (2)	0.0520 (6)
C5	0.1271 (4)	0.2521 (3)	0.3637 (3)	0.0671 (8)
H5A	0.2026	0.3399	0.4252	0.080*
C6	0.0536 (4)	0.1317 (3)	0.3987 (3)	0.0739 (9)
H6A	0.0812	0.1433	0.4849	0.089*
C7	-0.0932 (4)	-0.0074 (3)	0.1966 (3)	0.0754 (9)
H7A	-0.1703	-0.0962	0.1374	0.091*
C8	-0.0284 (4)	0.1062 (3)	0.1517 (3)	0.0668 (8)
H8A	-0.0621	0.0926	0.0650	0.080*
C9	0.2386 (3)	0.4986 (3)	-0.0558 (2)	0.0527 (6)
C10	0.1214 (4)	0.4006 (3)	-0.1700 (2)	0.0594 (7)
H10A	0.0342	0.3211	-0.1709	0.071*
C11	0.1334 (4)	0.4203 (3)	-0.2821 (3)	0.0708 (8)
H11A	0.0505	0.3537	-0.3572	0.085*
C12	0.3716 (4)	0.6201 (4)	-0.1801 (3)	0.0775 (9)
H12A	0.4600	0.6962	-0.1826	0.093*
C13	0.3698 (4)	0.6105 (3)	-0.0626 (3)	0.0681 (8)
H13A	0.4552	0.6778	0.0110	0.082*
C14	0.3805 (4)	0.7431 (3)	0.2368 (2)	0.0549 (7)
C15	0.3141 (4)	0.8416 (3)	0.1767 (3)	0.0675 (8)
H15A	0.2185	0.8049	0.1015	0.081*
C16	0.3925 (4)	0.9939 (3)	0.2305 (3)	0.0754 (9)
H16A	0.3470	1.0577	0.1890	0.090*
C17	0.5915 (4)	0.9608 (3)	0.3939 (3)	0.0735 (9)

supplementary materials

H17A	0.6879	1.0009	0.4685	0.088*
C18	0.5219 (4)	0.8055 (3)	0.3480 (3)	0.0650 (8)
H18A	0.5701	0.7446	0.3918	0.078*
O1	0.3155 (4)	0.6367 (2)	0.51602 (18)	0.1058 (10)
H1A	0.3597	0.7298	0.5581	0.159*
H1B	0.3044	0.5992	0.5745	0.159*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0703 (14)	0.0475 (12)	0.0430 (12)	0.0090 (10)	0.0198 (10)	0.0139 (9)
N2	0.0711 (15)	0.0475 (12)	0.0439 (12)	0.0088 (10)	0.0215 (11)	0.0167 (9)
N3	0.0983 (19)	0.0555 (14)	0.0683 (17)	0.0082 (13)	0.0323 (15)	0.0249 (13)
N4	0.102 (2)	0.0736 (16)	0.0558 (15)	0.0144 (15)	0.0375 (15)	0.0265 (13)
N5	0.0884 (19)	0.0549 (15)	0.0791 (18)	0.0055 (14)	0.0317 (16)	0.0152 (14)
C1	0.0689 (17)	0.0422 (13)	0.0430 (14)	0.0118 (12)	0.0187 (13)	0.0129 (11)
C2	0.0652 (16)	0.0482 (14)	0.0426 (14)	0.0091 (12)	0.0199 (12)	0.0153 (11)
C3	0.0660 (17)	0.0482 (14)	0.0446 (14)	0.0091 (12)	0.0201 (13)	0.0171 (11)
C4	0.0680 (17)	0.0428 (13)	0.0470 (15)	0.0122 (12)	0.0209 (13)	0.0174 (11)
C5	0.097 (2)	0.0510 (15)	0.0484 (16)	0.0064 (15)	0.0245 (15)	0.0162 (13)
C6	0.111 (3)	0.0599 (18)	0.0564 (18)	0.0163 (17)	0.0360 (18)	0.0236 (15)
C7	0.094 (2)	0.0519 (17)	0.068 (2)	-0.0007 (15)	0.0184 (17)	0.0212 (15)
C8	0.088 (2)	0.0522 (16)	0.0527 (16)	0.0074 (15)	0.0191 (15)	0.0174 (13)
C9	0.0659 (17)	0.0511 (14)	0.0462 (14)	0.0145 (12)	0.0225 (13)	0.0204 (12)
C10	0.0738 (18)	0.0566 (16)	0.0473 (15)	0.0113 (14)	0.0228 (14)	0.0169 (13)
C11	0.090 (2)	0.0732 (19)	0.0474 (17)	0.0160 (17)	0.0259 (16)	0.0173 (14)
C12	0.097 (2)	0.071 (2)	0.068 (2)	0.0054 (17)	0.0402 (19)	0.0273 (17)
C13	0.079 (2)	0.0656 (18)	0.0517 (17)	0.0000 (15)	0.0251 (15)	0.0158 (14)
C14	0.0670 (17)	0.0486 (14)	0.0492 (15)	0.0082 (12)	0.0251 (13)	0.0148 (12)
C15	0.078 (2)	0.0519 (16)	0.0657 (19)	0.0083 (14)	0.0172 (16)	0.0212 (14)
C16	0.087 (2)	0.0536 (17)	0.086 (2)	0.0151 (16)	0.0285 (19)	0.0265 (16)
C17	0.080 (2)	0.0604 (18)	0.0624 (19)	-0.0019 (16)	0.0216 (16)	0.0061 (15)
C18	0.077 (2)	0.0548 (16)	0.0552 (17)	0.0068 (14)	0.0185 (15)	0.0156 (13)
O1	0.187 (3)	0.0597 (13)	0.0500 (12)	-0.0125 (14)	0.0451 (14)	0.0087 (10)

Geometric parameters (\AA , $^\circ$)

N1—C1	1.363 (3)	C7—H7A	0.9300
N1—C3	1.381 (3)	C8—H8A	0.9300
N1—H1	0.8907	C9—C10	1.381 (3)
N2—C1	1.331 (3)	C9—C13	1.390 (3)
N2—C2	1.379 (3)	C10—C11	1.373 (4)
N3—C6	1.326 (4)	C10—H10A	0.9300
N3—C7	1.329 (4)	C11—H11A	0.9300
N4—C11	1.329 (4)	C12—C13	1.381 (4)
N4—C12	1.330 (4)	C12—H12A	0.9300
N5—C16	1.330 (4)	C13—H13A	0.9300
N5—C17	1.333 (4)	C14—C18	1.378 (4)
C1—C4	1.454 (3)	C14—C15	1.397 (4)

C2—C3	1.383 (3)	C15—C16	1.378 (4)
C2—C9	1.469 (3)	C15—H15A	0.9300
C3—C14	1.463 (3)	C16—H16A	0.9300
C4—C5	1.383 (3)	C17—C18	1.390 (4)
C4—C8	1.385 (3)	C17—H17A	0.9300
C5—C6	1.387 (4)	C18—H18A	0.9300
C5—H5A	0.9300	O1—H1A	0.8543
C6—H6A	0.9300	O1—H1B	0.8500
C7—C8	1.382 (4)		
C1—N1—C3	107.5 (2)	C10—C9—C13	116.4 (2)
C1—N1—H1	130.1	C10—C9—C2	120.8 (2)
C3—N1—H1	122.1	C13—C9—C2	122.8 (2)
C1—N2—C2	105.6 (2)	C11—C10—C9	120.2 (3)
C6—N3—C7	115.0 (2)	C11—C10—H10A	119.9
C11—N4—C12	115.6 (2)	C9—C10—H10A	119.9
C16—N5—C17	116.0 (3)	N4—C11—C10	124.1 (3)
N2—C1—N1	111.3 (2)	N4—C11—H11A	117.9
N2—C1—C4	124.4 (2)	C10—C11—H11A	117.9
N1—C1—C4	124.3 (2)	N4—C12—C13	124.6 (3)
N2—C2—C3	110.2 (2)	N4—C12—H12A	117.7
N2—C2—C9	119.8 (2)	C13—C12—H12A	117.7
C3—C2—C9	130.0 (2)	C12—C13—C9	119.1 (3)
N1—C3—C2	105.3 (2)	C12—C13—H13A	120.5
N1—C3—C14	120.4 (2)	C9—C13—H13A	120.5
C2—C3—C14	134.0 (2)	C18—C14—C15	117.3 (2)
C5—C4—C8	116.3 (2)	C18—C14—C3	122.0 (2)
C5—C4—C1	123.8 (2)	C15—C14—C3	120.6 (2)
C8—C4—C1	119.9 (2)	C16—C15—C14	118.9 (3)
C4—C5—C6	119.4 (3)	C16—C15—H15A	120.5
C4—C5—H5A	120.3	C14—C15—H15A	120.5
C6—C5—H5A	120.3	N5—C16—C15	124.5 (3)
N3—C6—C5	124.9 (3)	N5—C16—H16A	117.7
N3—C6—H6A	117.5	C15—C16—H16A	117.7
C5—C6—H6A	117.5	N5—C17—C18	124.1 (3)
N3—C7—C8	124.6 (3)	N5—C17—H17A	117.9
N3—C7—H7A	117.7	C18—C17—H17A	117.9
C8—C7—H7A	117.7	C14—C18—C17	119.1 (3)
C7—C8—C4	119.7 (3)	C14—C18—H18A	120.4
C7—C8—H8A	120.1	C17—C18—H18A	120.4
C4—C8—H8A	120.1	H1A—O1—H1B	100.9
C2—N2—C1—N1	-0.5 (3)	N2—C2—C9—C10	21.5 (4)
C2—N2—C1—C4	178.6 (3)	C3—C2—C9—C10	-161.6 (3)
C3—N1—C1—N2	1.3 (3)	N2—C2—C9—C13	-156.1 (3)
C3—N1—C1—C4	-177.8 (3)	C3—C2—C9—C13	20.8 (5)
C1—N2—C2—C3	-0.5 (3)	C13—C9—C10—C11	-2.7 (4)
C1—N2—C2—C9	176.9 (2)	C2—C9—C10—C11	179.5 (3)
C1—N1—C3—C2	-1.6 (3)	C12—N4—C11—C10	0.2 (5)
C1—N1—C3—C14	173.1 (2)	C9—C10—C11—N4	1.6 (5)

supplementary materials

N2—C2—C3—N1	1.3 (3)	C11—N4—C12—C13	-0.8 (5)
C9—C2—C3—N1	-175.8 (3)	N4—C12—C13—C9	-0.5 (5)
N2—C2—C3—C14	-172.3 (3)	C10—C9—C13—C12	2.2 (4)
C9—C2—C3—C14	10.6 (5)	C2—C9—C13—C12	179.9 (3)
N2—C1—C4—C5	170.9 (3)	N1—C3—C14—C18	45.8 (4)
N1—C1—C4—C5	-10.2 (4)	C2—C3—C14—C18	-141.4 (3)
N2—C1—C4—C8	-8.3 (4)	N1—C3—C14—C15	-131.1 (3)
N1—C1—C4—C8	170.7 (3)	C2—C3—C14—C15	41.7 (5)
C8—C4—C5—C6	0.8 (4)	C18—C14—C15—C16	0.2 (4)
C1—C4—C5—C6	-178.4 (3)	C3—C14—C15—C16	177.3 (3)
C7—N3—C6—C5	-2.3 (5)	C17—N5—C16—C15	0.7 (5)
C4—C5—C6—N3	1.1 (5)	C14—C15—C16—N5	-0.3 (5)
C6—N3—C7—C8	1.7 (5)	C16—N5—C17—C18	-0.9 (5)
N3—C7—C8—C4	0.1 (5)	C15—C14—C18—C17	-0.4 (4)
C5—C4—C8—C7	-1.3 (4)	C3—C14—C18—C17	-177.4 (3)
C1—C4—C8—C7	177.9 (3)	N5—C17—C18—C14	0.8 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1A \cdots N5 ⁱ	0.85	1.99	2.843 (3)	177.
N1—H1 \cdots O1	0.89	1.85	2.741 (3)	176.
O1—H1B \cdots N4 ⁱⁱ	0.85	1.94	2.787 (3)	172.

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $x, y, z+1$.

Fig. 1




Fig. 2

