

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Chloro-*N*-(4-chloro-3-iodophenyl)-4-(methylsulfonyl)benzamide


Dao-Cai Wang, Hang Song, Chang Yang, Wen-Cai Huang and Shun Yao*

Department of Pharmaceutical and Biological Engineering, College of Chemical Engineering, Sichuan University, Chengdu 610065, People's Republic of China
Correspondence e-mail: cusack@scu.edu.cn

Received 10 December 2011; accepted 13 December 2011

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.028; wR factor = 0.067; data-to-parameter ratio = 16.6.In the title compound, $\text{C}_{14}\text{H}_{10}\text{Cl}_2\text{INO}_3\text{S}$, the dihedral angle between the benzene rings is 52.13 (10)°. In the crystal, the components are linked by pairs of $\text{N}-\text{H}\cdots\text{O}(\text{sulfonyl})$ hydrogen bonds into centrosymmetric dimers.

Related literature

For background to benzamides, see: Castanedo *et al.* (2010); Tremblay *et al.* (2010); Mahindroo *et al.* (2010). For the preparation of the title compound, see: Robarge *et al.* (2009).

Experimental

Crystal data

 $\text{C}_{14}\text{H}_{10}\text{Cl}_2\text{INO}_3\text{S}$
 $M_r = 470.09$
 Triclinic, $P\bar{1}$
 $a = 8.8694$ (6) Å
 $b = 10.3837$ (8) Å
 $c = 10.4288$ (5) Å
 $\alpha = 103.862$ (5)°
 $\beta = 96.452$ (5)°

 $\gamma = 114.949$ (7)°
 $V = 820.25$ (9) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 2.41$ mm⁻¹
 $T = 290$ K
 $0.30 \times 0.30 \times 0.25$ mm

Data collection

Oxford Diffraction Xcalibur Eos diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2006)
 $T_{\min} = 0.887$, $T_{\max} = 1.000$ 6770 measured reflections
3323 independent reflections
2957 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.067$
 $S = 1.07$
3323 reflections200 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.43$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.55$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O2}^i$	0.86	2.15	2.991 (3)	167

Symmetry code: (i) $-x, -y + 1, -z + 3$.Data collection: *CrysAlis PRO* (Oxford Diffraction, 2006); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *OLEX2*.

We thank the Analytical and Testing Center of Sichuan University for the X-ray measurements.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5034).

References

- Castanedo, G. M., *et al.* (2010). *Bioorg. Med. Chem. Lett.* **20**, 6748–6753.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Mahindroo, N., Connelly, M. C., Punchihewa, C., Yang, L., Yan, B. & Fujii, N. (2010). *Bioorg. Med. Chem.* **18**, 4801–4811.
- Oxford Diffraction (2006). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Robarge, K. D., *et al.* (2009). *Bioorg. Med. Chem. Lett.* **19**, 5576–5581.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tremblay, M. R., McGovern, K., Read, M. A. & Castro, A. C. (2010). *Curr. Opin. Chem. Biol.* **14**, 428–435.

supplementary materials

Acta Cryst. (2012). E68, o189 [doi:10.1107/S1600536811053633]

2-Chloro-*N*-(4-chloro-3-iodophenyl)-4-(methylsulfonyl)benzamide

D.-C. Wang, H. Song, C. Yang, W.-C. Huang and S. Yao

Comment

The benzamide moiety is the constituent of many biologically significant compounds including anticancer compounds (Castanedo *et al.*, 2010; Tremblay *et al.*, 2010; Mahindroo *et al.*, 2010). The title compound is one of the key intermediates in our synthetic investigations of anticancer drugs. In this paper, we synthesized the title compound and report its crystal structure, Fig. 1. The dihedral angle between the two benzene rings is 52.13 (10)°. In the crystal, the molecules are connected *via* intermolecular N—H···O hydrogen bonds to form a centrosymmetric dimer, in which the amide H atom acts as a donor and the sulfonyl-O atom act as an acceptor, Table 1.

Experimental

The title compound was prepared by a method similar to that of Robarge *et al.* (2009). Crystals were obtained by slow evaporation from its methanol-acetic acid (5:2) solution.

Refinement

All H atoms were positioned geometrically (C—H = 0.93–0.96 Å; N—H = 0.86 Å) and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent})$.

Figures




Fig. 1. The molecular structure of the title compound, with displacement ellipsoids drawn at the 30% probability level.

2-Chloro-*N*-(4-chloro-3-iodophenyl)-4-(methylsulfonyl)benzamide

Crystal data

C₁₄H₁₀Cl₂INO₃S

$M_r = 470.09$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 8.8694$ (6) Å

$b = 10.3837$ (8) Å

$c = 10.4288$ (5) Å

$\alpha = 103.862$ (5)°

$\beta = 96.452$ (5)°

$Z = 2$

$F(000) = 456$

$D_x = 1.903$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.7107$ Å

Cell parameters from 2872 reflections

$\theta = 2.9$ – 26.3 °

$\mu = 2.41$ mm⁻¹

$T = 290$ K

Block, colourless

supplementary materials

$\gamma = 114.949 (7)^\circ$ $0.30 \times 0.30 \times 0.25$ mm
 $V = 820.25 (9) \text{ \AA}^3$

Data collection

Oxford Diffraction Xcalibur Eos diffractometer	3323 independent reflections
Radiation source: Enhance (Mo) X-ray Source graphite	2957 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.020$
Detector resolution: 16.0874 pixels mm^{-1} ω scans	$\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 2.9^\circ$ $h = -11 \rightarrow 11$
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2006) $T_{\text{min}} = 0.887$, $T_{\text{max}} = 1.000$	$k = -12 \rightarrow 12$ $l = -12 \rightarrow 13$
6770 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.028$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.067$	H-atom parameters constrained
$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.0257P)^2 + 0.4057P]$ where $P = (F_o^2 + 2F_c^2)/3$
3323 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
200 parameters	$\Delta\rho_{\text{max}} = 0.43 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.55 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.61447 (3)	0.68561 (3)	0.77089 (2)	0.04648 (9)
Cl2	0.29725 (14)	0.32782 (12)	0.59427 (9)	0.0627 (3)
Cl3	0.29306 (15)	0.99624 (11)	1.34350 (10)	0.0603 (3)

S2	0.11153 (10)	0.80692 (9)	1.76536 (8)	0.03759 (19)
O1	0.4697 (3)	0.8156 (3)	1.2278 (2)	0.0620 (8)
O2	0.0367 (3)	0.6656 (3)	1.7893 (2)	0.0536 (7)
O3	0.0121 (4)	0.8840 (4)	1.7587 (3)	0.0676 (8)
N1	0.2439 (3)	0.5816 (3)	1.1495 (2)	0.0344 (6)
H1	0.1576	0.5214	1.1721	0.041*
C1	0.4083 (4)	0.5502 (4)	0.8367 (3)	0.0328 (7)
C2	0.2898 (4)	0.4068 (4)	0.7585 (3)	0.0390 (8)
C3	0.1603 (4)	0.3219 (4)	0.8106 (3)	0.0478 (9)
H3	0.0821	0.2240	0.7588	0.057*
C4	0.1467 (4)	0.3817 (4)	0.9391 (3)	0.0412 (8)
H4	0.0578	0.3246	0.9729	0.049*
C5	0.2643 (4)	0.5261 (3)	1.0182 (3)	0.0303 (6)
C6	0.3978 (4)	0.6101 (3)	0.9675 (3)	0.0326 (7)
H6	0.4798	0.7061	1.0210	0.039*
C7	0.3431 (4)	0.7173 (4)	1.2433 (3)	0.0370 (7)
C8	0.2843 (4)	0.7407 (3)	1.3728 (3)	0.0315 (7)
C9	0.2612 (4)	0.8646 (3)	1.4266 (3)	0.0346 (7)
C10	0.2081 (4)	0.8860 (3)	1.5459 (3)	0.0346 (7)
H10	0.1917	0.9691	1.5806	0.042*
C11	0.1800 (3)	0.7808 (3)	1.6127 (3)	0.0307 (6)
C12	0.2046 (4)	0.6571 (3)	1.5633 (3)	0.0332 (7)
H12	0.1865	0.5880	1.6099	0.040*
C13	0.2566 (4)	0.6382 (4)	1.4433 (3)	0.0336 (7)
H13	0.2734	0.5553	1.4090	0.040*
C14	0.3027 (5)	0.9225 (5)	1.8925 (3)	0.0601 (11)
H14A	0.2762	0.9474	1.9789	0.090*
H14B	0.3667	1.0126	1.8717	0.090*
H14C	0.3697	0.8705	1.8963	0.090*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
II	0.05672 (15)	0.05233 (16)	0.04928 (15)	0.02995 (12)	0.03477 (12)	0.02942 (12)
C11	0.0823 (7)	0.0685 (7)	0.0327 (4)	0.0349 (6)	0.0224 (5)	0.0051 (4)
C12	0.0951 (7)	0.0446 (5)	0.0564 (5)	0.0341 (5)	0.0333 (5)	0.0322 (4)
S2	0.0338 (4)	0.0384 (4)	0.0326 (4)	0.0090 (3)	0.0162 (3)	0.0088 (3)
O1	0.0644 (16)	0.0378 (14)	0.0517 (15)	-0.0044 (13)	0.0346 (13)	0.0034 (12)
O2	0.0539 (14)	0.0436 (15)	0.0454 (13)	0.0031 (12)	0.0266 (12)	0.0140 (11)
O3	0.0703 (18)	0.094 (2)	0.0684 (18)	0.0555 (18)	0.0411 (15)	0.0315 (16)
N1	0.0369 (13)	0.0297 (14)	0.0319 (13)	0.0082 (11)	0.0193 (11)	0.0108 (11)
C1	0.0395 (16)	0.0390 (18)	0.0308 (15)	0.0225 (14)	0.0159 (13)	0.0181 (14)
C2	0.0490 (18)	0.048 (2)	0.0240 (15)	0.0270 (17)	0.0120 (14)	0.0086 (14)
C3	0.0491 (19)	0.040 (2)	0.0351 (18)	0.0103 (17)	0.0068 (15)	0.0020 (15)
C4	0.0386 (16)	0.0398 (19)	0.0350 (17)	0.0075 (15)	0.0120 (14)	0.0134 (14)
C5	0.0358 (15)	0.0330 (16)	0.0258 (14)	0.0157 (13)	0.0129 (12)	0.0134 (12)
C6	0.0356 (15)	0.0292 (16)	0.0325 (15)	0.0128 (13)	0.0116 (13)	0.0117 (13)
C7	0.0403 (17)	0.0300 (17)	0.0345 (16)	0.0089 (14)	0.0156 (14)	0.0107 (14)

supplementary materials

C8	0.0283 (14)	0.0296 (16)	0.0300 (15)	0.0065 (13)	0.0098 (12)	0.0104 (13)
C9	0.0421 (16)	0.0282 (16)	0.0324 (15)	0.0126 (14)	0.0123 (14)	0.0135 (13)
C10	0.0367 (16)	0.0297 (16)	0.0345 (16)	0.0132 (14)	0.0105 (13)	0.0083 (13)
C11	0.0267 (13)	0.0341 (17)	0.0260 (14)	0.0092 (13)	0.0078 (12)	0.0092 (12)
C12	0.0357 (15)	0.0314 (17)	0.0337 (15)	0.0135 (13)	0.0122 (13)	0.0147 (13)
C13	0.0390 (16)	0.0320 (17)	0.0332 (16)	0.0178 (14)	0.0142 (13)	0.0112 (13)
C14	0.056 (2)	0.051 (2)	0.0362 (19)	-0.0018 (19)	0.0054 (17)	0.0050 (17)

Geometric parameters (Å, °)

I1—C1	2.090 (3)	C4—C5	1.382 (4)
Cl1—C2	1.735 (3)	C5—C6	1.391 (4)
Cl2—C9	1.731 (3)	C6—H6	0.9300
S2—O2	1.431 (3)	C7—C8	1.504 (4)
S2—O3	1.426 (3)	C8—C9	1.384 (4)
S2—C11	1.769 (3)	C8—C13	1.388 (4)
S2—C14	1.757 (4)	C9—C10	1.381 (4)
O1—C7	1.218 (4)	C10—H10	0.9300
N1—H1	0.8600	C10—C11	1.383 (4)
N1—C5	1.416 (4)	C11—C12	1.380 (4)
N1—C7	1.348 (4)	C12—H12	0.9300
C1—C2	1.375 (5)	C12—C13	1.381 (4)
C1—C6	1.388 (4)	C13—H13	0.9300
C2—C3	1.381 (5)	C14—H14A	0.9600
C3—H3	0.9300	C14—H14B	0.9600
C3—C4	1.378 (4)	C14—H14C	0.9600
C4—H4	0.9300		
O2—S2—C11	107.98 (15)	O1—C7—N1	124.5 (3)
O2—S2—C14	106.74 (19)	O1—C7—C8	121.2 (3)
O3—S2—O2	118.76 (18)	N1—C7—C8	114.3 (2)
O3—S2—C11	108.53 (16)	C9—C8—C7	121.8 (3)
O3—S2—C14	109.7 (2)	C9—C8—C13	118.5 (3)
C14—S2—C11	104.21 (16)	C13—C8—C7	119.7 (3)
C5—N1—H1	116.0	C8—C9—Cl2	120.7 (2)
C7—N1—H1	116.0	C10—C9—Cl2	117.7 (3)
C7—N1—C5	128.0 (2)	C10—C9—C8	121.6 (3)
C2—C1—H1	123.1 (2)	C9—C10—H10	120.8
C2—C1—C6	120.3 (3)	C9—C10—C11	118.4 (3)
C6—C1—H1	116.6 (2)	C11—C10—H10	120.8
C1—C2—Cl1	121.8 (2)	C10—C11—S2	118.7 (2)
C1—C2—C3	119.8 (3)	C12—C11—S2	119.6 (2)
C3—C2—Cl1	118.3 (3)	C12—C11—C10	121.7 (3)
C2—C3—H3	119.9	C11—C12—H12	120.7
C4—C3—C2	120.3 (3)	C11—C12—C13	118.6 (3)
C4—C3—H3	119.9	C13—C12—H12	120.7
C3—C4—H4	119.8	C8—C13—H13	119.4
C3—C4—C5	120.4 (3)	C12—C13—C8	121.3 (3)
C5—C4—H4	119.8	C12—C13—H13	119.4
C4—C5—N1	117.9 (3)	S2—C14—H14A	109.5

C4—C5—C6	119.4 (3)	S2—C14—H14B	109.5
C6—C5—N1	122.7 (3)	S2—C14—H14C	109.5
C1—C6—C5	119.8 (3)	H14A—C14—H14B	109.5
C1—C6—H6	120.1	H14A—C14—H14C	109.5
C5—C6—H6	120.1	H14B—C14—H14C	109.5
I1—C1—C2—C11	-2.5 (4)	C4—C5—C6—C1	2.2 (5)
I1—C1—C2—C3	177.7 (3)	C5—N1—C7—O1	0.1 (6)
I1—C1—C6—C5	-179.8 (2)	C5—N1—C7—C8	179.6 (3)
C11—C2—C3—C4	-178.1 (3)	C6—C1—C2—C11	179.6 (2)
C12—C9—C10—C11	179.2 (2)	C6—C1—C2—C3	-0.2 (5)
S2—C11—C12—C13	179.4 (2)	C7—N1—C5—C4	179.5 (3)
O1—C7—C8—C9	51.1 (5)	C7—N1—C5—C6	0.2 (5)
O1—C7—C8—C13	-127.0 (4)	C7—C8—C9—C12	2.0 (4)
O2—S2—C11—C10	160.6 (2)	C7—C8—C9—C10	-179.6 (3)
O2—S2—C11—C12	-19.6 (3)	C7—C8—C13—C12	179.2 (3)
O3—S2—C11—C10	30.7 (3)	C8—C9—C10—C11	0.8 (5)
O3—S2—C11—C12	-149.6 (3)	C9—C8—C13—C12	1.0 (4)
N1—C5—C6—C1	-178.5 (3)	C9—C10—C11—S2	-179.9 (2)
N1—C7—C8—C9	-128.4 (3)	C9—C10—C11—C12	0.4 (4)
N1—C7—C8—C13	53.5 (4)	C10—C11—C12—C13	-0.8 (4)
C1—C2—C3—C4	1.8 (6)	C11—C12—C13—C8	0.1 (4)
C2—C1—C6—C5	-1.7 (5)	C13—C8—C9—C12	-179.8 (2)
C2—C3—C4—C5	-1.3 (6)	C13—C8—C9—C10	-1.5 (4)
C3—C4—C5—N1	-180.0 (3)	C14—S2—C11—C10	-86.2 (3)
C3—C4—C5—C6	-0.7 (5)	C14—S2—C11—C12	93.6 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N1-H1\cdots O2^i$	0.86	2.15	2.991 (3)	167

Symmetry codes: (i) $-x, -y+1, -z+3$.

Fig. 1

