

(2,3-Di-2-pyridylpyrazine- κ^2N^2,N^3)bis-(thiocyanato- κS)palladium(II)

Kwang Ha

School of Applied Chemical Engineering, The Research Institute of Catalysis, Chonnam National University, Gwangju 500-757, Republic of Korea
Correspondence e-mail: hakwang@chonnam.ac.kr


Received 9 January 2012; accepted 9 January 2012

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(C-C) = 0.008$ Å; R factor = 0.046; wR factor = 0.100; data-to-parameter ratio = 14.4.

The Pd^{II} ion in the title complex, [Pd(NCS)₂(C₁₄H₁₀N₄)], is four-coordinated in a distorted square-planar environment by the two pyridine N atoms of the chelating 2,3-di-2-pyridylpyrazine (dpp) ligand and two S atoms from two thiocyanate anions. The pyridine rings are considerably inclined to the least-squares plane of the PdS₂N₂ unit [maximum deviation = 0.027 (1) Å], making dihedral angles of 70.3 (2) and 69.2 (1)°. The pyrazine ring is almost perpendicular to the PdS₂N₂ plane, with a dihedral angle of 86.3 (1)°. The thiocyanate ligands are located on opposite sides of the PdS₂N₂ unit plane and are almost linear [N—C—S angles = 177.8 (6) and 178.9 (6)°]. The complex molecules are stacked in columns along the b axis and are connected by intermolecular C—H...N hydrogen bonds, forming chains along the a axis.

Related literature

For related crystal structures of [PdX₂(dpp)] ($X = Cl, Br$ or I), see: Ha (2011a,b,c). For related Pt and Pd complexes, see: Granifo *et al.* (2000); Cai *et al.* (2009).



Experimental

Crystal data

[Pd(NCS)₂(C₁₄H₁₀N₄)]
 $M_r = 456.82$
 Monoclinic, $P2_1/c$
 $a = 15.8236$ (11) Å
 $b = 13.5901$ (9) Å
 $c = 7.9189$ (6) Å
 $\beta = 102.960$ (1)°
 $V = 1659.5$ (2) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 1.38$ mm⁻¹
 $T = 200$ K
 0.27 × 0.25 × 0.13 mm

Data collection

Bruker SMART 1000 CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{min} = 0.835$, $T_{max} = 1.000$
 10174 measured reflections
 3247 independent reflections
 2248 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.071$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.100$
 $S = 1.00$
 3247 reflections
 226 parameters
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.89$ e Å⁻³
 $\Delta\rho_{min} = -0.66$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Pd1—N3	2.059 (4)	Pd1—S1	2.3090 (17)
Pd1—N4	2.039 (5)	Pd1—S2	2.3045 (15)
N4—Pd1—N3	86.73 (16)	S2—Pd1—S1	82.78 (6)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C12—H12...N2 ⁱ	0.95	2.54	3.306 (7)	138
C14—H14...N5 ⁱⁱ	0.95	2.49	3.277 (8)	140

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $-x, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Priority Research Centers Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2010-0029626).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5782).

References

- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cai, X., Donzello, M. P., Viola, E., Rizzoli, C., Ercolani, C. & Kadish, K. M. (2009). *Inorg. Chem.* **48**, 7086–7098.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Granifo, J., Vargas, M. E., Garland, M. T. & Baggio, R. (2000). *Inorg. Chim. Acta*, **305**, 143–150.
- Ha, K. (2011a). *Acta Cryst.* **E67**, m1615.
- Ha, K. (2011b). *Acta Cryst.* **E67**, m1896.
- Ha, K. (2011c). *Acta Cryst.* **E67**, m1626.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2012). E68, m144 [doi:10.1107/S1600536812000980]

(2,3-Di-2-pyridylpyrazine- κ^2N^2,N^3)bis(thiocyanato- κS)palladium(II)

K. Ha

Comment

The title complex, [Pd(SCN)₂(dpp)] (dpp = 2,3-di-2-pyridylpyrazine, C₁₄H₁₀N₄), is isomorphous with the previously reported analogous halogen complexes [PdX₂(dpp)] (X = Cl, Br or I) (Ha, 2011*a,b,c*). The Pd^{II} ion is four-coordinated in a distorted square-planar environment by the two pyridine N atoms of the chelating dpp ligand and two S atoms from two thiocyanate anions (Fig. 1). The coordination mode of the dpp ligand is similar to that found in the mononuclear Pt(II) and Pd(II) complexes [PtCl₂(dpq)] (dpq = 2,3-di-2-pyridylquinoxaline) (Granifo *et al.*, 2000) and [MCl₂(dcdpp)] (M = Pt, Pd; dcdpp = 2,3-dicyano-5,6-di-2-pyridylpyrazine) (Cai *et al.*, 2009).

In the crystal, the pyridine rings are considerably inclined to the least-squares plane of the PdS₂N₂ unit [maximum deviation = 0.027 (1) Å], making dihedral angles of 70.3 (2)° and 69.2 (1)°. The nearly planar pyrazine ring [maximum deviation = 0.013 (4) Å] is almost perpendicular to the unit plane with a dihedral angle of 86.3 (1)°. The dihedral angle between the two pyridine rings is 76.8 (2)°. The thiocyanato ligands are located on opposite sides of the PdS₂N₂ unit plane and are almost linear with the bond angles of <S1—C15—N5 = 177.8 (6)° and <S2—C16—N6 = 178.9 (6)°. The Pd—N and Pd—S bond lengths are nearly equivalent, respectively (Table 1). The complex molecules are stacked in columns along the *b* axis and are connected by intermolecular C—H...N hydrogen bonds, forming chains along the *a* axis (Fig.2 and Table 2). In the columns, numerous inter- and intramolecular π - π interactions between the six-membered rings are present, the shortest ring centroid-centroid distance being 4.061 (3) Å.

Experimental

To a suspension of Na₂PdCl₄ (0.1475 g, 0.501 mmol) and KSCN (0.5456 g, 5.614 mmol) in MeOH (30 ml) was added 2,3-di-2-pyridylpyrazine (0.1176 g, 0.502 mmol), and stirred for 24 h at room temperature. After removal of the formed brown precipitate by filtration, the solvent of the filtrate was evaporated. The residue was washed with H₂O, and dried at 50 °C, to give a yellow powder (0.2119 g). Crystals suitable for X-ray analysis were obtained by slow evaporation from a CH₃CN solution.

Refinement

H atoms were positioned geometrically and allowed to ride on their respective parent atoms [C—H = 0.95 Å and $U_{iso}(H) = 1.2U_{eq}(C)$]. The highest peak (0.89 e Å⁻³) and the deepest hole (-0.66 e Å⁻³) in the difference Fourier map are located 1.05 Å and 0.99 Å from the atoms S1 and Pd1, respectively.

Figures




Fig. 1. A view of the molecular structure of the title complex, with displacement ellipsoids drawn at the 40% probability level and the atom numbering.




Fig. 2. A partial view of the crystal packing of the title complex. Hydrogen-bond interactions are drawn with dashed lines.

(2,3-Di-2-pyridylpyrazine- κ^2N^2,N^3)bis(thiocyanato- κS)palladium(II)

Crystal data

[Pd(NCS)₂(C₁₄H₁₀N₄)]

$M_r = 456.82$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 15.8236$ (11) Å

$b = 13.5901$ (9) Å

$c = 7.9189$ (6) Å

$\beta = 102.960$ (1)°

$V = 1659.5$ (2) Å³

$Z = 4$

$F(000) = 904$

$D_x = 1.828$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2951 reflections

$\theta = 2.6$ – 25.6 °

$\mu = 1.38$ mm⁻¹

$T = 200$ K

Plate, yellow

$0.27 \times 0.25 \times 0.13$ mm

Data collection

Bruker SMART 1000 CCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2000)

$T_{\min} = 0.835$, $T_{\max} = 1.000$

10174 measured reflections

3247 independent reflections

2248 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.071$

$\theta_{\text{max}} = 26.0$ °, $\theta_{\text{min}} = 2.0$ °

$h = -19 \rightarrow 19$

$k = -16 \rightarrow 14$

$l = -9 \rightarrow 9$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct
methods

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.046$$

$$wR(F^2) = 0.100$$

$$S = 1.00$$

3247 reflections

226 parameters

0 restraints

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0351P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.89 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.66 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pd1	0.19166 (3)	0.47138 (3)	0.53401 (6)	0.02445 (15)
S1	0.11316 (10)	0.43246 (12)	0.2597 (2)	0.0379 (4)
S2	0.23273 (12)	0.30866 (11)	0.5347 (2)	0.0455 (5)
N1	0.3412 (3)	0.7055 (3)	0.3986 (6)	0.0299 (12)
N2	0.4466 (3)	0.5833 (3)	0.6359 (6)	0.0298 (11)
N3	0.1593 (3)	0.6182 (3)	0.5309 (5)	0.0229 (10)
N4	0.2666 (3)	0.4960 (3)	0.7755 (6)	0.0236 (11)
N5	-0.0358 (3)	0.5536 (4)	0.2015 (7)	0.0434 (14)
N6	0.3992 (4)	0.3047 (4)	0.7615 (8)	0.0519 (16)
C1	0.3097 (3)	0.6576 (4)	0.5210 (7)	0.0237 (13)
C2	0.3637 (3)	0.5968 (4)	0.6413 (7)	0.0249 (13)
C3	0.4761 (4)	0.6298 (4)	0.5124 (8)	0.0330 (15)
H3	0.5350	0.6215	0.5063	0.040*
C4	0.4231 (4)	0.6892 (4)	0.3940 (8)	0.0317 (14)
H4	0.4459	0.7197	0.3058	0.038*
C5	0.2195 (3)	0.6873 (4)	0.5241 (7)	0.0230 (13)
C6	0.1985 (4)	0.7861 (4)	0.5149 (8)	0.0323 (14)
H6	0.2410	0.8338	0.5051	0.039*
C7	0.1161 (4)	0.8153 (4)	0.5200 (8)	0.0328 (15)
H7	0.1013	0.8832	0.5138	0.039*
C8	0.0551 (4)	0.7456 (4)	0.5339 (8)	0.0331 (14)
H8	-0.0020	0.7645	0.5399	0.040*
C9	0.0785 (4)	0.6479 (4)	0.5391 (7)	0.0256 (13)

supplementary materials

H9	0.0365	0.5995	0.5487	0.031*
C10	0.3414 (3)	0.5479 (3)	0.7954 (7)	0.0218 (12)
C11	0.3978 (4)	0.5543 (4)	0.9550 (7)	0.0298 (14)
H11	0.4494	0.5921	0.9681	0.036*
C12	0.3797 (4)	0.5061 (4)	1.0957 (8)	0.0308 (14)
H12	0.4190	0.5086	1.2056	0.037*
C13	0.3029 (4)	0.4541 (4)	1.0736 (8)	0.0323 (14)
H13	0.2884	0.4206	1.1684	0.039*
C14	0.2480 (4)	0.4517 (4)	0.9131 (8)	0.0292 (14)
H14	0.1947	0.4171	0.8992	0.035*
C15	0.0257 (4)	0.5056 (4)	0.2271 (8)	0.0326 (14)
C16	0.3312 (5)	0.3072 (4)	0.6693 (9)	0.0371 (16)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pd1	0.0243 (3)	0.0191 (2)	0.0282 (3)	0.00066 (18)	0.00218 (18)	-0.00024 (19)
S1	0.0347 (10)	0.0434 (10)	0.0325 (10)	-0.0005 (7)	0.0006 (8)	-0.0074 (7)
S2	0.0538 (12)	0.0205 (8)	0.0555 (12)	0.0066 (7)	-0.0018 (9)	-0.0058 (7)
N1	0.029 (3)	0.028 (3)	0.034 (3)	0.002 (2)	0.009 (2)	0.004 (2)
N2	0.026 (3)	0.028 (3)	0.036 (3)	0.003 (2)	0.008 (2)	0.002 (2)
N3	0.027 (3)	0.022 (2)	0.020 (3)	-0.0008 (19)	0.005 (2)	0.0022 (19)
N4	0.022 (3)	0.022 (2)	0.027 (3)	0.0048 (18)	0.006 (2)	0.0001 (19)
N5	0.028 (3)	0.052 (4)	0.046 (4)	0.001 (3)	-0.002 (3)	0.004 (3)
N6	0.039 (4)	0.052 (4)	0.065 (5)	0.016 (3)	0.013 (3)	0.009 (3)
C1	0.021 (3)	0.019 (3)	0.029 (3)	-0.004 (2)	0.002 (3)	-0.007 (2)
C2	0.025 (3)	0.020 (3)	0.027 (3)	-0.002 (2)	0.000 (3)	-0.004 (2)
C3	0.029 (4)	0.032 (3)	0.042 (4)	0.003 (3)	0.014 (3)	-0.004 (3)
C4	0.033 (4)	0.024 (3)	0.040 (4)	-0.003 (3)	0.011 (3)	-0.002 (3)
C5	0.024 (3)	0.025 (3)	0.017 (3)	0.001 (2)	-0.001 (3)	0.001 (2)
C6	0.033 (4)	0.028 (3)	0.032 (4)	0.000 (3)	0.000 (3)	0.000 (3)
C7	0.043 (4)	0.019 (3)	0.037 (4)	0.009 (3)	0.010 (3)	0.002 (3)
C8	0.026 (3)	0.034 (3)	0.037 (4)	0.009 (3)	0.003 (3)	-0.001 (3)
C9	0.029 (3)	0.028 (3)	0.019 (3)	-0.004 (2)	0.004 (3)	0.001 (2)
C10	0.016 (3)	0.018 (3)	0.029 (3)	0.005 (2)	0.001 (2)	0.001 (2)
C11	0.027 (3)	0.027 (3)	0.034 (4)	0.006 (2)	0.004 (3)	-0.002 (3)
C12	0.029 (4)	0.026 (3)	0.032 (4)	0.011 (3)	-0.004 (3)	-0.001 (3)
C13	0.043 (4)	0.025 (3)	0.029 (4)	0.014 (3)	0.008 (3)	0.008 (3)
C14	0.023 (3)	0.029 (3)	0.037 (4)	0.003 (2)	0.010 (3)	0.002 (3)
C15	0.034 (4)	0.041 (4)	0.021 (3)	-0.014 (3)	0.002 (3)	0.003 (3)
C16	0.047 (4)	0.021 (3)	0.049 (5)	0.006 (3)	0.022 (4)	0.006 (3)

Geometric parameters (\AA , $^\circ$)

Pd1—N3	2.059 (4)	C3—C4	1.372 (8)
Pd1—N4	2.039 (5)	C3—H3	0.9500
Pd1—S1	2.3090 (17)	C4—H4	0.9500
Pd1—S2	2.3045 (15)	C5—C6	1.382 (7)
S1—C15	1.677 (7)	C6—C7	1.371 (7)

S2—C16	1.679 (7)	C6—H6	0.9500
N1—C4	1.324 (7)	C7—C8	1.374 (8)
N1—C1	1.352 (7)	C7—H7	0.9500
N2—C3	1.333 (7)	C8—C9	1.377 (7)
N2—C2	1.336 (6)	C8—H8	0.9500
N3—C5	1.347 (6)	C9—H9	0.9500
N3—C9	1.356 (6)	C10—C11	1.378 (7)
N4—C14	1.334 (7)	C11—C12	1.377 (8)
N4—C10	1.356 (6)	C11—H11	0.9500
N5—C15	1.151 (7)	C12—C13	1.383 (8)
N6—C16	1.157 (8)	C12—H12	0.9500
C1—C2	1.398 (7)	C13—C14	1.369 (8)
C1—C5	1.488 (7)	C13—H13	0.9500
C2—C10	1.499 (7)	C14—H14	0.9500
N4—Pd1—N3	86.73 (16)	C6—C5—C1	118.9 (5)
N4—Pd1—S2	92.92 (12)	C7—C6—C5	120.0 (5)
N3—Pd1—S2	177.82 (13)	C7—C6—H6	120.0
N4—Pd1—S1	175.50 (12)	C5—C6—H6	120.0
N3—Pd1—S1	97.51 (13)	C6—C7—C8	119.4 (5)
S2—Pd1—S1	82.78 (6)	C6—C7—H7	120.3
C15—S1—Pd1	105.1 (2)	C8—C7—H7	120.3
C16—S2—Pd1	103.5 (2)	C7—C8—C9	118.6 (5)
C4—N1—C1	117.4 (5)	C7—C8—H8	120.7
C3—N2—C2	117.8 (5)	C9—C8—H8	120.7
C5—N3—C9	118.5 (4)	N3—C9—C8	122.4 (5)
C5—N3—Pd1	120.0 (4)	N3—C9—H9	118.8
C9—N3—Pd1	121.5 (3)	C8—C9—H9	118.8
C14—N4—C10	119.1 (5)	N4—C10—C11	120.6 (5)
C14—N4—Pd1	120.2 (4)	N4—C10—C2	119.7 (5)
C10—N4—Pd1	120.2 (4)	C11—C10—C2	119.8 (5)
N1—C1—C2	120.4 (5)	C12—C11—C10	120.2 (5)
N1—C1—C5	113.0 (5)	C12—C11—H11	119.9
C2—C1—C5	126.2 (5)	C10—C11—H11	119.9
N2—C2—C1	120.9 (5)	C11—C12—C13	118.5 (6)
N2—C2—C10	112.2 (5)	C11—C12—H12	120.8
C1—C2—C10	126.7 (5)	C13—C12—H12	120.8
N2—C3—C4	121.3 (5)	C14—C13—C12	119.1 (5)
N2—C3—H3	119.4	C14—C13—H13	120.4
C4—C3—H3	119.4	C12—C13—H13	120.4
N1—C4—C3	122.1 (5)	N4—C14—C13	122.4 (5)
N1—C4—H4	118.9	N4—C14—H14	118.8
C3—C4—H4	118.9	C13—C14—H14	118.8
N3—C5—C6	121.0 (5)	N5—C15—S1	177.8 (6)
N3—C5—C1	120.1 (5)	N6—C16—S2	178.9 (6)
N3—Pd1—S1—C15	-32.7 (2)	N1—C1—C5—N3	133.2 (5)
S2—Pd1—S1—C15	149.5 (2)	C2—C1—C5—N3	-54.5 (8)
N4—Pd1—S2—C16	-26.8 (3)	N1—C1—C5—C6	-44.9 (7)
S1—Pd1—S2—C16	151.9 (2)	C2—C1—C5—C6	127.3 (6)

supplementary materials

N4—Pd1—N3—C5	68.5 (4)	N3—C5—C6—C7	2.8 (8)
S1—Pd1—N3—C5	-110.0 (4)	C1—C5—C6—C7	-179.1 (5)
N4—Pd1—N3—C9	-110.1 (4)	C5—C6—C7—C8	-0.1 (9)
S1—Pd1—N3—C9	71.4 (4)	C6—C7—C8—C9	-1.2 (9)
N3—Pd1—N4—C14	115.9 (4)	C5—N3—C9—C8	2.6 (8)
S2—Pd1—N4—C14	-66.3 (4)	Pd1—N3—C9—C8	-178.8 (4)
N3—Pd1—N4—C10	-72.1 (4)	C7—C8—C9—N3	0.0 (8)
S2—Pd1—N4—C10	105.8 (4)	C14—N4—C10—C11	0.4 (7)
C4—N1—C1—C2	2.5 (8)	Pd1—N4—C10—C11	-171.7 (4)
C4—N1—C1—C5	175.3 (5)	C14—N4—C10—C2	179.1 (4)
C3—N2—C2—C1	0.0 (8)	Pd1—N4—C10—C2	7.0 (6)
C3—N2—C2—C10	-174.9 (5)	N2—C2—C10—N4	-135.8 (5)
N1—C1—C2—N2	-1.2 (8)	C1—C2—C10—N4	49.7 (7)
C5—C1—C2—N2	-173.0 (5)	N2—C2—C10—C11	42.9 (6)
N1—C1—C2—C10	172.9 (5)	C1—C2—C10—C11	-131.6 (6)
C5—C1—C2—C10	1.2 (9)	N4—C10—C11—C12	1.5 (7)
C2—N2—C3—C4	-0.3 (8)	C2—C10—C11—C12	-177.2 (5)
C1—N1—C4—C3	-2.8 (8)	C10—C11—C12—C13	-1.9 (8)
N2—C3—C4—N1	1.7 (9)	C11—C12—C13—C14	0.5 (8)
C9—N3—C5—C6	-4.0 (8)	C10—N4—C14—C13	-1.9 (7)
Pd1—N3—C5—C6	177.4 (4)	Pd1—N4—C14—C13	170.2 (4)
C9—N3—C5—C1	177.9 (5)	C12—C13—C14—N4	1.4 (8)
Pd1—N3—C5—C1	-0.7 (7)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C12—H12 \cdots N2 ⁱ	0.95	2.54	3.306 (7)	138.
C14—H14 \cdots N5 ⁱⁱ	0.95	2.49	3.277 (8)	140.

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $-x, -y+1, -z+1$.

Fig. 1




Fig. 2

