

{4,4',6,6'-Tetrachloro-2,2'-[2,2-dimethylpropane-1,3-diylbis(nitrilomethanylylidene)]copper(II)}

Hadi Kargar,^{a*} Reza Kia,^{b,c} Saeideh Abbasian^a and Muhammad Nawaz Tahir^{d*}

^aDepartment of Chemistry, Payame Noor University, PO Box 19395-3697 Tehran, I. R. of IRAN, ^bX-ray Crystallography Lab., Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran, ^cDepartment of Chemistry, Science and Research Branch, Islamic Azad University, Tehran, Iran, and ^dDepartment of Physics, University of Sargodha, Punjab, Pakistan
Correspondence e-mail: hkargar@pnu.ac.ir, dmntahir_uos@yahoo.com


Received 7 January 2012; accepted 16 January 2012

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.047; wR factor = 0.105; data-to-parameter ratio = 19.6.

In the title Schiff base complex, $[\text{Cu}(\text{C}_{19}\text{H}_{16}\text{Cl}_4\text{N}_2\text{O}_2)]$, the geometry around the Cu^{II} atom is distorted square-planar defined by the N_2O_2 donor atoms of the coordinated ligand. The dihedral angle between the substituted benzene rings is 29.95 (16)°. In the crystal, molecules are linked along the b axis, forming individual dimers through $\text{C}-\text{H}\cdots\text{O}$ interactions. The crystal structure is further stabilized by intermolecular $\pi-\pi$ interactions [centroid-centroid distance = 3.6131 (17) Å].

Related literature

For standard values of bond lengths, see: Allen *et al.* (1987). For applications of Schiff bases in coordination chemistry, see, for example: Granovski *et al.* (1993); Blower (1998). For related structures see, for example: Ghaemi *et al.* (2011); Kargar *et al.* (2011, 2012).



Experimental

Crystal data

$[\text{Cu}(\text{C}_{19}\text{H}_{16}\text{Cl}_4\text{N}_2\text{O}_2)]$
 $M_r = 509.68$
 Monoclinic, $P2_1/n$
 $a = 12.4002$ (10) Å
 $b = 8.4570$ (7) Å
 $c = 20.0316$ (19) Å
 $\beta = 97.278$ (4)°
 $V = 2083.8$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.58$ mm⁻¹
 $T = 291$ K
 $0.25 \times 0.18 \times 0.09$ mm

Data collection

Bruker SMART APEXII CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\text{min}} = 0.694$, $T_{\text{max}} = 0.871$
 18291 measured reflections
 4988 independent reflections
 2882 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.060$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.105$
 $S = 1.00$
 4988 reflections
 255 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.38$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.36$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C8}-\text{H8A}\cdots\text{O1}^i$	0.97	2.56	3.331 (4)	136

Symmetry code: (i) $-x + 1, -y + 2, -z$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HK and SA thank PNU for financial support. MNT thanks GC University of Sargodha, Pakistan for the research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HP2025).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Blower, P. J. (1998). *Transition Met. Chem.*, **23**, 109–112.
 Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Ghaemi, A., Rayati, S., Elahi, E., Ng, S. W. & Tiekink, E. R. T. (2011). *Acta Cryst. E* **67**, m1445–m1446.
 Granovski, A. D., Nivorozhkin, A. L. & Minkin, V. I. (1993). *Coord. Chem. Rev.* **126**, 1–69.
 Kargar, H., Kia, R., Pahlavani, E. & Tahir, M. N. (2011). *Acta Cryst. E* **67**, m941.
 Kargar, H., Kia, R., Sharafi, Z. & Tahir, M. N. (2012). *Acta Cryst. E* **68**, m82.
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2012). E68, m182 [doi:10.1107/S160053681200195X]

{4,4',6,6'-Tetrachloro-2,2'-[2,2-dimethylpropane-1,3-diylbis(nitrilomethanylylidene)]}copper(II)

H. Kargar, R. Kia, S. Abbasian and M. N. Tahir

Comment

Schiff base complexes are one of the most important stereochemical models in transition metal coordination chemistry, with the ease of preparation and structural variations (Granovski *et al.*, 1993; Blower *et al.*, (1998). In continuation of our work on the crystal structure of Schiff base metal complexes (Kargar *et al.*, 2012; Kargar *et al.*, 2011; Ghaemi, *et al.*, (2011), we determined the X-ray structure of the title compound.

The asymmetric unit of the title compound, Fig. 1, comprises a Schiff base complex. The bond lengths (Allen *et al.*, 1987) and angles are within the normal ranges and are comparable to the related structure (Kargar *et al.*, 2012; Kargar *et al.*, 2011; Ghaemi, *et al.*, (2011).

The geometry around Cu^{II} is a distorted square-planar which is supported by the N₂O₂ donor atoms of the coordinated Schiff base ligand. The dihedral angle between the substituted benzene rings is 29.95 (16)°. In the crystal structure the molecules are linked together along the *b*-axis, forming individual dimers through the intermolecular C—H...O interactions (Table 1, Fig. 2). The crystal structure is further stabilized by the intermolecular π - π interaction [Cg1...Cg1ⁱⁱ = 3.6131 (17)Å; (ii) 1 - *X*, 1 - *Y*, -*Z*; Cg1 is the centroid of Cu(1)/O(2)/C(19)/C(14)/C(13)/N(2) ring].

Experimental

The title compound was synthesized by adding 3,5-dichloro-salicylaldehyde-2,2-dimethyl-1, 3-propanediamine (2 mmol) to a solution of CuCl₂. 4H₂O (2.1 mmol) in ethanol (30 ml). The mixture was refluxed with stirring for half an hour. The resultant solution was filtered. Dark-green single crystals of the title compound suitable for X-ray structure determination were recrystallized from ethanol by slow evaporation of the solvents at room temperature over several days.

Refinement

The H-atoms were included in calculated positions and treated as riding atoms: C—H = 0.93, 0.96 and 0.97 Å for CH, CH₃ and CH₂ H-atoms, respectively, with U_{iso}(H) = k x U_{eq}(C), where k = 1.5 for CH₃ H-atoms, and k = 1.2 for all other H-atoms.

Figures




Fig. 1. The ORTEP plot of the title compound, showing 40% probability displacement ellipsoids and the atomic numbering.




Fig. 2. A part of the packing diagram of the title compound showing individual dimer formation through the intermolecular C—H...O interactions (dashed lines). Only the H atoms involved in the interactions are shown.

{4,4',6,6'-Tetrachloro-2,2'-[2,2-dimethylpropane-1,3- diylbis(nitrilomethanylidene)]}copper(II)

Crystal data

[Cu(C₁₉H₁₆Cl₄N₂O₂)]

$M_r = 509.68$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 12.4002$ (10) Å

$b = 8.4570$ (7) Å

$c = 20.0316$ (19) Å

$\beta = 97.278$ (4)°

$V = 2083.8$ (3) Å³

$Z = 4$

$F(000) = 1028$

$D_x = 1.625$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2540 reflections

$\theta = 2.5$ – 27.4 °

$\mu = 1.58$ mm⁻¹

$T = 291$ K

Block, dark-green

$0.25 \times 0.18 \times 0.09$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\min} = 0.694$, $T_{\max} = 0.871$

18291 measured reflections

4988 independent reflections

2882 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.060$

$\theta_{\max} = 27.9$ °, $\theta_{\min} = 2.1$ °

$h = -16 \rightarrow 14$

$k = -9 \rightarrow 11$

$l = -26 \rightarrow 26$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.105$

$S = 1.00$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0376P)^2]$

4988 reflections
255 parameters
0 restraints

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5462 (3)	0.8875 (4)	0.14363 (17)	0.0370 (8)
C2	0.6238 (3)	0.9020 (4)	0.20186 (18)	0.0428 (9)
C3	0.6049 (3)	0.9834 (4)	0.25795 (18)	0.0468 (9)
H3	0.6577	0.9874	0.2953	0.056*
C4	0.5058 (3)	1.0605 (4)	0.25883 (18)	0.0486 (9)
C5	0.4286 (3)	1.0532 (4)	0.20422 (18)	0.0430 (9)
H5	0.3629	1.1057	0.2051	0.052*
C6	0.4466 (3)	0.9681 (4)	0.14670 (17)	0.0365 (8)
C7	0.3634 (3)	0.9733 (4)	0.08983 (17)	0.0398 (8)
H7	0.3024	1.0343	0.0944	0.048*
C8	0.2764 (3)	0.9331 (4)	-0.02093 (17)	0.0442 (9)
H8A	0.3077	0.9657	-0.0607	0.053*
H8B	0.2327	1.0201	-0.0078	0.053*
C9	0.2022 (3)	0.7905 (4)	-0.03888 (18)	0.0414 (8)
C10	0.1184 (3)	0.7758 (5)	0.0105 (2)	0.0725 (13)
H10A	0.1548	0.7761	0.0557	0.109*
H10B	0.0788	0.6788	0.0022	0.109*
H10C	0.0689	0.8634	0.0045	0.109*
C11	0.1447 (3)	0.8178 (5)	-0.1105 (2)	0.0722 (13)
H11A	0.0918	0.7361	-0.1217	0.108*
H11B	0.1973	0.8156	-0.1418	0.108*
H11C	0.1091	0.9188	-0.1127	0.108*
C12	0.2678 (2)	0.6363 (4)	-0.03537 (18)	0.0436 (9)
H12A	0.2730	0.5953	0.0101	0.052*
H12B	0.2283	0.5593	-0.0649	0.052*
C13	0.4035 (3)	0.5747 (4)	-0.10521 (18)	0.0411 (9)
H13	0.3483	0.5186	-0.1305	0.049*
C14	0.5102 (3)	0.5676 (4)	-0.12688 (18)	0.0394 (8)

supplementary materials

C15	0.5173 (3)	0.4938 (4)	-0.18871 (19)	0.0490 (10)
H15	0.4548	0.4545	-0.2138	0.059*
C16	0.6153 (3)	0.4793 (4)	-0.21236 (18)	0.0505 (10)
C17	0.7093 (3)	0.5351 (4)	-0.17506 (18)	0.0464 (9)
H17	0.7761	0.5236	-0.1910	0.056*
C18	0.7030 (2)	0.6076 (4)	-0.11442 (17)	0.0374 (8)
C19	0.6035 (2)	0.6294 (4)	-0.08710 (17)	0.0354 (8)
C11	0.74867 (8)	0.80938 (13)	0.20120 (6)	0.0735 (4)
C12	0.48425 (9)	1.16671 (15)	0.33020 (6)	0.0774 (4)
C13	0.62354 (9)	0.38486 (16)	-0.28877 (6)	0.0822 (4)
C14	0.82198 (6)	0.67591 (10)	-0.06837 (5)	0.0476 (2)
Cu1	0.48016 (3)	0.76529 (5)	0.01057 (2)	0.03872 (14)
N1	0.3649 (2)	0.9025 (3)	0.03362 (14)	0.0384 (7)
N2	0.37799 (19)	0.6517 (3)	-0.05406 (14)	0.0374 (7)
O1	0.57082 (17)	0.8082 (3)	0.09215 (12)	0.0448 (6)
O2	0.60372 (17)	0.6982 (3)	-0.02938 (12)	0.0431 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0357 (19)	0.035 (2)	0.040 (2)	-0.0035 (15)	0.0030 (16)	0.0027 (16)
C2	0.0335 (19)	0.038 (2)	0.055 (2)	0.0000 (15)	-0.0015 (17)	0.0029 (18)
C3	0.048 (2)	0.045 (2)	0.045 (2)	-0.0118 (17)	-0.0026 (18)	0.0024 (18)
C4	0.050 (2)	0.054 (2)	0.042 (2)	-0.0112 (19)	0.0095 (19)	-0.0025 (18)
C5	0.040 (2)	0.045 (2)	0.045 (2)	0.0019 (17)	0.0106 (18)	0.0005 (18)
C6	0.0335 (19)	0.036 (2)	0.040 (2)	-0.0006 (15)	0.0041 (16)	0.0032 (16)
C7	0.0345 (19)	0.037 (2)	0.049 (2)	0.0021 (15)	0.0108 (17)	0.0023 (17)
C8	0.046 (2)	0.045 (2)	0.040 (2)	0.0141 (17)	-0.0017 (17)	0.0079 (17)
C9	0.0311 (18)	0.047 (2)	0.046 (2)	0.0060 (15)	0.0005 (16)	0.0039 (17)
C10	0.044 (2)	0.084 (3)	0.094 (4)	0.007 (2)	0.028 (2)	-0.004 (3)
C11	0.070 (3)	0.067 (3)	0.070 (3)	0.010 (2)	-0.027 (2)	-0.001 (2)
C12	0.0301 (18)	0.046 (2)	0.055 (2)	-0.0014 (16)	0.0085 (17)	0.0046 (18)
C13	0.0324 (19)	0.036 (2)	0.053 (2)	-0.0017 (15)	-0.0012 (17)	0.0003 (17)
C14	0.0326 (19)	0.036 (2)	0.049 (2)	0.0016 (15)	0.0036 (17)	-0.0009 (17)
C15	0.040 (2)	0.051 (2)	0.055 (3)	-0.0033 (17)	0.0005 (19)	-0.0141 (19)
C16	0.044 (2)	0.057 (2)	0.052 (3)	-0.0002 (18)	0.0116 (19)	-0.018 (2)
C17	0.039 (2)	0.048 (2)	0.056 (2)	0.0011 (17)	0.0177 (19)	-0.0066 (19)
C18	0.0329 (18)	0.0348 (19)	0.045 (2)	-0.0006 (14)	0.0066 (16)	0.0010 (16)
C19	0.0337 (19)	0.0294 (18)	0.043 (2)	0.0025 (14)	0.0034 (16)	0.0011 (16)
C11	0.0471 (6)	0.0767 (8)	0.0895 (9)	0.0202 (5)	-0.0185 (6)	-0.0163 (6)
C12	0.0730 (8)	0.1044 (9)	0.0569 (7)	-0.0119 (7)	0.0172 (6)	-0.0287 (6)
C13	0.0600 (7)	0.1171 (10)	0.0709 (8)	-0.0016 (6)	0.0130 (6)	-0.0493 (7)
C14	0.0320 (5)	0.0540 (6)	0.0568 (6)	-0.0024 (4)	0.0058 (4)	-0.0025 (5)
Cu1	0.0315 (2)	0.0416 (3)	0.0435 (3)	0.00221 (18)	0.00652 (19)	-0.0001 (2)
N1	0.0312 (15)	0.0406 (17)	0.0418 (18)	0.0036 (12)	-0.0011 (13)	0.0022 (14)
N2	0.0293 (15)	0.0356 (16)	0.0481 (18)	-0.0012 (12)	0.0086 (13)	0.0008 (14)
O1	0.0317 (13)	0.0533 (15)	0.0483 (16)	0.0058 (11)	0.0006 (11)	-0.0046 (12)
O2	0.0321 (13)	0.0509 (15)	0.0469 (15)	-0.0002 (10)	0.0073 (11)	-0.0083 (12)

Geometric parameters (Å, °)

C1—O1	1.299 (4)	C11—H11A	0.9600
C1—C6	1.418 (4)	C11—H11B	0.9600
C1—C2	1.420 (4)	C11—H11C	0.9600
C2—C3	1.362 (5)	C12—N2	1.467 (4)
C2—C11	1.737 (3)	C12—H12A	0.9700
C3—C4	1.394 (5)	C12—H12B	0.9700
C3—H3	0.9300	C13—N2	1.287 (4)
C4—C5	1.360 (5)	C13—C14	1.444 (4)
C4—C12	1.737 (4)	C13—H13	0.9300
C5—C6	1.400 (4)	C14—C15	1.399 (5)
C5—H5	0.9300	C14—C19	1.419 (4)
C6—C7	1.437 (4)	C15—C16	1.365 (4)
C7—N1	1.278 (4)	C15—H15	0.9300
C7—H7	0.9300	C16—C17	1.385 (5)
C8—N1	1.470 (4)	C16—C13	1.741 (3)
C8—C9	1.532 (4)	C17—C18	1.372 (4)
C8—H8A	0.9700	C17—H17	0.9300
C8—H8B	0.9700	C18—C19	1.424 (4)
C9—C10	1.528 (5)	C18—C14	1.737 (3)
C9—C12	1.534 (4)	C19—O2	1.294 (4)
C9—C11	1.537 (5)	Cu1—O1	1.898 (2)
C10—H10A	0.9600	Cu1—O2	1.903 (2)
C10—H10B	0.9600	Cu1—N1	1.942 (3)
C10—H10C	0.9600	Cu1—N2	1.947 (3)
O1—C1—C6	125.2 (3)	H11A—C11—H11C	109.5
O1—C1—C2	119.6 (3)	H11B—C11—H11C	109.5
C6—C1—C2	115.2 (3)	N2—C12—C9	114.7 (3)
C3—C2—C1	123.5 (3)	N2—C12—H12A	108.6
C3—C2—C11	118.7 (3)	C9—C12—H12A	108.6
C1—C2—C11	117.8 (3)	N2—C12—H12B	108.6
C2—C3—C4	119.3 (3)	C9—C12—H12B	108.6
C2—C3—H3	120.3	H12A—C12—H12B	107.6
C4—C3—H3	120.3	N2—C13—C14	126.0 (3)
C5—C4—C3	120.0 (3)	N2—C13—H13	117.0
C5—C4—C12	121.3 (3)	C14—C13—H13	117.0
C3—C4—C12	118.7 (3)	C15—C14—C19	121.5 (3)
C4—C5—C6	121.1 (3)	C15—C14—C13	116.6 (3)
C4—C5—H5	119.5	C19—C14—C13	121.9 (3)
C6—C5—H5	119.5	C16—C15—C14	120.4 (3)
C5—C6—C1	120.8 (3)	C16—C15—H15	119.8
C5—C6—C7	117.6 (3)	C14—C15—H15	119.8
C1—C6—C7	121.5 (3)	C15—C16—C17	120.5 (3)
N1—C7—C6	126.5 (3)	C15—C16—C13	120.0 (3)
N1—C7—H7	116.7	C17—C16—C13	119.5 (3)
C6—C7—H7	116.7	C18—C17—C16	119.4 (3)
N1—C8—C9	113.9 (3)	C18—C17—H17	120.3

supplementary materials

N1—C8—H8A	108.8	C16—C17—H17	120.3
C9—C8—H8A	108.8	C17—C18—C19	123.2 (3)
N1—C8—H8B	108.8	C17—C18—C14	118.6 (2)
C9—C8—H8B	108.8	C19—C18—C14	118.2 (3)
H8A—C8—H8B	107.7	O2—C19—C14	125.2 (3)
C10—C9—C8	110.5 (3)	O2—C19—C18	119.9 (3)
C10—C9—C12	107.6 (3)	C14—C19—C18	114.9 (3)
C8—C9—C12	111.1 (3)	O1—Cu1—O2	89.91 (10)
C10—C9—C11	110.1 (3)	O1—Cu1—N1	93.14 (11)
C8—C9—C11	107.1 (3)	O2—Cu1—N1	159.14 (11)
C12—C9—C11	110.6 (3)	O1—Cu1—N2	158.73 (10)
C9—C10—H10A	109.5	O2—Cu1—N2	93.67 (10)
C9—C10—H10B	109.5	N1—Cu1—N2	90.94 (11)
H10A—C10—H10B	109.5	C7—N1—C8	118.7 (3)
C9—C10—H10C	109.5	C7—N1—Cu1	125.7 (2)
H10A—C10—H10C	109.5	C8—N1—Cu1	115.6 (2)
H10B—C10—H10C	109.5	C13—N2—C12	119.3 (3)
C9—C11—H11A	109.5	C13—N2—Cu1	125.0 (2)
C9—C11—H11B	109.5	C12—N2—Cu1	115.0 (2)
H11A—C11—H11B	109.5	C1—O1—Cu1	127.5 (2)
C9—C11—H11C	109.5	C19—O2—Cu1	126.8 (2)
O1—C1—C2—C3	179.8 (3)	C15—C14—C19—C18	-1.1 (5)
C6—C1—C2—C3	1.6 (5)	C13—C14—C19—C18	177.7 (3)
O1—C1—C2—C11	-0.7 (4)	C17—C18—C19—O2	179.7 (3)
C6—C1—C2—C11	-178.9 (2)	C14—C18—C19—O2	-0.1 (4)
C1—C2—C3—C4	-1.5 (5)	C17—C18—C19—C14	1.1 (5)
C11—C2—C3—C4	179.0 (3)	C14—C18—C19—C14	-178.7 (2)
C2—C3—C4—C5	0.4 (5)	C6—C7—N1—C8	-174.7 (3)
C2—C3—C4—C12	-178.8 (3)	C6—C7—N1—Cu1	1.0 (5)
C3—C4—C5—C6	0.5 (5)	C9—C8—N1—C7	-112.5 (3)
C12—C4—C5—C6	179.7 (3)	C9—C8—N1—Cu1	71.4 (3)
C4—C5—C6—C1	-0.3 (5)	O1—Cu1—N1—C7	-5.6 (3)
C4—C5—C6—C7	-176.7 (3)	O2—Cu1—N1—C7	-103.6 (4)
O1—C1—C6—C5	-178.7 (3)	N2—Cu1—N1—C7	153.5 (3)
C2—C1—C6—C5	-0.6 (5)	O1—Cu1—N1—C8	170.2 (2)
O1—C1—C6—C7	-2.5 (5)	O2—Cu1—N1—C8	72.2 (4)
C2—C1—C6—C7	175.6 (3)	N2—Cu1—N1—C8	-30.7 (2)
C5—C6—C7—N1	-179.3 (3)	C14—C13—N2—C12	-174.9 (3)
C1—C6—C7—N1	4.3 (5)	C14—C13—N2—Cu1	-4.7 (5)
N1—C8—C9—C10	81.1 (4)	C9—C12—N2—C13	-118.0 (3)
N1—C8—C9—C12	-38.1 (4)	C9—C12—N2—Cu1	70.9 (3)
N1—C8—C9—C11	-159.0 (3)	O1—Cu1—N2—C13	-103.4 (3)
C10—C9—C12—N2	-153.9 (3)	O2—Cu1—N2—C13	-4.1 (3)
C8—C9—C12—N2	-33.0 (4)	N1—Cu1—N2—C13	155.5 (3)
C11—C9—C12—N2	85.8 (4)	O1—Cu1—N2—C12	67.2 (4)
N2—C13—C14—C15	-172.0 (3)	O2—Cu1—N2—C12	166.4 (2)
N2—C13—C14—C19	9.1 (5)	N1—Cu1—N2—C12	-33.9 (2)
C19—C14—C15—C16	0.1 (5)	C6—C1—O1—Cu1	-4.6 (5)
C13—C14—C15—C16	-178.7 (3)	C2—C1—O1—Cu1	177.4 (2)

C14—C15—C16—C17	1.0 (6)	O2—Cu1—O1—C1	166.7 (3)
C14—C15—C16—C13	179.1 (3)	N1—Cu1—O1—C1	7.4 (3)
C15—C16—C17—C18	-1.0 (6)	N2—Cu1—O1—C1	-93.3 (4)
C13—C16—C17—C18	-179.1 (3)	C14—C19—O2—Cu1	-10.9 (5)
C16—C17—C18—C19	-0.1 (5)	C18—C19—O2—Cu1	170.7 (2)
C16—C17—C18—C14	179.7 (3)	O1—Cu1—O2—C19	170.8 (3)
C15—C14—C19—O2	-179.6 (3)	N1—Cu1—O2—C19	-90.6 (4)
C13—C14—C19—O2	-0.8 (5)	N2—Cu1—O2—C19	11.8 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C8—H8A \cdots O1 ⁱ	0.97	2.56	3.331 (4)	136

Symmetry codes: (i) $-x+1, -y+2, -z$.

Fig. 1




Fig. 2

