

(Nitrato- κ^2O,O')bis(triptanthrin- κN)-silver(I)Jie Wu,^a Chao Huang,^b Guo-Qiang Li,^a Hai-Yan Tian^a and Ren-Wang Jiang^{a*}^aGuangdong Province Key Laboratory of Pharmacodynamic Constituents of Traditional Chinese Medicine and New Drugs Research, Institute of Traditional Chinese Medicine and Natural Products, Jinan University, Guangzhou 510632, People's Republic of China, and ^bSinopharmtc Shenzhen Ltd, Shenzhen 518029, People's Republic of China

Correspondence e-mail: trwjiang@jnu.edu.cn


Received 7 October 2011; accepted 15 January 2012

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.043; wR factor = 0.108; data-to-parameter ratio = 13.3.

In the crystal structure of the title compound, $[\text{Ag}(\text{NO}_3)\text{-}(\text{C}_{15}\text{H}_{8}\text{N}_2\text{O}_2)_2]$, triptanthrin (indolo[2,1-*b*]quinazoline-6,12-dione) and silver nitrate form a 2:1 complex. The silver ion is surrounded by two triptanthrin ligands, each coordinating through the N atoms, with $\text{Ag}-\text{N}$ bond lengths of 2.247 (3) and 2.264 (3) Å, and an anionic nitrate ligand coordinating through two O atoms, with $\text{Ag}-\text{O}$ bond lengths of 2.499 (3) and 2.591 (3) Å. The $\text{N}-\text{Ag}-\text{N}$ plane and the $\text{O}-\text{Ag}-\text{O}$ plane are roughly perpendicular, making a dihedral angle of 81.6 (2)°. In the crystal, C—H···O interactions between aromatic H atoms and keto and nitrate O atoms as well as $\pi-\pi$ interactions [centroid-centroid distance = 3.706 (4) Å] give rise to a three-dimensional network.

Related literature

For the biological activity of triptanthrin, see: Yu *et al.* (2007); Chan *et al.* (2009); Bandekar *et al.* (2010). For the synthesis and structural modification of triptanthrin, see: Jao *et al.* (2008); Kumar *et al.* (2011); Chen *et al.* (2011). For related $\pi-\pi$ interactions in natural flavonoids, see: Jiang *et al.* (2002, 2009). For standard bond lengths, see: Allen *et al.* (1987). For bond lengths and angles in a silver nitrate complex with 4,4'-trimethylenedipiperidine, see: Kokunov *et al.* (2011).



Experimental

Crystal data

$[\text{Ag}(\text{NO}_3)(\text{C}_{15}\text{H}_{8}\text{N}_2\text{O}_2)_2]$	$\gamma = 84.447\text{ (4)}^\circ$
$M_r = 666.35$	$V = 1219.0\text{ (5)}\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.0598\text{ (19)}\text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.873\text{ (3)}\text{ \AA}$	$\mu = 0.89\text{ mm}^{-1}$
$c = 14.541\text{ (3)}\text{ \AA}$	$T = 150\text{ K}$
$\alpha = 76.010\text{ (4)}^\circ$	$0.34 \times 0.26 \times 0.22\text{ mm}$
$\beta = 81.019\text{ (4)}^\circ$	

Data collection

Bruker SMART 1000 CCD diffractometer	9843 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2004)	5150 independent reflections
$T_{\min} = 0.626$, $T_{\max} = 1.000$	3893 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$	388 parameters
$wR(F^2) = 0.108$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\max} = 1.08\text{ e \AA}^{-3}$
5150 reflections	$\Delta\rho_{\min} = -0.63\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$C3'-\text{H}3'\cdots O1^{\text{i}}$	0.95	2.52	3.348 (4)	145
$C4'-\text{H}4'\cdots O4^{\text{ii}}$	0.95	2.42	3.300 (3)	153
$C3-\text{H}3\cdots O1^{\text{iii}}$	0.95	2.45	3.139 (4)	130
$C6'-\text{H}6'\cdots O2$	0.95	2.40	3.313 (4)	161
$C4-\text{H}4\cdots O3^{\text{iv}}$	0.95	2.53	3.190 (2)	127
$C6-\text{H}6\cdots O2'$	0.95	2.53	3.454 (3)	164
$C13'-\text{H}13B\cdots O4^{\text{v}}$	0.95	2.53	3.453 (2)	163
$C14-\text{H}14A\cdots O1$	0.95	2.47	2.996 (5)	115
$C14'-\text{H}14B\cdots O1'$	0.95	2.43	2.970 (3)	116

Symmetry codes: (i) $x, y, z - 1$; (ii) $-x, -y + 1, -z + 1$; (iii) $x, y, z + 1$; (iv) $-x + 1, -y, -z + 2$; (v) $-x + 1, -y, -z + 1$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SMART* and *SAINT* (Bruker, 1998); data reduction: *XPREP* (Bruker, 1998); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *XP* in *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by grants from the New Century Excellent Talents Scheme of the Ministry of Education (NCET-08-0612), the National Science Foundation of China (30801433) and the Fundamental Research Funds for the Central Universities (21609202).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RN2096).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bandekar, P. P., Roopnarine, K. A., Parekh, V. J., Mitchell, T. R., Novak, M. J. & Sinden, R. R. (2010). *J. Med. Chem.* **53**, 3558–3565.
- Bruker (1998). *SMART*, *SAINT* and *XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chan, H. L., Yip, H. Y., Mak, N. K. & Leung, K. N. (2009). *Cell. Mol. Immunol.* **6**, 335–342.
- Chen, H. J., Tsao, H. H., Lo, J. G., Chiu, K. H. & Jen, J. F. (2011). *Separ. Sci. Technol.* **46**, 972–977.
- Jao, C. W., Lin, W. C., Wu, Y. T. & Wu, P. L. (2008). *J. Nat. Prod.* **71**, 1275–1279.
- Jiang, R. W., Wang, Y., Gao, H., Zhang, D. M. & Ye, W. C. (2009). *J. Mol. Struct.* **920**, 383–386.
- Jiang, R. W., Ye, W. C., Woo, K. Y., Du, J., Che, C. T., But, P. P. H. & Mak, T. C. W. (2002). *J. Mol. Struct.* **642**, 77–84.
- Kokunov, Y. V., Gorbunova, Y. E. & Kovalev, V. V. (2011). *Russ. J. Inorg. Chem.* **56**, 39–43.
- Kumar, A., Tripathi, V. D. & Kumar, P. (2011). *Green Chem.* **13**, 51–54.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yu, S. T., Chen, T. M., Tseng, S. Y. & Chen, Y. H. (2007). *Biochem. Biophys. Res. Commun.* **358**, 79–84.

supplementary materials

Acta Cryst. (2012). E68, m185-m186 [doi:10.1107/S1600536812001821]

(Nitrato- $\kappa^2 O,O'$)bis(trypanthrin- κN)silver(I)

J. Wu, C. Huang, G.-Q. Li, H.-Y. Tian and R.-W. Jiang

Comment

Tryptanthrin is an indole quinazoline alkaloid isolated from *Folium Isatidis* (Chen *et al.*, 2011). It was found to inhibit the murine myelomonocytic leukemia cells by causing cell cycle arrest and triggering cell differentiation (Chan *et al.*, 2009), and reverse the doxorubicin resistance in breast cancer cells by inhibition of MDR1 (Yu *et al.*, 2007). It was also found to inhibit the growth of *Escherichia coli* through intercalation into DNA (Bandekar *et al.*, 2010). In addition, synthesis of tryptanthrin (Kumar *et al.*, 2011) and its derivatives (Chen *et al.*, 2011) were reported.

Tryptanthrin and silver nitrate form a 2:1 complex. The same proportions were used in the synthesis. Both the two tryptanthrin molecules are essentially planar with mean deviations of 0.0066 (4) Å and 0.0054 (3) Å, respectively, and make a dihedral angle of 2.9 (2)°. The nitrate group makes dihedral angles of 89.5 (3)° and 105.1 (4)° with the two tryptanthrin molecules. The silver ion is coordinated with two oxygen atoms from the anionic nitrate ligand with bond distances of 2.499 (3) Å and 2.591 (3) Å, and two nitrogen atoms from the tryptanthrin ligands with bond distances of 2.266 (3) Å and 2.249 (3) Å, which are slightly longer than those reported in silver nitrate complex with 4,4'-trimethylenedipiperidine [2.192 (5) Å and 2.212 (5) Å] (Kokunov *et al.*, 2011). The N—Ag—N plane and the O—Ag—O plane are roughly perpendicular with a dihedral angle of 81.6 (2)°. The bond distances and bond angles in both tryptanthrin molecules are all normal (Allen *et al.*, 1987).

Short intermolecular C—H···O interactions (Table 1) between the tryptanthrin methane H atoms and the nitrate ligand [C4'—H···O4, 3.300 (3) Å, C13'—H···O4, 3.453 (2) Å] linked adjacent molecules into layers. Adjacent layers were linked by π — π interactions between the benzene rings [centroid-centroid distance 3.706 (4) Å and displacement angle 4.5 (2)°]. The centroid-centroid distance observed in title compound is similar to those in natural flavonoids (Jiang, *et al.*, 2002 and 2009).

Experimental

Tryptanthrin (12.4 mg, 0.05 mmol) was dissolved in a solution including methanol (1 ml) and chloroform (3 ml). silver nitrate (17.0 mg, 0.1 mmol) was dissolved in methanol (1 ml) and was added into the tryptanthrin solution. Three days later, the crystals were obtained *via* slow evaporation at room temperature.

Refinement

The C-bound H atoms were positioned geometrically and were included in the refinement in the riding-model approximation, with C—H = 0.96 Å (CH₃) and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$; 0.97 Å (CH₂) and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$; 0.93 Å (aryl H) and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$; O—H = 0.82 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. There was a residue peak with height of 1.08 and distance of 0.93 Å from Ag after the final refinement.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound showing 30% probability displacement ellipsoids.




Fig. 2. The packing diagram viewed down the *c* axis. The dashed lines represent intermolecular C—H···O interactions.

Bis(indolo[2,1-*b*]quinazoline-6,12-dione-κ*N*)(nitrato- κ²*O,O'*)silver(I)

Crystal data

[Ag(No ₃)(C ₁₅ H ₈ N ₂ O ₂) ₂]	<i>Z</i> = 2
<i>M_r</i> = 666.35	<i>F</i> (000) = 668
Triclinic, <i>P</i> 1	<i>D_x</i> = 1.815 Mg m ⁻³
<i>a</i> = 8.0598 (19) Å	Mo <i>K</i> α radiation, λ = 0.71074 Å
<i>b</i> = 10.873 (3) Å	Cell parameters from 9843 reflections
<i>c</i> = 14.541 (3) Å	θ = 3.2–80.9°
α = 76.010 (4)°	μ = 0.89 mm ⁻¹
β = 81.019 (4)°	<i>T</i> = 150 K
γ = 84.447 (4)°	Block, yellow
<i>V</i> = 1219.0 (5) Å ³	0.34 × 0.26 × 0.22 mm

Data collection

Bruker SMART 1000 CCD	5150 independent reflections
diffractometer	
Radiation source: fine-focus sealed tube	3893 reflections with $I > 2\sigma(I)$
graphite	$R_{\text{int}} = 0.034$
ω scans	$\theta_{\text{max}} = 27.1^\circ$, $\theta_{\text{min}} = 1.5^\circ$
Absorption correction: multi-scan	$h = -10 \rightarrow 10$
(SADABS; Sheldrick, 2004)	$k = -13 \rightarrow 13$
$T_{\text{min}} = 0.626$, $T_{\text{max}} = 1.000$	$l = -18 \rightarrow 18$
9843 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.043$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.108$	H-atom parameters constrained
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0565P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
5150 reflections	$(\Delta/\sigma)_{\max} = 0.001$
388 parameters	$\Delta\rho_{\max} = 1.08 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.63 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.16812 (4)	0.20968 (3)	0.75957 (2)	0.01945 (11)
O1	-0.1263 (4)	0.2779 (3)	1.17705 (19)	0.0228 (7)
O2	-0.0402 (3)	0.4293 (3)	0.76517 (19)	0.0203 (6)
N1	0.0735 (4)	0.2171 (3)	0.9133 (2)	0.0153 (7)
N2	-0.1018 (4)	0.3331 (3)	1.0138 (2)	0.0152 (7)
C1	-0.0672 (5)	0.2537 (4)	1.1009 (3)	0.0181 (9)
C2	0.0488 (5)	0.1456 (4)	1.0870 (3)	0.0159 (8)
C3	0.0992 (5)	0.0596 (4)	1.1675 (3)	0.0206 (9)
H3A	0.0532	0.0691	1.2298	0.025*
C4	0.2162 (5)	-0.0393 (4)	1.1561 (3)	0.0239 (10)
H4A	0.2493	-0.0986	1.2108	0.029*
C5	0.2856 (5)	-0.0522 (4)	1.0649 (3)	0.0229 (9)
H5A	0.3680	-0.1191	1.0579	0.028*
C6	0.2361 (5)	0.0311 (4)	0.9844 (3)	0.0196 (9)
H6A	0.2821	0.0206	0.9223	0.024*
C7	0.1178 (5)	0.1310 (4)	0.9955 (3)	0.0155 (8)
C8	-0.0295 (5)	0.3110 (4)	0.9265 (3)	0.0152 (8)
C9	-0.0852 (5)	0.4188 (4)	0.8498 (3)	0.0181 (9)

supplementary materials

C10	-0.1937 (5)	0.5033 (4)	0.9006 (3)	0.0155 (8)
C11	-0.2774 (5)	0.6174 (4)	0.8666 (3)	0.0206 (9)
H11A	-0.2692	0.6537	0.7998	0.025*
C12	-0.3733 (5)	0.6781 (4)	0.9311 (3)	0.0236 (9)
H12A	-0.4334	0.7566	0.9089	0.028*
C13	-0.3827 (5)	0.6249 (4)	1.0287 (3)	0.0215 (9)
H13A	-0.4495	0.6685	1.0721	0.026*
C14	-0.2977 (5)	0.5102 (4)	1.0649 (3)	0.0198 (9)
H14A	-0.3052	0.4746	1.1317	0.024*
C15	-0.2014 (5)	0.4501 (4)	0.9990 (3)	0.0171 (8)
O1'	0.1922 (4)	0.0444 (3)	0.37176 (19)	0.0238 (7)
O2'	0.3255 (4)	-0.0370 (3)	0.76130 (19)	0.0211 (6)
N1'	0.1282 (4)	0.1466 (3)	0.6288 (2)	0.0151 (7)
N2'	0.2432 (4)	0.0131 (3)	0.5259 (2)	0.0146 (7)
C1'	0.1695 (5)	0.0793 (4)	0.4463 (3)	0.0162 (8)
C2'	0.0650 (5)	0.1903 (4)	0.4645 (3)	0.0147 (8)
C3'	-0.0166 (5)	0.2676 (4)	0.3917 (3)	0.0168 (8)
H3'A	-0.0020	0.2486	0.3304	0.020*
C4'	-0.1183 (5)	0.3713 (4)	0.4083 (3)	0.0189 (9)
H4'A	-0.1730	0.4244	0.3584	0.023*
C5'	-0.1411 (5)	0.3985 (4)	0.4989 (3)	0.0223 (9)
H5'A	-0.2129	0.4695	0.5102	0.027*
C6'	-0.0611 (5)	0.3240 (4)	0.5718 (3)	0.0202 (9)
H6'A	-0.0778	0.3431	0.6332	0.024*
C7'	0.0446 (5)	0.2203 (4)	0.5550 (3)	0.0143 (8)
C8'	0.2189 (5)	0.0498 (4)	0.6114 (3)	0.0146 (8)
C9'	0.3156 (5)	-0.0455 (4)	0.6808 (3)	0.0168 (8)
C10'	0.3883 (5)	-0.1396 (4)	0.6284 (3)	0.0172 (8)
C11'	0.4838 (5)	-0.2517 (4)	0.6561 (3)	0.0212 (9)
H11B	0.5066	-0.2801	0.7203	0.025*
C12'	0.5456 (5)	-0.3218 (4)	0.5889 (3)	0.0234 (9)
H12B	0.6109	-0.3991	0.6070	0.028*
C13'	0.5123 (5)	-0.2793 (4)	0.4945 (3)	0.0223 (9)
H13B	0.5586	-0.3270	0.4488	0.027*
C14'	0.4125 (5)	-0.1686 (4)	0.4660 (3)	0.0188 (9)
H14B	0.3877	-0.1409	0.4022	0.023*
C15'	0.3514 (5)	-0.1012 (4)	0.5342 (3)	0.0152 (8)
N3	0.4813 (4)	0.3383 (3)	0.7421 (2)	0.0211 (8)
O3	0.4698 (4)	0.2218 (3)	0.7760 (3)	0.0418 (9)
O4	0.3608 (4)	0.3988 (3)	0.7064 (2)	0.0355 (8)
O5	0.6079 (4)	0.3907 (3)	0.7475 (2)	0.0393 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.02012 (18)	0.02486 (19)	0.01477 (17)	0.00208 (12)	-0.00300 (12)	-0.00824 (13)
O1	0.0281 (16)	0.0265 (16)	0.0124 (14)	0.0007 (13)	0.0007 (12)	-0.0053 (12)
O2	0.0196 (15)	0.0273 (16)	0.0136 (14)	-0.0003 (13)	-0.0010 (12)	-0.0053 (12)

N1	0.0112 (16)	0.0187 (17)	0.0167 (17)	-0.0020 (14)	-0.0016 (13)	-0.0051 (14)
N2	0.0131 (16)	0.0203 (17)	0.0115 (16)	-0.0012 (14)	0.0017 (13)	-0.0043 (14)
C1	0.018 (2)	0.022 (2)	0.015 (2)	-0.0074 (17)	-0.0030 (17)	-0.0028 (17)
C2	0.016 (2)	0.018 (2)	0.0147 (19)	-0.0034 (16)	-0.0044 (16)	-0.0036 (16)
C3	0.020 (2)	0.025 (2)	0.016 (2)	-0.0025 (18)	-0.0014 (17)	-0.0045 (18)
C4	0.022 (2)	0.027 (2)	0.021 (2)	-0.0049 (19)	-0.0064 (18)	0.0017 (19)
C5	0.017 (2)	0.022 (2)	0.030 (2)	0.0029 (18)	-0.0062 (18)	-0.0070 (19)
C6	0.018 (2)	0.023 (2)	0.019 (2)	-0.0032 (17)	-0.0033 (17)	-0.0063 (18)
C7	0.0135 (19)	0.018 (2)	0.0160 (19)	-0.0045 (16)	-0.0042 (16)	-0.0035 (16)
C8	0.0145 (19)	0.020 (2)	0.0123 (19)	-0.0090 (17)	-0.0009 (15)	-0.0037 (16)
C9	0.0124 (19)	0.023 (2)	0.018 (2)	-0.0071 (17)	-0.0027 (16)	-0.0007 (17)
C10	0.0113 (19)	0.021 (2)	0.016 (2)	-0.0022 (16)	-0.0004 (16)	-0.0079 (17)
C11	0.016 (2)	0.027 (2)	0.019 (2)	-0.0009 (18)	-0.0040 (17)	-0.0050 (18)
C12	0.022 (2)	0.021 (2)	0.031 (2)	-0.0011 (18)	-0.0100 (19)	-0.0066 (19)
C13	0.014 (2)	0.024 (2)	0.027 (2)	-0.0031 (17)	0.0012 (17)	-0.0087 (19)
C14	0.017 (2)	0.022 (2)	0.020 (2)	-0.0067 (17)	0.0038 (17)	-0.0063 (18)
C15	0.015 (2)	0.0149 (19)	0.022 (2)	-0.0061 (16)	-0.0017 (17)	-0.0039 (17)
O1'	0.0287 (17)	0.0304 (17)	0.0143 (14)	0.0009 (13)	-0.0066 (13)	-0.0076 (13)
O2'	0.0225 (15)	0.0246 (16)	0.0161 (15)	0.0026 (12)	-0.0059 (12)	-0.0041 (12)
N1'	0.0143 (16)	0.0175 (17)	0.0127 (16)	-0.0034 (14)	-0.0008 (13)	-0.0018 (14)
N2'	0.0185 (17)	0.0136 (16)	0.0117 (16)	-0.0010 (14)	-0.0022 (14)	-0.0027 (13)
C1'	0.0128 (19)	0.020 (2)	0.017 (2)	-0.0006 (16)	-0.0020 (16)	-0.0056 (17)
C2'	0.0135 (19)	0.0169 (19)	0.0121 (18)	-0.0051 (16)	-0.0008 (15)	0.0005 (16)
C3'	0.0123 (19)	0.023 (2)	0.0137 (19)	-0.0034 (17)	-0.0002 (16)	-0.0027 (17)
C4'	0.0123 (19)	0.021 (2)	0.019 (2)	-0.0041 (17)	-0.0036 (16)	0.0054 (17)
C5'	0.016 (2)	0.016 (2)	0.031 (2)	0.0045 (17)	-0.0034 (18)	-0.0005 (18)
C6'	0.020 (2)	0.021 (2)	0.021 (2)	-0.0048 (18)	-0.0043 (17)	-0.0047 (18)
C7'	0.0108 (18)	0.0157 (19)	0.0151 (19)	-0.0038 (16)	-0.0007 (15)	-0.0010 (16)
C8'	0.0119 (19)	0.017 (2)	0.0139 (19)	-0.0077 (16)	0.0006 (15)	-0.0003 (16)
C9'	0.0132 (19)	0.018 (2)	0.017 (2)	-0.0021 (16)	0.0000 (16)	-0.0006 (17)
C10'	0.0132 (19)	0.022 (2)	0.0155 (19)	-0.0035 (17)	0.0005 (16)	-0.0037 (17)
C11'	0.016 (2)	0.021 (2)	0.025 (2)	0.0013 (17)	-0.0059 (18)	-0.0017 (18)
C12'	0.019 (2)	0.019 (2)	0.031 (2)	0.0002 (18)	-0.0050 (19)	-0.0039 (19)
C13'	0.019 (2)	0.021 (2)	0.028 (2)	-0.0047 (18)	0.0025 (18)	-0.0094 (19)
C14'	0.016 (2)	0.019 (2)	0.021 (2)	-0.0078 (17)	0.0020 (17)	-0.0041 (17)
C15'	0.0147 (19)	0.0148 (19)	0.0155 (19)	-0.0029 (16)	-0.0036 (16)	-0.0007 (16)
N3	0.0160 (18)	0.026 (2)	0.0206 (19)	0.0022 (16)	-0.0019 (15)	-0.0065 (16)
O3	0.0239 (18)	0.0247 (18)	0.071 (3)	0.0026 (14)	-0.0111 (17)	0.0015 (17)
O4	0.0246 (18)	0.0298 (18)	0.047 (2)	0.0059 (14)	-0.0132 (16)	0.0029 (16)
O5	0.0257 (18)	0.048 (2)	0.045 (2)	-0.0132 (16)	-0.0080 (16)	-0.0051 (17)

Geometric parameters (\AA , $^\circ$)

Ag1—N1'	2.247 (3)	O1'—C1'	1.215 (5)
Ag1—N1	2.264 (3)	O2'—C9'	1.212 (5)
Ag1—O3	2.499 (3)	N1'—C8'	1.277 (5)
Ag1—O4	2.591 (3)	N1'—C7'	1.398 (5)
O1—C1	1.214 (5)	N2'—C8'	1.374 (5)
O2—C9	1.209 (5)	N2'—C1'	1.393 (5)

supplementary materials

N1—C8	1.284 (5)	N2'—C15'	1.439 (5)
N1—C7	1.400 (5)	C1'—C2'	1.457 (5)
N2—C8	1.377 (5)	C2'—C3'	1.392 (5)
N2—C1	1.401 (5)	C2'—C7'	1.413 (5)
N2—C15	1.426 (5)	C3'—C4'	1.376 (6)
C1—C2	1.463 (6)	C3'—H3'A	0.9500
C2—C3	1.398 (5)	C4'—C5'	1.400 (6)
C2—C7	1.399 (5)	C4'—H4'A	0.9500
C3—C4	1.384 (6)	C5'—C6'	1.377 (6)
C3—H3A	0.9500	C5'—H5'A	0.9500
C4—C5	1.390 (6)	C6'—C7'	1.393 (5)
C4—H4A	0.9500	C6'—H6'A	0.9500
C5—C6	1.383 (6)	C8'—C9'	1.511 (5)
C5—H5A	0.9500	C9'—C10'	1.447 (5)
C6—C7	1.397 (6)	C10'—C11'	1.386 (6)
C6—H6A	0.9500	C10'—C15'	1.402 (5)
C8—C9	1.497 (5)	C11'—C12'	1.386 (6)
C9—C10	1.458 (5)	C11'—H11B	0.9500
C10—C11	1.373 (6)	C12'—C13'	1.397 (6)
C10—C15	1.401 (5)	C12'—H12B	0.9500
C11—C12	1.373 (6)	C13'—C14'	1.394 (6)
C11—H11A	0.9500	C13'—H13B	0.9500
C12—C13	1.390 (6)	C14'—C15'	1.375 (5)
C12—H12A	0.9500	C14'—H14B	0.9500
C13—C14	1.390 (6)	N3—O4	1.231 (4)
C13—H13A	0.9500	N3—O5	1.237 (4)
C14—C15	1.385 (6)	N3—O3	1.250 (4)
C14—H14A	0.9500		
N1'—Ag1—N1	147.76 (11)	C8'—N1'—C7'	116.9 (3)
N1'—Ag1—O3	114.47 (12)	C8'—N1'—Ag1	116.4 (3)
N1—Ag1—O3	94.10 (12)	C7'—N1'—Ag1	124.8 (2)
N1'—Ag1—O4	108.49 (11)	C8'—N2'—C1'	123.1 (3)
N1—Ag1—O4	101.36 (11)	C8'—N2'—C15'	109.4 (3)
O3—Ag1—O4	49.28 (10)	C1'—N2'—C15'	127.5 (3)
C8—N1—C7	116.5 (3)	O1'—C1'—N2'	121.8 (4)
C8—N1—Ag1	116.7 (3)	O1'—C1'—C2'	125.8 (4)
C7—N1—Ag1	126.7 (2)	N2'—C1'—C2'	112.4 (3)
C8—N2—C1	122.7 (3)	C3'—C2'—C7'	119.5 (4)
C8—N2—C15	109.3 (3)	C3'—C2'—C1'	119.6 (3)
C1—N2—C15	127.9 (3)	C7'—C2'—C1'	120.9 (3)
O1—C1—N2	121.7 (4)	C4'—C3'—C2'	120.2 (4)
O1—C1—C2	126.3 (4)	C4'—C3'—H3'A	119.9
N2—C1—C2	112.0 (3)	C2'—C3'—H3'A	119.9
C3—C2—C7	119.6 (4)	C3'—C4'—C5'	119.9 (4)
C3—C2—C1	118.7 (3)	C3'—C4'—H4'A	120.1
C7—C2—C1	121.6 (3)	C5'—C4'—H4'A	120.1
C4—C3—C2	119.8 (4)	C6'—C5'—C4'	121.0 (4)
C4—C3—H3A	120.1	C6'—C5'—H5'A	119.5
C2—C3—H3A	120.1	C4'—C5'—H5'A	119.5

C3—C4—C5	120.2 (4)	C5'—C6'—C7'	119.4 (4)
C3—C4—H4A	119.9	C5'—C6'—H6'A	120.3
C5—C4—H4A	119.9	C7'—C6'—H6'A	120.3
C6—C5—C4	120.9 (4)	C6'—C7'—N1'	119.0 (3)
C6—C5—H5A	119.6	C6'—C7'—C2'	120.0 (4)
C4—C5—H5A	119.6	N1'—C7'—C2'	121.0 (3)
C5—C6—C7	119.2 (4)	N1'—C8'—N2'	125.6 (4)
C5—C6—H6A	120.4	N1'—C8'—C9'	126.3 (3)
C7—C6—H6A	120.4	N2'—C8'—C9'	108.1 (3)
C6—C7—C2	120.3 (4)	O2'—C9'—C10'	131.1 (4)
C6—C7—N1	118.4 (3)	O2'—C9'—C8'	124.3 (4)
C2—C7—N1	121.2 (3)	C10'—C9'—C8'	104.6 (3)
N1—C8—N2	126.0 (4)	C11'—C10'—C15'	119.7 (4)
N1—C8—C9	125.9 (3)	C11'—C10'—C9'	131.0 (4)
N2—C8—C9	108.1 (3)	C15'—C10'—C9'	109.2 (3)
O2—C9—C10	130.4 (4)	C10'—C11'—C12'	119.0 (4)
O2—C9—C8	124.4 (4)	C10'—C11'—H11B	120.5
C10—C9—C8	105.1 (3)	C12'—C11'—H11B	120.5
C11—C10—C15	121.3 (4)	C11'—C12'—C13'	120.3 (4)
C11—C10—C9	130.6 (4)	C11'—C12'—H12B	119.9
C15—C10—C9	108.1 (3)	C13'—C12'—H12B	119.9
C12—C11—C10	118.6 (4)	C14'—C13'—C12'	121.5 (4)
C12—C11—H11A	120.7	C14'—C13'—H13B	119.2
C10—C11—H11A	120.7	C12'—C13'—H13B	119.2
C11—C12—C13	120.2 (4)	C15'—C14'—C13'	117.2 (4)
C11—C12—H12A	119.9	C15'—C14'—H14B	121.4
C13—C12—H12A	119.9	C13'—C14'—H14B	121.4
C12—C13—C14	122.2 (4)	C14'—C15'—C10'	122.3 (4)
C12—C13—H13A	118.9	C14'—C15'—N2'	129.2 (4)
C14—C13—H13A	118.9	C10'—C15'—N2'	108.5 (3)
C15—C14—C13	116.9 (4)	O4—N3—O5	121.6 (4)
C15—C14—H14A	121.5	O4—N3—O3	117.8 (3)
C13—C14—H14A	121.5	O5—N3—O3	120.6 (4)
C14—C15—C10	120.7 (4)	N3—O3—Ag1	98.2 (2)
C14—C15—N2	129.9 (4)	N3—O4—Ag1	94.3 (2)
C10—C15—N2	109.4 (3)		
N1'—Ag1—N1—C8	-87.6 (3)	O4—Ag1—N1'—C7'	-64.6 (3)
O3—Ag1—N1—C8	119.3 (3)	C8'—N2'—C1'—O1'	-179.5 (4)
O4—Ag1—N1—C8	70.0 (3)	C15'—N2'—C1'—O1'	-0.3 (6)
N1'—Ag1—N1—C7	93.5 (3)	C8'—N2'—C1'—C2'	-0.1 (5)
O3—Ag1—N1—C7	-59.6 (3)	C15'—N2'—C1'—C2'	179.1 (3)
O4—Ag1—N1—C7	-108.8 (3)	O1'—C1'—C2'—C3'	-1.2 (6)
C8—N2—C1—O1	176.6 (4)	N2'—C1'—C2'—C3'	179.4 (3)
C15—N2—C1—O1	1.8 (6)	O1'—C1'—C2'—C7'	178.6 (4)
C8—N2—C1—C2	-1.6 (5)	N2'—C1'—C2'—C7'	-0.8 (5)
C15—N2—C1—C2	-176.4 (3)	C7'—C2'—C3'—C4'	-1.0 (6)
O1—C1—C2—C3	0.0 (6)	C1'—C2'—C3'—C4'	178.8 (3)
N2—C1—C2—C3	178.2 (3)	C2'—C3'—C4'—C5'	-0.6 (6)
O1—C1—C2—C7	-175.9 (4)	C3'—C4'—C5'—C6'	1.0 (6)

supplementary materials

N2—C1—C2—C7	2.3 (5)	C4'—C5'—C6'—C7'	0.3 (6)
C7—C2—C3—C4	-0.3 (6)	C5'—C6'—C7'—N1'	178.7 (3)
C1—C2—C3—C4	-176.3 (4)	C5'—C6'—C7'—C2'	-1.9 (6)
C2—C3—C4—C5	1.0 (6)	C8'—N1'—C7'—C6'	177.5 (3)
C3—C4—C5—C6	-1.6 (6)	Ag1—N1'—C7'—C6'	-18.6 (5)
C4—C5—C6—C7	1.5 (6)	C8'—N1'—C7'—C2'	-2.0 (5)
C5—C6—C7—C2	-0.8 (6)	Ag1—N1'—C7'—C2'	162.0 (3)
C5—C6—C7—N1	177.8 (3)	C3'—C2'—C7'—C6'	2.3 (5)
C3—C2—C7—C6	0.2 (6)	C1'—C2'—C7'—C6'	-177.5 (3)
C1—C2—C7—C6	176.0 (4)	C3'—C2'—C7'—N1'	-178.3 (3)
C3—C2—C7—N1	-178.4 (3)	C1'—C2'—C7'—N1'	1.9 (5)
C1—C2—C7—N1	-2.5 (6)	C7'—N1'—C8'—N2'	1.0 (5)
C8—N1—C7—C6	-176.8 (3)	Ag1—N1'—C8'—N2'	-164.3 (3)
Ag1—N1—C7—C6	2.1 (5)	C7'—N1'—C8'—C9'	-177.9 (3)
C8—N1—C7—C2	1.8 (5)	Ag1—N1'—C8'—C9'	16.8 (5)
Ag1—N1—C7—C2	-179.3 (3)	C1'—N2'—C8'—N1'	0.0 (6)
C7—N1—C8—N2	-1.2 (5)	C15'—N2'—C8'—N1'	-179.3 (3)
Ag1—N1—C8—N2	179.9 (3)	C1'—N2'—C8'—C9'	179.1 (3)
C7—N1—C8—C9	175.6 (3)	C15'—N2'—C8'—C9'	-0.2 (4)
Ag1—N1—C8—C9	-3.3 (5)	N1'—C8'—C9'—O2'	-4.1 (6)
C1—N2—C8—N1	1.2 (6)	N2'—C8'—C9'—O2'	176.8 (4)
C15—N2—C8—N1	176.9 (4)	N1'—C8'—C9'—C10'	176.7 (4)
C1—N2—C8—C9	-176.0 (3)	N2'—C8'—C9'—C10'	-2.4 (4)
C15—N2—C8—C9	-0.4 (4)	O2'—C9'—C10'—C11'	3.8 (7)
N1—C8—C9—O2	0.9 (6)	C8'—C9'—C10'—C11'	-177.0 (4)
N2—C8—C9—O2	178.2 (4)	O2'—C9'—C10'—C15'	-174.9 (4)
N1—C8—C9—C10	-177.1 (4)	C8'—C9'—C10'—C15'	4.2 (4)
N2—C8—C9—C10	0.2 (4)	C15'—C10'—C11'—C12'	2.0 (6)
O2—C9—C10—C11	1.2 (7)	C9'—C10'—C11'—C12'	-176.6 (4)
C8—C9—C10—C11	178.9 (4)	C10'—C11'—C12'—C13'	0.3 (6)
O2—C9—C10—C15	-177.7 (4)	C11'—C12'—C13'—C14'	-2.1 (6)
C8—C9—C10—C15	0.0 (4)	C12'—C13'—C14'—C15'	1.5 (6)
C15—C10—C11—C12	-1.5 (6)	C13'—C14'—C15'—C10'	0.8 (6)
C9—C10—C11—C12	179.7 (4)	C13'—C14'—C15'—N2'	-178.2 (4)
C10—C11—C12—C13	0.8 (6)	C11'—C10'—C15'—C14'	-2.6 (6)
C11—C12—C13—C14	-0.2 (6)	C9'—C10'—C15'—C14'	176.3 (3)
C12—C13—C14—C15	0.2 (6)	C11'—C10'—C15'—N2'	176.6 (3)
C13—C14—C15—C10	-0.8 (6)	C9'—C10'—C15'—N2'	-4.5 (4)
C13—C14—C15—N2	-179.8 (4)	C8'—N2'—C15'—C14'	-177.9 (4)
C11—C10—C15—C14	1.5 (6)	C1'—N2'—C15'—C14'	2.8 (6)
C9—C10—C15—C14	-179.4 (3)	C8'—N2'—C15'—C10'	2.9 (4)
C11—C10—C15—N2	-179.3 (3)	C1'—N2'—C15'—C10'	-176.3 (3)
C9—C10—C15—N2	-0.3 (4)	O4—N3—O3—Ag1	-7.0 (4)
C8—N2—C15—C14	179.5 (4)	O5—N3—O3—Ag1	170.6 (3)
C1—N2—C15—C14	-5.2 (6)	N1'—Ag1—O3—N3	97.8 (3)
C8—N2—C15—C10	0.4 (4)	N1—Ag1—O3—N3	-97.6 (3)
C1—N2—C15—C10	175.8 (4)	O4—Ag1—O3—N3	3.9 (2)
N1—Ag1—N1'—C8'	-103.8 (3)	O5—N3—O4—Ag1	-170.9 (3)
O3—Ag1—N1'—C8'	46.6 (3)	O3—N3—O4—Ag1	6.7 (4)

O4—Ag1—N1'—C8'	99.4 (3)	N1'—Ag1—O4—N3	-110.7 (2)
N1—Ag1—N1'—C7'	92.3 (3)	N1—Ag1—O4—N3	81.7 (3)
O3—Ag1—N1'—C7'	-117.4 (3)	O3—Ag1—O4—N3	-3.9 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C3'—H3'A···O1 ⁱ	0.95	2.52	3.348 (4)	145
C4'—H4'A···O4 ⁱⁱ	0.95	2.42	3.300 (3)	153
C3—H3A···O1 ⁱⁱⁱ	0.95	2.45	3.139 (4)	130
C6'—H6'A···O2	0.95	2.40	3.313 (4)	161
C4—H4A···O3 ^{iv}	0.95	2.53	3.190 (2)	127
C6—H6A···O2'	0.95	2.53	3.454 (3)	164
C13'—H13B···O4 ^v	0.95	2.53	3.453 (2)	163
C14—H14A···O1	0.95	2.47	2.996 (5)	115
C14'—H14B···O1'	0.95	2.43	2.970 (3)	116

Symmetry codes: (i) $x, y, z-1$; (ii) $-x, -y+1, -z+1$; (iii) $x, y, z+1$; (iv) $-x+1, -y, -z+2$; (v) $-x+1, -y, -z+1$.

supplementary materials

Fig. 1




Fig. 2

