

catena-Poly[[[bis[μ -3-(4-carboxyphenoxy)propionato]- κ^3 O¹,O^{1'}:O¹; κ^3 O¹:O¹,-O^{1'}-bis[aqua(N,N-dimethylformamide- κ O)cadmium]]- μ -4,4'-bipyridine- κ^2 N:N'] dinitrate]

Shan Gao^a and Seik Weng Ng^{b,c*}

^aKey Laboratory of Functional Inorganic Materials Chemistry, Ministry of Education, Heilongjiang University, Harbin 150080, People's Republic of China; ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: seiweng@um.edu.my


Received 24 January 2012; accepted 24 January 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.030; wR factor = 0.077; data-to-parameter ratio = 16.9.

In the title coordination polymer, $\{[Cd_2(C_{10}H_9O_5)_2(C_{10}H_8N_2)_2(C_3H_7NO)_2(H_2O)_2](NO_3)_2\}_n$, the 3-(4-carboxyphenoxy)-propionate monoanion O,O' -chelates to a Cd^{II} cation through the aliphatic carboxylate end. One of these O atoms is also connected to the metal cation from an inversion-related metal atom. The five O atoms bonded to the metal centre form a pentagon, above and below which are located the N atoms of the 4,4'-bipyridine molecules. The polycationic ribbon propagates along the b axis of the unit cell. The (aromatic) carboxyl end of the monoanion connects adjacent ribbons into a layer motif in the (102) plane. The nitrate ions are hydrogen bonded to the layer. The geometry of the Cd^{II} atom is a *trans*-N₂O₅Cd pentagonal bipyramidal.

Related literature

For 3-(4-carboxyphenoxy)propionic acid, see: Gao & Ng (2006).



Experimental

Crystal data

$[Cd_2(C_{10}H_9O_5)_2(C_{10}H_8N_2)_2(C_3H_7NO)_2(H_2O)_2](NO_3)_2$	$\beta = 84.2052$ (16) $^\circ$
$M_r = 1261.76$	$\gamma = 76.9358$ (16) $^\circ$
Triclinic, $P\bar{1}$	$V = 1292.84$ (10) Å ³
$a = 9.1020$ (5) Å	$Z = 1$
$b = 11.6866$ (5) Å	Mo $K\alpha$ radiation
$c = 13.3534$ (5) Å	$\mu = 0.91$ mm ⁻¹
$\alpha = 69.1646$ (11) $^\circ$	$T = 293$ K
	$0.19 \times 0.12 \times 0.11$ mm

Data collection

Rigaku R-AXIS RAPID IP diffractometer	12594 measured reflections
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	5822 independent reflections
$T_{\min} = 0.847$, $T_{\max} = 0.907$	4760 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$	345 parameters
$wR(F^2) = 0.077$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\max} = 0.95$ e Å ⁻³
5822 reflections	$\Delta\rho_{\min} = -0.59$ e Å ⁻³

Table 1
Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
O5—H5···O4 ⁱ	0.84	1.80	2.629 (3)	171
O1W—H11···O7	0.84	1.94	2.754 (4)	164
O1W—H12···O2 ⁱⁱ	0.84	2.03	2.763 (3)	145

Symmetry codes: (i) $-x + 2, -y + 1, -z$; (ii) $-x + 1, -y + 2, -z + 1$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

This work is supported by the Key Project of the Natural Science Foundation of Heilongjiang Province (grant No. ZD200903), the Key Project of the Education Bureau of Heilongjiang Province (grants No. 12511z023, No. 2011CJHB006), the Innovation Team of the Education Bureau of Heilongjiang Province (grant No. 2010t d03), Heilongjiang University (Hdtd2010-04) and the Ministry of Higher Education of Malaysia (grant No. UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5796).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Gao, S. & Ng, S. W. (2006). *Acta Cryst. E62*, o3420–o3421.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, m214 [doi:10.1107/S1600536812002991]

catena-Poly[[[bis[μ -3-(4-carboxyphenoxy)propionato]- $\kappa^3 O^1,O^{1'}:O^1,\kappa^3 O^1:O^1,O^{1'}\text{-bis[aqua}(N,N\text{-dimethylformamide-}\kappa O\text{)cadmium}]]-\mu\text{-4,4'\text{-bipyridine-}\kappa^2 N:N'] dinitrate]$

S. Gao and S. W. Ng

Comment

We reported the crystal structure of 3-(4-carboxyphenoxy)propionic acid (Gao & Ng, 2006). We also reported the crystal structures of some metal derivatives. In the coordination polymer, $[\text{Cd}_2(\text{H}_2\text{O})_2(\text{C}_{10}\text{H}_8\text{N}_2)_2(\text{DMF})_2(\text{C}_{10}\text{H}_8\text{O}_5)_2]_n 2n(\text{NO}_3)$ (Scheme I), the 3-(4-carboxyphenoxy)propionate monoanion O,O' -chelates to a Cd^{II} atom through the aliphatic (negatively-charged) carboxyl $-\text{CO}_2$ end; one of the O atoms is also connected to an inversion-related metal atom. The carboxyl O atoms, the carboxyl O atom of an inversion-related dianion, and the O atoms of the water and DMF molecules comprise a pentagon, above and below which are located the N atoms of the 4,4'-bipyridine heterocycle (Fig. 1). The polycationic ribbon propagates along the *b*-axis of the unit cell; the (aromatic) carboxylic acid end of the monoanion connects adjacent ribbons (Fig. 2) into a layer motif; the nitrate ions are hydrogen bonded to the layer (Table 1).

Experimental

Cadmium nitrate and 3-(4-carboxyphenoxy)propionic acid (1 mmol) were mixed in a water-DMF (3/1) mixture (10 ml). 4,4'-Bipyridine dissolved in methanol (5 ml) was added. The mixture was stirred until the reactants dissolved. Yellow crystals were isolated after a few days.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93–0.97 Å) and were included in the refinement in the riding model approximation, with *U*(H) set to 1.2–1.5*U*(C). The acid and water H-atoms were similarly treated (O–H 0.84 Å) and their displacement parameters were similarly tied.

Omitted owing to bad disagreement, the (0 2 0), (10 2 8), (9 2 11), (4 0 13), (4 - 1 14), (1 - 1 14), (10 4 9), (3 - 2 13), (2 - 2 13), (9 3 11), (10 2 9) and (10 5 10) reflections were omitted from refinement.

Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of a portion of polymeric $[\text{Cd}_2(\text{H}_2\text{O})_2(\text{C}_{10}\text{H}_8\text{N}_2)_2(\text{C}_3\text{H}_7\text{NO})_2(\text{C}_{10}\text{H}_8\text{O}_5)_2]_n 2n(\text{NO}_3)$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

supplementary materials




Fig. 2. Polycationic ribbon motif.

catena-Poly[[[bis[μ -3-(4-carboxyphenoxy)propionato]- $\kappa^3O^1,O^{1\prime}:O^1;\kappa^3O^{1\prime}:O^1,O^{1\prime}$ -bis[aqua(N,N -dimethylformamide- κO)cadmium]]- μ -4,4'-bipyridine- $\kappa^2N:N'$] dinitrate]

Crystal data

[Cd₂(C₁₀H₉O₅)₂(C₁₀H₈N₂)₂(C₃H₇NO)₂(H₂O)₂](NO₃) $Z = 1$

$M_r = 1261.76$

$F(000) = 640$

Triclinic, $P\bar{1}$

$D_x = 1.621 \text{ Mg m}^{-3}$

Hall symbol: -P 1

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

$a = 9.1020 (5) \text{ \AA}$

Cell parameters from 10397 reflections

$b = 11.6866 (5) \text{ \AA}$

$\theta = 3.2\text{--}27.5^\circ$

$c = 13.3534 (5) \text{ \AA}$

$\mu = 0.91 \text{ mm}^{-1}$

$\alpha = 69.1646 (11)^\circ$

$T = 293 \text{ K}$

$\beta = 84.2052 (16)^\circ$

Prism, yellow

$\gamma = 76.9358 (16)^\circ$

$0.19 \times 0.12 \times 0.11 \text{ mm}$

$V = 1292.84 (10) \text{ \AA}^3$

Data collection

Rigaku R-AXIS RAPID IP
diffractometer

5822 independent reflections

Radiation source: fine-focus sealed tube
graphite

4760 reflections with $I > 2\sigma(I)$

ω scan

$R_{\text{int}} = 0.031$

Absorption correction: multi-scan
(*ABSCOR*; Higashi, 1995)

$\theta_{\max} = 27.5^\circ, \theta_{\min} = 3.2^\circ$

$T_{\min} = 0.847, T_{\max} = 0.907$

$h = -11 \rightarrow 11$

12594 measured reflections

$k = -15 \rightarrow 14$

$l = -17 \rightarrow 16$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.030$

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.077$

H-atom parameters constrained

$S = 1.08$

$w = 1/[\sigma^2(F_o^2) + (0.0367P)^2 + 0.2058P]$

5822 reflections

where $P = (F_o^2 + 2F_c^2)/3$

345 parameters

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.95 \text{ e \AA}^{-3}$

0 restraints

 $\Delta\rho_{\min} = -0.59 \text{ e } \text{\AA}^{-3}$ *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.18002 (2)	0.944005 (16)	0.589093 (15)	0.02562 (7)
O1	0.0904 (2)	1.01688 (18)	0.40717 (15)	0.0361 (4)
O2	0.3311 (2)	1.00556 (19)	0.41763 (15)	0.0390 (5)
O3	0.3901 (3)	0.8936 (2)	0.23010 (18)	0.0493 (6)
O4	0.8546 (3)	0.4867 (2)	0.0878 (2)	0.0620 (7)
O5	0.9522 (3)	0.6549 (2)	0.0014 (2)	0.0741 (8)
H5	1.0193	0.6153	-0.0288	0.111*
O6	0.1426 (3)	0.8763 (2)	0.77857 (16)	0.0479 (5)
O7	0.5403 (4)	0.6153 (3)	0.6982 (3)	0.0893 (10)
O8	0.7250 (5)	0.6396 (4)	0.7677 (4)	0.1299 (16)
O9	0.7435 (3)	0.4764 (3)	0.7307 (3)	0.0829 (9)
O1W	0.4333 (2)	0.8691 (2)	0.64542 (17)	0.0486 (5)
H11	0.4611	0.7941	0.6493	0.073*
H12	0.4890	0.9124	0.6010	0.073*
N1	0.1779 (3)	0.74731 (19)	0.58879 (18)	0.0307 (5)
N2	0.1773 (3)	0.1380 (2)	0.59570 (18)	0.0315 (5)
N3	0.2606 (4)	0.7602 (3)	0.9348 (2)	0.0646 (9)
N4	0.6706 (4)	0.5765 (3)	0.7332 (3)	0.0596 (8)
C1	0.2171 (3)	1.0305 (2)	0.3627 (2)	0.0282 (5)
C2	0.2299 (4)	1.0791 (3)	0.2418 (2)	0.0423 (7)
H2A	0.2158	1.1695	0.2170	0.051*
H2B	0.1497	1.0586	0.2130	0.051*
C3	0.3801 (4)	1.0261 (3)	0.1987 (2)	0.0440 (7)
H3A	0.3853	1.0634	0.1213	0.053*
H3B	0.4624	1.0431	0.2284	0.053*
C4	0.5115 (4)	0.8267 (3)	0.1927 (2)	0.0413 (7)
C5	0.5131 (4)	0.7005 (3)	0.2196 (2)	0.0426 (7)
H5A	0.4373	0.6657	0.2639	0.051*
C6	0.6268 (4)	0.6263 (3)	0.1807 (2)	0.0412 (7)
H6	0.6275	0.5416	0.1990	0.049*
C7	0.7403 (4)	0.6773 (3)	0.1143 (2)	0.0415 (7)
C8	0.7405 (4)	0.8018 (3)	0.0913 (3)	0.0515 (8)
H8	0.8178	0.8360	0.0485	0.062*
C9	0.6276 (4)	0.8772 (3)	0.1307 (3)	0.0514 (8)
H9	0.6299	0.9608	0.1156	0.062*
C10	0.8552 (4)	0.5984 (3)	0.0665 (3)	0.0481 (8)
C11	0.0810 (3)	0.6805 (2)	0.6513 (2)	0.0339 (6)
H11A	0.0148	0.7150	0.6957	0.041*
C12	0.0742 (3)	0.5627 (2)	0.6532 (2)	0.0330 (6)
H12A	0.0044	0.5198	0.6978	0.040*
C13	0.1722 (3)	0.5090 (2)	0.5881 (2)	0.0279 (5)
C14	0.2714 (3)	0.5799 (2)	0.5215 (2)	0.0366 (6)
H14	0.3376	0.5486	0.4752	0.044*

supplementary materials

C15	0.2709 (3)	0.6963 (3)	0.5248 (2)	0.0362 (6)
H15	0.3385	0.7419	0.4802	0.043*
C16	0.0616 (3)	0.2007 (3)	0.6371 (2)	0.0376 (7)
H16	-0.0188	0.1622	0.6681	0.045*
C17	0.0552 (3)	0.3203 (3)	0.6364 (2)	0.0381 (7)
H17	-0.0282	0.3605	0.6661	0.046*
C18	0.1735 (3)	0.3800 (2)	0.5914 (2)	0.0278 (5)
C19	0.2945 (4)	0.3138 (3)	0.5491 (3)	0.0428 (7)
H19	0.3768	0.3498	0.5180	0.051*
C20	0.2921 (4)	0.1946 (3)	0.5536 (3)	0.0448 (8)
H20	0.3748	0.1515	0.5258	0.054*
C21	0.2327 (4)	0.7857 (3)	0.8328 (3)	0.0483 (8)
H21	0.2858	0.7304	0.7998	0.058*
C22	0.1831 (6)	0.8376 (5)	0.9937 (4)	0.0898 (16)
H22A	0.1137	0.9066	0.9477	0.135*
H22B	0.2548	0.8687	1.0197	0.135*
H22C	0.1286	0.7895	1.0532	0.135*
C23	0.3709 (7)	0.6488 (6)	0.9891 (4)	0.118 (2)
H23A	0.4105	0.6049	0.9405	0.177*
H23B	0.3234	0.5955	1.0496	0.177*
H23C	0.4516	0.6725	1.0131	0.177*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.02636 (11)	0.01654 (10)	0.03726 (12)	-0.00540 (7)	-0.00062 (7)	-0.01261 (7)
O1	0.0260 (11)	0.0386 (11)	0.0424 (11)	-0.0067 (9)	0.0057 (8)	-0.0140 (9)
O2	0.0319 (12)	0.0424 (12)	0.0433 (11)	-0.0115 (9)	0.0017 (9)	-0.0139 (9)
O3	0.0467 (14)	0.0443 (13)	0.0622 (14)	-0.0143 (11)	0.0238 (11)	-0.0283 (11)
O4	0.0743 (19)	0.0414 (14)	0.0691 (16)	-0.0119 (13)	0.0278 (13)	-0.0256 (12)
O5	0.071 (2)	0.0525 (16)	0.101 (2)	-0.0199 (14)	0.0489 (16)	-0.0380 (14)
O6	0.0440 (14)	0.0504 (14)	0.0415 (12)	-0.0020 (11)	-0.0028 (10)	-0.0107 (10)
O7	0.053 (2)	0.0577 (19)	0.136 (3)	-0.0091 (15)	-0.0075 (18)	-0.0079 (17)
O8	0.117 (3)	0.103 (3)	0.207 (5)	-0.016 (3)	-0.031 (3)	-0.094 (3)
O9	0.0556 (19)	0.0490 (17)	0.140 (3)	-0.0094 (14)	0.0235 (17)	-0.0346 (17)
O1W	0.0314 (12)	0.0417 (13)	0.0651 (14)	-0.0108 (10)	-0.0036 (10)	-0.0068 (10)
N1	0.0271 (13)	0.0192 (11)	0.0485 (13)	-0.0045 (9)	0.0000 (10)	-0.0154 (9)
N2	0.0354 (14)	0.0206 (11)	0.0442 (13)	-0.0074 (10)	0.0013 (10)	-0.0176 (9)
N3	0.067 (2)	0.085 (2)	0.0376 (16)	-0.0242 (19)	-0.0111 (14)	-0.0075 (15)
N4	0.048 (2)	0.0435 (18)	0.082 (2)	-0.0156 (15)	0.0143 (16)	-0.0162 (16)
C1	0.0261 (15)	0.0209 (13)	0.0382 (14)	-0.0021 (10)	0.0038 (11)	-0.0136 (10)
C2	0.0413 (19)	0.0419 (18)	0.0391 (16)	-0.0022 (14)	0.0030 (13)	-0.0133 (13)
C3	0.045 (2)	0.0462 (19)	0.0393 (16)	-0.0087 (15)	0.0126 (13)	-0.0168 (13)
C4	0.0386 (18)	0.0471 (19)	0.0437 (16)	-0.0108 (14)	0.0100 (13)	-0.0241 (14)
C5	0.046 (2)	0.0458 (19)	0.0416 (16)	-0.0171 (15)	0.0112 (13)	-0.0201 (14)
C6	0.0432 (19)	0.0402 (17)	0.0424 (16)	-0.0103 (14)	0.0013 (13)	-0.0165 (13)
C7	0.0402 (19)	0.0414 (18)	0.0432 (16)	-0.0053 (14)	0.0039 (13)	-0.0178 (13)
C8	0.043 (2)	0.048 (2)	0.064 (2)	-0.0125 (16)	0.0207 (16)	-0.0238 (16)

C9	0.053 (2)	0.0385 (18)	0.064 (2)	-0.0141 (16)	0.0206 (16)	-0.0226 (15)
C10	0.052 (2)	0.044 (2)	0.0466 (18)	-0.0062 (16)	0.0070 (15)	-0.0173 (14)
C11	0.0347 (16)	0.0250 (14)	0.0473 (16)	-0.0079 (12)	0.0073 (12)	-0.0198 (12)
C12	0.0342 (16)	0.0243 (14)	0.0431 (15)	-0.0092 (12)	0.0067 (12)	-0.0146 (11)
C13	0.0296 (15)	0.0186 (13)	0.0395 (14)	-0.0058 (10)	-0.0031 (11)	-0.0135 (10)
C14	0.0381 (17)	0.0232 (14)	0.0516 (17)	-0.0076 (12)	0.0113 (13)	-0.0192 (12)
C15	0.0321 (16)	0.0223 (14)	0.0550 (17)	-0.0101 (12)	0.0089 (13)	-0.0142 (12)
C16	0.0368 (17)	0.0273 (15)	0.0560 (18)	-0.0141 (13)	0.0092 (13)	-0.0212 (13)
C17	0.0345 (17)	0.0276 (15)	0.0587 (18)	-0.0085 (12)	0.0114 (13)	-0.0247 (13)
C18	0.0285 (15)	0.0208 (13)	0.0392 (14)	-0.0050 (10)	-0.0014 (11)	-0.0161 (11)
C19	0.0383 (18)	0.0313 (16)	0.069 (2)	-0.0170 (13)	0.0179 (15)	-0.0288 (14)
C20	0.0414 (19)	0.0325 (16)	0.072 (2)	-0.0119 (14)	0.0160 (15)	-0.0332 (15)
C21	0.045 (2)	0.052 (2)	0.0461 (18)	-0.0156 (16)	0.0003 (15)	-0.0119 (15)
C22	0.113 (4)	0.113 (4)	0.062 (3)	-0.055 (3)	0.011 (3)	-0.037 (3)
C23	0.119 (5)	0.124 (5)	0.078 (3)	-0.006 (4)	-0.051 (3)	0.007 (3)

Geometric parameters (Å, °)

Cd1—O1	2.4304 (19)	C4—C9	1.380 (4)
Cd1—O1 ⁱ	2.398 (2)	C4—C5	1.386 (4)
Cd1—O2	2.5080 (19)	C5—C6	1.378 (4)
Cd1—O6	2.381 (2)	C5—H5A	0.9300
Cd1—O1W	2.370 (2)	C6—C7	1.388 (4)
Cd1—N1	2.305 (2)	C6—H6	0.9300
Cd1—N2 ⁱⁱ	2.294 (2)	C7—C8	1.376 (4)
O1—C1	1.259 (3)	C7—C10	1.482 (4)
O1—Cd1 ⁱ	2.3985 (19)	C8—C9	1.387 (4)
O2—C1	1.251 (3)	C8—H8	0.9300
O3—C4	1.363 (4)	C9—H9	0.9300
O3—C3	1.436 (4)	C11—C12	1.384 (3)
O4—C10	1.235 (4)	C11—H11A	0.9300
O5—C10	1.293 (4)	C12—C13	1.387 (4)
O5—H5	0.8400	C12—H12A	0.9300
O6—C21	1.225 (4)	C13—C14	1.394 (4)
O7—N4	1.245 (4)	C13—C18	1.490 (3)
O8—N4	1.206 (4)	C14—C15	1.376 (4)
O9—N4	1.218 (4)	C14—H14	0.9300
O1W—H11	0.8400	C15—H15	0.9300
O1W—H12	0.8400	C16—C17	1.382 (4)
N1—C11	1.335 (4)	C16—H16	0.9300
N1—C15	1.338 (3)	C17—C18	1.385 (4)
N2—C20	1.332 (4)	C17—H17	0.9300
N2—C16	1.333 (4)	C18—C19	1.389 (4)
N2—Cd1 ⁱⁱⁱ	2.294 (2)	C19—C20	1.379 (4)
N3—C21	1.324 (4)	C19—H19	0.9300
N3—C22	1.428 (5)	C20—H20	0.9300
N3—C23	1.453 (6)	C21—H21	0.9300
C1—C2	1.510 (4)	C22—H22A	0.9600

supplementary materials

C2—C3	1.515 (4)	C22—H22B	0.9600
C2—H2A	0.9700	C22—H22C	0.9600
C2—H2B	0.9700	C23—H23A	0.9600
C3—H3A	0.9700	C23—H23B	0.9600
C3—H3B	0.9700	C23—H23C	0.9600
N2 ⁱⁱ —Cd1—N1	177.79 (8)	C6—C5—H5A	120.0
N2 ⁱⁱ —Cd1—O1W	92.00 (8)	C4—C5—H5A	120.0
N1—Cd1—O1W	88.38 (8)	C5—C6—C7	120.4 (3)
N2 ⁱⁱ —Cd1—O6	87.14 (8)	C5—C6—H6	119.8
N1—Cd1—O6	90.79 (8)	C7—C6—H6	119.8
O1W—Cd1—O6	79.50 (8)	C8—C7—C6	119.0 (3)
N2 ⁱⁱ —Cd1—O1 ⁱ	91.36 (8)	C8—C7—C10	121.4 (3)
N1—Cd1—O1 ⁱ	87.59 (7)	C6—C7—C10	119.7 (3)
O1W—Cd1—O1 ⁱ	161.18 (7)	C7—C8—C9	121.2 (3)
O6—Cd1—O1 ⁱ	82.19 (7)	C7—C8—H8	119.4
N2 ⁱⁱ —Cd1—O1	96.38 (7)	C9—C8—H8	119.4
N1—Cd1—O1	85.10 (7)	C4—C9—C8	119.2 (3)
O1W—Cd1—O1	127.31 (7)	C4—C9—H9	120.4
O6—Cd1—O1	152.60 (8)	C8—C9—H9	120.4
O1 ⁱ —Cd1—O1	70.59 (7)	O4—C10—O5	123.4 (3)
N2 ⁱⁱ —Cd1—O2	87.11 (7)	O4—C10—C7	121.0 (3)
N1—Cd1—O2	95.10 (7)	O5—C10—C7	115.6 (3)
O1W—Cd1—O2	76.40 (7)	N1—C11—C12	123.3 (3)
O6—Cd1—O2	154.98 (8)	N1—C11—H11A	118.4
O1 ⁱ —Cd1—O2	122.27 (6)	C12—C11—H11A	118.4
O1—Cd1—O2	52.37 (6)	C11—C12—C13	119.5 (3)
C1—O1—Cd1 ⁱ	153.61 (18)	C11—C12—H12A	120.3
C1—O1—Cd1	95.24 (16)	C13—C12—H12A	120.3
Cd1 ⁱ —O1—Cd1	109.41 (7)	C12—C13—C14	117.1 (2)
C1—O2—Cd1	91.80 (16)	C12—C13—C18	121.2 (2)
C4—O3—C3	117.4 (2)	C14—C13—C18	121.7 (2)
C10—O5—H5	120.0	C15—C14—C13	119.6 (3)
C21—O6—Cd1	117.7 (2)	C15—C14—H14	120.2
Cd1—O1W—H11	109.5	C13—C14—H14	120.2
Cd1—O1W—H12	109.5	N1—C15—C14	123.3 (3)
H11—O1W—H12	109.5	N1—C15—H15	118.4
C11—N1—C15	117.2 (2)	C14—C15—H15	118.4
C11—N1—Cd1	121.05 (17)	N2—C16—C17	123.2 (3)
C15—N1—Cd1	121.72 (18)	N2—C16—H16	118.4
C20—N2—C16	117.1 (2)	C17—C16—H16	118.4
C20—N2—Cd1 ⁱⁱⁱ	120.16 (18)	C16—C17—C18	119.8 (3)
C16—N2—Cd1 ⁱⁱⁱ	122.71 (18)	C16—C17—H17	120.1
C21—N3—C22	121.9 (4)	C18—C17—H17	120.1
C21—N3—C23	119.2 (4)	C17—C18—C19	116.8 (2)
C22—N3—C23	119.0 (4)	C17—C18—C13	122.0 (2)
O8—N4—O9	120.3 (4)	C19—C18—C13	121.2 (2)

O8—N4—O7	119.6 (4)	C20—C19—C18	119.7 (3)
O9—N4—O7	120.1 (4)	C20—C19—H19	120.2
O2—C1—O1	120.6 (2)	C18—C19—H19	120.2
O2—C1—C2	120.1 (2)	N2—C20—C19	123.4 (3)
O1—C1—C2	119.3 (2)	N2—C20—H20	118.3
C1—C2—C3	113.3 (3)	C19—C20—H20	118.3
C1—C2—H2A	108.9	O6—C21—N3	125.3 (4)
C3—C2—H2A	108.9	O6—C21—H21	117.4
C1—C2—H2B	108.9	N3—C21—H21	117.4
C3—C2—H2B	108.9	N3—C22—H22A	109.5
H2A—C2—H2B	107.7	N3—C22—H22B	109.5
O3—C3—C2	106.8 (2)	H22A—C22—H22B	109.5
O3—C3—H3A	110.4	N3—C22—H22C	109.5
C2—C3—H3A	110.4	H22A—C22—H22C	109.5
O3—C3—H3B	110.4	H22B—C22—H22C	109.5
C2—C3—H3B	110.4	N3—C23—H23A	109.5
H3A—C3—H3B	108.6	N3—C23—H23B	109.5
O3—C4—C9	124.3 (3)	H23A—C23—H23B	109.5
O3—C4—C5	115.8 (3)	N3—C23—H23C	109.5
C9—C4—C5	120.0 (3)	H23A—C23—H23C	109.5
C6—C5—C4	120.1 (3)	H23B—C23—H23C	109.5
N2 ⁱⁱ —Cd1—O1—C1	81.21 (16)	C3—O3—C4—C9	2.8 (5)
N1—Cd1—O1—C1	-100.43 (16)	C3—O3—C4—C5	-176.5 (3)
O1W—Cd1—O1—C1	-16.23 (19)	O3—C4—C5—C6	176.5 (3)
O6—Cd1—O1—C1	177.34 (16)	C9—C4—C5—C6	-2.8 (5)
O1 ⁱ —Cd1—O1—C1	170.41 (19)	C4—C5—C6—C7	-0.1 (5)
O2—Cd1—O1—C1	-0.14 (14)	C5—C6—C7—C8	2.3 (5)
N2 ⁱⁱ —Cd1—O1—Cd1 ⁱ	-89.19 (9)	C5—C6—C7—C10	-175.3 (3)
N1—Cd1—O1—Cd1 ⁱ	89.16 (9)	C6—C7—C8—C9	-1.7 (5)
O1W—Cd1—O1—Cd1 ⁱ	173.37 (7)	C10—C7—C8—C9	175.9 (3)
O6—Cd1—O1—Cd1 ⁱ	6.93 (19)	O3—C4—C9—C8	-175.8 (3)
O1 ⁱ —Cd1—O1—Cd1 ⁱ	0.0	C5—C4—C9—C8	3.4 (5)
O2—Cd1—O1—Cd1 ⁱ	-170.55 (12)	C7—C8—C9—C4	-1.2 (6)
N2 ⁱⁱ —Cd1—O2—C1	-100.20 (16)	C8—C7—C10—O4	179.8 (3)
N1—Cd1—O2—C1	79.95 (16)	C6—C7—C10—O4	-2.6 (5)
O1W—Cd1—O2—C1	167.04 (16)	C8—C7—C10—O5	-1.0 (5)
O6—Cd1—O2—C1	-177.12 (17)	C6—C7—C10—O5	176.6 (3)
O1 ⁱ —Cd1—O2—C1	-10.41 (17)	C15—N1—C11—C12	0.7 (4)
O1—Cd1—O2—C1	0.14 (14)	Cd1—N1—C11—C12	179.3 (2)
N2 ⁱⁱ —Cd1—O6—C21	-125.3 (2)	N1—C11—C12—C13	0.3 (4)
N1—Cd1—O6—C21	55.5 (2)	C11—C12—C13—C14	-1.4 (4)
O1W—Cd1—O6—C21	-32.7 (2)	C11—C12—C13—C18	177.1 (3)
O1 ⁱ —Cd1—O6—C21	142.9 (2)	C12—C13—C14—C15	1.5 (4)
O1—Cd1—O6—C21	136.3 (2)	C18—C13—C14—C15	-177.0 (3)
O2—Cd1—O6—C21	-48.4 (3)	C11—N1—C15—C14	-0.6 (4)
O1W—Cd1—N1—C11	119.7 (2)	Cd1—N1—C15—C14	-179.2 (2)

supplementary materials

O6—Cd1—N1—C11	40.2 (2)	C13—C14—C15—N1	-0.5 (5)
O1 ⁱ —Cd1—N1—C11	-41.9 (2)	C20—N2—C16—C17	1.1 (5)
O1—Cd1—N1—C11	-112.6 (2)	Cd1 ⁱⁱⁱ —N2—C16—C17	-176.5 (2)
O2—Cd1—N1—C11	-164.1 (2)	N2—C16—C17—C18	-0.3 (5)
O1W—Cd1—N1—C15	-61.8 (2)	C16—C17—C18—C19	-0.3 (4)
O6—Cd1—N1—C15	-141.2 (2)	C16—C17—C18—C13	179.7 (3)
O1 ⁱ —Cd1—N1—C15	136.6 (2)	C12—C13—C18—C17	17.2 (4)
O1—Cd1—N1—C15	65.9 (2)	C14—C13—C18—C17	-164.4 (3)
O2—Cd1—N1—C15	14.4 (2)	C12—C13—C18—C19	-162.9 (3)
Cd1—O2—C1—O1	-0.3 (2)	C14—C13—C18—C19	15.6 (4)
Cd1—O2—C1—C2	178.9 (2)	C17—C18—C19—C20	0.0 (5)
Cd1 ⁱ —O1—C1—O2	159.6 (3)	C13—C18—C19—C20	180.0 (3)
Cd1—O1—C1—O2	0.3 (3)	C16—N2—C20—C19	-1.4 (5)
Cd1 ⁱ —O1—C1—C2	-19.6 (5)	Cd1 ⁱⁱⁱ —N2—C20—C19	176.2 (3)
Cd1—O1—C1—C2	-178.9 (2)	C18—C19—C20—N2	0.9 (5)
O2—C1—C2—C3	33.5 (4)	Cd1—O6—C21—N3	156.6 (3)
O1—C1—C2—C3	-147.4 (3)	C22—N3—C21—O6	1.3 (6)
C4—O3—C3—C2	174.9 (3)	C23—N3—C21—O6	179.9 (4)
C1—C2—C3—O3	63.2 (3)		


Symmetry codes: (i) $-x, -y+2, -z+1$; (ii) $x, y+1, z$; (iii) $x, y-1, z$.

Hydrogen-bond geometry (\AA , °)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
O5—H5 ^{iv} —O4 ^{iv}	0.84	1.80	2.629 (3)	171
O1W—H11 ^v —O7	0.84	1.94	2.754 (4)	164
O1W—H12 ^v —O2 ^v	0.84	2.03	2.763 (3)	145

Symmetry codes: (iv) $-x+2, -y+1, -z$; (v) $-x+1, -y+2, -z+1$.

Fig. 1



supplementary materials

Fig. 2

