

{4-Bromo-2-[5-chloro-2-oxidophenyl]-iminomethylphenolato- $\kappa^3 O,N,O'$ -(methanol- κO)(methanolato- κO)oxido-vanadium(V)}

Gholam Hossein Shahverdizadeh,^a‡ Seik Weng Ng,^{b,c}
Edward R. T. Tiekkink^{b*} and Babak Mirtamizdoust^d

^aDepartment of Chemistry, Faculty of Science, Tabriz Branch, Islamic Azad University, PO Box 1655, Tabriz, Iran, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, ^cChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia, and ^dDepartment of Inorganic Chemistry, Faculty of Chemistry, University of Tabriz, PO Box 5166616471, Tabriz, Iran
Correspondence e-mail: edward.tiekkink@gmail.com


Received 24 January 2012; accepted 25 January 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.004 \text{ \AA}$; R factor = 0.046; wR factor = 0.131; data-to-parameter ratio = 15.6.

The title Schiff base complex, $[V(C_{13}H_7BrClNO_2)(CH_3O)O(CH_3OH)]$, features a vanadyl group, a tridentate Schiff base ligand, and coordinated methanol and methanolate ligands. The NO_5^- donor set is based on a distorted octahedron. Helical supramolecular chains along [010] are found in the crystal structure mediated by $O-H\cdots O$ hydrogen bonds formed between the coordinating methanol molecule and the phenolate O atom of the chlorobenzene residue.

Related literature

For the structures of (*E*)-2-(2-hydroxybenzylideneamino)-phenolates containing halide atoms on the aromatic ring(s), see: Yenişehirli *et al.* (2010). For related Schiff base vanadyl complexes containing alcohol and alkoxide ligands, see: Hartung *et al.* (2007); Clague *et al.* (1993). For the crystallization procedure, see: Harrowfield *et al.* (1996).



‡ Additional correspondence author, e-mail: shahverdizadeh@iaut.ac.ir.

Experimental

Crystal data

$[V(C_{13}H_7BrClNO_2)^{-}(CH_3O)O(CH_3O)]$
 $M_r = 454.57$
Monoclinic, $P2_1/n$
 $a = 9.9585 (2) \text{ \AA}$
 $b = 9.8949 (2) \text{ \AA}$
 $c = 17.3612 (3) \text{ \AA}$

$\beta = 100.746 (2)^\circ$
 $V = 1680.74 (6) \text{ \AA}^3$
 $Z = 4$
Cu $K\alpha$ radiation
 $\mu = 9.42 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
 $0.20 \times 0.20 \times 0.02 \text{ mm}$

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.255$, $T_{\max} = 0.834$

6974 measured reflections
3453 independent reflections
3125 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.131$
 $S = 1.04$
3453 reflections
221 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 1.37 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.93 \text{ e \AA}^{-3}$

Table 1
Selected bond lengths (Å).

V—O1	1.872 (2)	V—O4	1.766 (2)
V—O2	1.937 (2)	V—O5	1.596 (2)
V—O3	2.266 (2)	V—N1	2.170 (3)

Table 2
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H3 \cdots O2 ⁱ	0.84 (1)	1.89 (2)	2.702 (3)	163 (5)

Symmetry code: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We gratefully acknowledge support of this study by Tabriz Azad University, and thank the Ministry of Higher Education (Malaysia) for funding structural studies through the High-Impact Research scheme (UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5167).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.

metal-organic compounds

- Clague, M. J., Keder, N. L. & Butler, A. (1993). *Inorg. Chem.* **32**, 4754–4761.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Harrowfield, J. M., Miyamae, H., Skelton, B. W., Soudi, A. A. & White, A. H. (1996). *Aust. J. Chem.* **49**, 1165–1169.
Hartung, J., Ludwig, A., Svoboda, I. & Fuess, H. (2007). *Acta Cryst. E* **63**, m1422–m1423.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
Yenişehirli, G., Öztaş, N. A., Şahin, E., Çelebier, M., Ancin, N. & Öztaş, S. G. (2010). *Heteroat. Chem.* **21**, 373–385.

supplementary materials

Acta Cryst. (2012), E68, m221-m222 [doi:10.1107/S160053681200311X]

{4-Bromo-2-[{(5-chloro-2-oxidophenyl)iminomethyl]phenolato- κ^3O,N,O' }(methanol- κO)(methanolato- κO)oxidovanadium(V)}

G. H. Shahverdizadeh, S. W. Ng, E. R. T. Tiekkink and B. Mirtamizdoust

Comment

Structural studies of complexes with (*E*)-2-(2-hydroxybenzylideneamino)phenolates containing halide atoms on the aromatic ring(s) are comparatively rare (Yenişehirli *et al.*, 2010) and those of complexes containing the (*E*)-4-bromo-2-((5-chloro-2-hydroxyphenylimino)methyl)phenolate ligand have not been reported. Herein we report the oxo-vanadium(V) complex of this ligand, (I).

The V atom in (I) is coordinated by the *O,N,O*-tridentate Schiff base ligand, an oxido-O atom, an O atom of the methoxido ligand and an O atom of the methanol ligand. The resulting NO₅ donor set is based on an octahedron. The methanol ligand is *trans* to the oxido group and the V—O(methanol) bond length is significantly longer than the V—O(methanolate) bond, Table 1. The coordination geometry resembles those found in related V=O Schiff base compounds containing neutral and anionic forms of alcohols (Clague *et al.*, 1993; Hartung *et al.*, 2007).

The most prominent feature of the crystal packing is the formation of helical supramolecular chains along [010], Fig. 1 and Table 2. The connections between molecules are of the type O—H···O and involve the coordinated methanol molecule as the donor and the phenoxide-O atom of the chloro-substituted benzene ring as the acceptor.

Experimental

A solution of 4-chlorosalicylaldehyde (10 mmol) in EtOH (25 ml) was added drop-wise to a solution of 2-(aminomethyl)-4-bromophenol (10 mmol) in EtOH (15 ml). The mixture was refluxed for 5 h. The yellow precipitate was removed by filtration and recrystallized from MeOH solution. Then the ligand (0.8 mmol) was placed in one arm of a branched tube (Harrowfield *et al.*, 1996) and oxovanadium(IV) bis(acetylacetone) (0.8 mmol) placed in the other. Methanol was then added to fill both arms. The tube was sealed and the ligand-containing arm immersed in a bath at 333 K, while the other was left at ambient temperature. After two weeks, crystals were deposited in the arm held at ambient temperature. They were filtered off, washed with acetone and ether, and air-dried. Yield: 61%. *M.pt.*: 517 K.

Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H 0.95–0.98 Å, $U_{\text{iso}}(\text{H})$ 1.2–1.5 $U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation. The hydroxy H-atom was located in a difference Fourier map and was refined with a distance restraint of O—H 0.84±0.01 Å; U_{iso} was refined. The final difference Fourier map had a peak *ca* 1 Å from Br.

supplementary materials

Figures




Fig. 1. The molecular of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Fig. 2. A view of the helical supramolecular chain along [010] in (I). The O—H···O hydrogen bonds are shown as orange dashed lines.



Crystal data

[V(C ₁₃ H ₇ BrClNO ₂)(CH ₃ O)O(CH ₄ O)]	$F(000) = 904$
$M_r = 454.57$	$D_x = 1.796 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Cu $K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 3765 reflections
$a = 9.9585 (2) \text{ \AA}$	$\theta = 4.5\text{--}76.3^\circ$
$b = 9.8949 (2) \text{ \AA}$	$\mu = 9.42 \text{ mm}^{-1}$
$c = 17.3612 (3) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 100.746 (2)^\circ$	Plate, brown
$V = 1680.74 (6) \text{ \AA}^3$	$0.20 \times 0.20 \times 0.02 \text{ mm}$
$Z = 4$	

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector	3453 independent reflections
Radiation source: SuperNova (Cu) X-ray Source	3125 reflections with $I > 2\sigma(I)$
Mirror	$R_{\text{int}} = 0.029$
Detector resolution: 10.4041 pixels mm^{-1}	$\theta_{\text{max}} = 76.5^\circ, \theta_{\text{min}} = 4.8^\circ$
ω scan	$h = -12 \rightarrow 12$
Absorption correction: multi-scan (CrysAlis PRO; Agilent, 2010)	$k = -12 \rightarrow 12$
$T_{\text{min}} = 0.255, T_{\text{max}} = 0.834$	$l = -10 \rightarrow 21$
6974 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.046$	Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.131$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0869P)^2 + 1.3682P]$
3453 reflections	where $P = (F_o^2 + 2F_c^2)/3$
221 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
1 restraint	$\Delta\rho_{\text{max}} = 1.37 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.93 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.21340 (4)	0.40275 (4)	0.54299 (2)	0.03558 (16)
V	0.58605 (5)	1.02259 (5)	0.71783 (3)	0.01590 (16)
Cl	0.86189 (8)	1.03418 (8)	0.37755 (4)	0.02228 (19)
O1	0.4952 (2)	0.8625 (2)	0.73417 (12)	0.0202 (4)
O2	0.7315 (2)	1.1284 (2)	0.68762 (12)	0.0177 (4)
O3	0.7673 (2)	0.8869 (2)	0.76214 (13)	0.0194 (4)
H3	0.752 (5)	0.8057 (17)	0.771 (3)	0.041 (13)*
O4	0.6177 (2)	1.0892 (2)	0.81386 (13)	0.0211 (5)
O5	0.4594 (2)	1.1120 (2)	0.67670 (13)	0.0220 (5)
N1	0.6091 (3)	0.9280 (3)	0.60826 (15)	0.0169 (5)
C1	0.4304 (3)	0.7660 (3)	0.68900 (17)	0.0181 (6)
C2	0.3401 (3)	0.6813 (3)	0.72029 (19)	0.0216 (6)
H2	0.3230	0.6980	0.7715	0.026*
C3	0.2760 (3)	0.5739 (3)	0.6771 (2)	0.0234 (7)
H3A	0.2160	0.5164	0.6986	0.028*
C4	0.3008 (3)	0.5516 (3)	0.6017 (2)	0.0237 (7)
C5	0.3868 (3)	0.6329 (3)	0.56906 (19)	0.0218 (6)
H5	0.4016	0.6157	0.5175	0.026*
C6	0.4529 (3)	0.7419 (3)	0.61230 (18)	0.0191 (6)
C7	0.5457 (3)	0.8232 (3)	0.57644 (17)	0.0182 (6)
H7	0.5606	0.7972	0.5260	0.022*
C8	0.7007 (3)	1.0031 (3)	0.57077 (18)	0.0171 (6)
C9	0.7317 (3)	0.9777 (3)	0.49641 (17)	0.0172 (6)
H9	0.6910	0.9041	0.4654	0.021*
C10	0.8226 (3)	1.0625 (3)	0.46973 (17)	0.0189 (6)
C11	0.8838 (3)	1.1709 (3)	0.51455 (18)	0.0212 (6)
H11	0.9462	1.2277	0.4946	0.025*
C12	0.8540 (3)	1.1961 (3)	0.58772 (18)	0.0202 (6)
H12	0.8950	1.2701	0.6182	0.024*
C13	0.7623 (3)	1.1110 (3)	0.61637 (17)	0.0169 (6)
C14	0.8841 (3)	0.9337 (4)	0.8166 (2)	0.0259 (7)
H14A	0.9514	0.8607	0.8278	0.039*
H14B	0.8559	0.9619	0.8653	0.039*
H14C	0.9250	1.0107	0.7939	0.039*
C15	0.5543 (4)	1.2004 (4)	0.84499 (19)	0.0256 (7)
H15A	0.5968	1.2136	0.9001	0.038*
H15B	0.4567	1.1817	0.8412	0.038*

supplementary materials

H15C 0.5660 1.2822 0.8152 0.038*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br	0.0332 (2)	0.0270 (2)	0.0455 (3)	-0.01347 (15)	0.00452 (18)	-0.01039 (15)
V	0.0176 (3)	0.0136 (3)	0.0168 (3)	-0.00055 (19)	0.0040 (2)	-0.00131 (18)
Cl	0.0255 (4)	0.0234 (4)	0.0193 (4)	0.0001 (3)	0.0078 (3)	0.0003 (3)
O1	0.0231 (11)	0.0182 (10)	0.0201 (10)	-0.0041 (9)	0.0057 (8)	-0.0012 (8)
O2	0.0203 (10)	0.0146 (10)	0.0188 (10)	-0.0018 (8)	0.0051 (8)	-0.0001 (8)
O3	0.0208 (11)	0.0135 (10)	0.0228 (10)	-0.0017 (8)	0.0008 (8)	0.0025 (8)
O4	0.0231 (11)	0.0203 (11)	0.0209 (10)	-0.0001 (9)	0.0068 (8)	-0.0028 (8)
O5	0.0212 (11)	0.0203 (11)	0.0241 (11)	0.0011 (9)	0.0035 (9)	-0.0008 (8)
N1	0.0175 (12)	0.0151 (12)	0.0182 (11)	0.0005 (10)	0.0036 (9)	-0.0008 (9)
C1	0.0169 (13)	0.0140 (14)	0.0225 (14)	0.0021 (11)	0.0017 (11)	0.0013 (11)
C2	0.0185 (14)	0.0193 (15)	0.0276 (15)	-0.0004 (12)	0.0058 (12)	0.0011 (12)
C3	0.0174 (15)	0.0196 (15)	0.0334 (17)	-0.0030 (12)	0.0052 (12)	0.0026 (13)
C4	0.0184 (15)	0.0161 (14)	0.0338 (17)	-0.0032 (12)	-0.0025 (12)	-0.0014 (13)
C5	0.0213 (15)	0.0183 (15)	0.0244 (15)	0.0007 (12)	0.0006 (12)	-0.0023 (12)
C6	0.0179 (14)	0.0146 (14)	0.0242 (14)	-0.0001 (12)	0.0024 (11)	0.0018 (11)
C7	0.0198 (14)	0.0160 (14)	0.0190 (13)	0.0020 (12)	0.0039 (11)	0.0003 (11)
C8	0.0178 (14)	0.0132 (13)	0.0203 (14)	0.0002 (11)	0.0038 (11)	0.0005 (11)
C9	0.0203 (15)	0.0142 (14)	0.0171 (14)	0.0016 (11)	0.0031 (11)	-0.0001 (10)
C10	0.0197 (14)	0.0201 (15)	0.0167 (13)	0.0032 (12)	0.0031 (11)	-0.0003 (11)
C11	0.0220 (15)	0.0202 (15)	0.0219 (14)	-0.0014 (12)	0.0055 (11)	0.0040 (12)
C12	0.0205 (14)	0.0169 (14)	0.0224 (14)	-0.0014 (12)	0.0016 (11)	-0.0012 (11)
C13	0.0173 (14)	0.0148 (13)	0.0181 (13)	0.0023 (11)	0.0019 (11)	0.0012 (11)
C14	0.0204 (16)	0.0208 (15)	0.0322 (17)	-0.0014 (13)	-0.0063 (13)	0.0010 (13)
C15	0.0266 (16)	0.0259 (16)	0.0260 (15)	0.0014 (14)	0.0092 (13)	-0.0092 (13)

Geometric parameters (\AA , $^\circ$)

Br—C4	1.906 (3)	C4—C5	1.372 (5)
V—O1	1.872 (2)	C5—C6	1.405 (4)
V—O2	1.937 (2)	C5—H5	0.9500
V—O3	2.266 (2)	C6—C7	1.451 (4)
V—O4	1.766 (2)	C7—H7	0.9500
V—O5	1.596 (2)	C8—C13	1.400 (4)
V—N1	2.170 (3)	C8—C9	1.405 (4)
Cl—C10	1.740 (3)	C9—C10	1.377 (4)
O1—C1	1.324 (4)	C9—H9	0.9500
O2—C13	1.340 (4)	C10—C11	1.397 (4)
O3—C14	1.433 (4)	C11—C12	1.380 (4)
O3—H3	0.835 (10)	C11—H11	0.9500
O4—C15	1.424 (4)	C12—C13	1.400 (4)
N1—C7	1.284 (4)	C12—H12	0.9500
N1—C8	1.425 (4)	C14—H14A	0.9800
C1—C6	1.411 (4)	C14—H14B	0.9800
C1—C2	1.411 (4)	C14—H14C	0.9800

C2—C3	1.385 (5)	C15—H15A	0.9800
C2—H2	0.9500	C15—H15B	0.9800
C3—C4	1.396 (5)	C15—H15C	0.9800
C3—H3A	0.9500		
O5—V—O4	101.68 (11)	C6—C5—H5	120.1
O5—V—O1	99.96 (11)	C5—C6—C1	119.6 (3)
O4—V—O1	100.33 (10)	C5—C6—C7	118.0 (3)
O5—V—O2	98.49 (11)	C1—C6—C7	122.5 (3)
O4—V—O2	92.46 (10)	N1—C7—C6	124.4 (3)
O1—V—O2	154.86 (10)	N1—C7—H7	117.8
O5—V—N1	92.90 (11)	C6—C7—H7	117.8
O4—V—N1	163.59 (11)	C13—C8—C9	120.5 (3)
O1—V—N1	84.30 (10)	C13—C8—N1	112.9 (3)
O2—V—N1	77.84 (9)	C9—C8—N1	126.5 (3)
O5—V—O3	173.36 (10)	C10—C9—C8	118.1 (3)
O4—V—O3	84.85 (10)	C10—C9—H9	120.9
O1—V—O3	79.86 (9)	C8—C9—H9	120.9
O2—V—O3	79.85 (9)	C9—C10—C11	121.8 (3)
N1—V—O3	80.47 (9)	C9—C10—Cl	119.1 (2)
C1—O1—V	135.74 (19)	C11—C10—Cl	119.1 (2)
C13—O2—V	119.48 (19)	C12—C11—C10	120.3 (3)
C14—O3—V	122.00 (19)	C12—C11—H11	119.8
C14—O3—H3	110 (3)	C10—C11—H11	119.8
V—O3—H3	118 (4)	C11—C12—C13	119.0 (3)
C15—O4—V	129.3 (2)	C11—C12—H12	120.5
C7—N1—C8	122.0 (3)	C13—C12—H12	120.5
C7—N1—V	127.0 (2)	O2—C13—C12	121.8 (3)
C8—N1—V	110.89 (19)	O2—C13—C8	117.9 (3)
O1—C1—C6	122.4 (3)	C12—C13—C8	120.3 (3)
O1—C1—C2	118.4 (3)	O3—C14—H14A	109.5
C6—C1—C2	119.2 (3)	O3—C14—H14B	109.5
C3—C2—C1	120.7 (3)	H14A—C14—H14B	109.5
C3—C2—H2	119.7	O3—C14—H14C	109.5
C1—C2—H2	119.7	H14A—C14—H14C	109.5
C2—C3—C4	119.0 (3)	H14B—C14—H14C	109.5
C2—C3—H3A	120.5	O4—C15—H15A	109.5
C4—C3—H3A	120.5	O4—C15—H15B	109.5
C5—C4—C3	121.8 (3)	H15A—C15—H15B	109.5
C5—C4—Br	119.4 (3)	O4—C15—H15C	109.5
C3—C4—Br	118.8 (3)	H15A—C15—H15C	109.5
C4—C5—C6	119.7 (3)	H15B—C15—H15C	109.5
C4—C5—H5	120.1		
O5—V—O1—C1	69.0 (3)	C2—C3—C4—C5	0.0 (5)
O4—V—O1—C1	172.9 (3)	C2—C3—C4—Br	179.5 (2)
O2—V—O1—C1	−67.7 (4)	C3—C4—C5—C6	0.2 (5)
N1—V—O1—C1	−23.0 (3)	Br—C4—C5—C6	−179.2 (2)
O3—V—O1—C1	−104.3 (3)	C4—C5—C6—C1	0.3 (5)
O5—V—O2—C13	−82.2 (2)	C4—C5—C6—C7	178.4 (3)

supplementary materials


O4—V—O2—C13	175.6 (2)	O1—C1—C6—C5	176.3 (3)
O1—V—O2—C13	54.6 (3)	C2—C1—C6—C5	-1.0 (4)
N1—V—O2—C13	8.9 (2)	O1—C1—C6—C7	-1.7 (5)
O3—V—O2—C13	91.2 (2)	C2—C1—C6—C7	-179.1 (3)
O4—V—O3—C14	-36.8 (2)	C8—N1—C7—C6	179.1 (3)
O1—V—O3—C14	-138.3 (2)	V—N1—C7—C6	-5.0 (4)
O2—V—O3—C14	56.6 (2)	C5—C6—C7—N1	177.6 (3)
N1—V—O3—C14	135.9 (2)	C1—C6—C7—N1	-4.3 (5)
O5—V—O4—C15	-4.4 (3)	C7—N1—C8—C13	-178.1 (3)
O1—V—O4—C15	-106.9 (3)	V—N1—C8—C13	5.4 (3)
O2—V—O4—C15	94.8 (3)	C7—N1—C8—C9	1.7 (5)
N1—V—O4—C15	147.9 (4)	V—N1—C8—C9	-174.8 (3)
O3—V—O4—C15	174.4 (3)	C13—C8—C9—C10	-0.7 (5)
O5—V—N1—C7	-85.7 (3)	N1—C8—C9—C10	179.6 (3)
O4—V—N1—C7	121.4 (4)	C8—C9—C10—C11	0.2 (5)
O1—V—N1—C7	14.0 (3)	C8—C9—C10—Cl	-179.5 (2)
O2—V—N1—C7	176.2 (3)	C9—C10—C11—C12	0.0 (5)
O3—V—N1—C7	94.6 (3)	Cl—C10—C11—C12	179.7 (2)
O5—V—N1—C8	90.5 (2)	C10—C11—C12—C13	0.4 (5)
O4—V—N1—C8	-62.3 (4)	V—O2—C13—C12	171.7 (2)
O1—V—N1—C8	-169.7 (2)	V—O2—C13—C8	-8.8 (4)
O2—V—N1—C8	-7.54 (19)	C11—C12—C13—O2	178.6 (3)
O3—V—N1—C8	-89.1 (2)	C11—C12—C13—C8	-0.9 (5)
V—O1—C1—C6	21.3 (5)	C9—C8—C13—O2	-178.4 (3)
V—O1—C1—C2	-161.3 (2)	N1—C8—C13—O2	1.4 (4)
O1—C1—C2—C3	-176.2 (3)	C9—C8—C13—C12	1.0 (5)
C6—C1—C2—C3	1.3 (5)	N1—C8—C13—C12	-179.2 (3)
C1—C2—C3—C4	-0.8 (5)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O3—H3 ⁱ —O2 ⁱ	0.84 (1)	1.89 (2)	2.702 (3)	163 (5)

Symmetry codes: (i) $-x+3/2, y-1/2, -z+3/2$.

Fig. 1



supplementary materials

Fig. 2

