

**(Diaminomethylidene)sulfonium chloride–thiourea (3/2)**

Hafid Zouihri

Laboratoire Privé de Cristallographie (L.P.C), Kénitra, Morocco.  
Correspondence e-mail: hafid.zouihri@gmail.com


Received 7 December 2011; accepted 7 December 2011

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{N}-\text{C}) = 0.002$  Å;  $R$  factor = 0.026;  $wR$  factor = 0.068; data-to-parameter ratio = 16.3.

The asymmetric unit of the title salt,  $3\text{CH}_5\text{N}_2\text{S}^+\cdot 3\text{Cl}^-\cdot 2\text{CH}_4\text{N}_2\text{S}$ , contains two molecules of thiourea, three (diaminomethylidene)sulfonium cations and three chloride anions. The crystal packing is stabilized by  $\text{N}-\text{H}\cdots\text{Cl}$ ,  $\text{N}-\text{H}\cdots\text{S}$ ,  $\text{S}-\text{H}\cdots\text{Cl}$  and  $\text{S}-\text{H}\cdots\text{S}$  hydrogen bonds, forming a three-dimensional network.

**Related literature**

For applications of thiourea salts, see: Xing *et al.* (1987); Velsko *et al.* (1990).

**Experimental***Crystal data*

$3\text{CH}_5\text{N}_2\text{S}^+\cdot 3\text{Cl}^-\cdot 2\text{CH}_4\text{N}_2\text{S}$   
 $M_r = 489.98$   
 Monoclinic,  $P2_1/c$   
 $a = 16.3469$  (6) Å  
 $b = 8.9579$  (3) Å  
 $c = 16.1505$  (5) Å  
 $\beta = 109.105$  (2)°

$V = 2234.72$  (13) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.89$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.45 \times 0.32 \times 0.29$  mm

*Data collection*

Bruker APEXII CCD detector  
 diffractometer  
 25994 measured reflections

4885 independent reflections  
 4190 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.036$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.068$   
 $S = 1.06$   
 4885 reflections  
 300 parameters

15 restraints  
 All H-atom parameters refined  
 $\Delta\rho_{\text{max}} = 0.45$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.39$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{S3}$	0.82 (2)	2.76 (2)	3.5403 (17)	161.9 (18)
$\text{N1}-\text{H2N}\cdots\text{Cl1}$	0.848 (18)	2.474 (18)	3.2683 (16)	156.2 (18)
$\text{N10}-\text{H17N}\cdots\text{Cl3}^{\text{i}}$	0.86 (2)	2.45 (2)	3.2261 (17)	152 (2)
$\text{N10}-\text{H18N}\cdots\text{Cl1}^{\text{ii}}$	0.85 (2)	2.35 (2)	3.2019 (14)	175 (2)
$\text{N2}-\text{H3N}\cdots\text{Cl1}$	0.815 (18)	2.624 (18)	3.3656 (16)	152.0 (17)
$\text{N2}-\text{H4N}\cdots\text{Cl1}^{\text{ii}}$	0.86 (2)	2.45 (2)	3.3060 (17)	174.5 (17)
$\text{N3}-\text{H5N}\cdots\text{Cl2}$	0.86 (2)	2.51 (2)	3.3333 (16)	161 (2)
$\text{N3}-\text{H6N}\cdots\text{Cl1}^{\text{iii}}$	0.86 (2)	2.47 (2)	3.2799 (16)	157 (2)
$\text{N4}-\text{H7N}\cdots\text{Cl3}$	0.84 (2)	2.43 (2)	3.2422 (17)	163 (2)
$\text{N4}-\text{H8N}\cdots\text{Cl1}^{\text{iii}}$	0.84 (2)	2.57 (2)	3.3327 (16)	151 (2)
$\text{N4}-\text{H8N}\cdots\text{S1}^{\text{iv}}$	0.84 (2)	2.83 (2)	3.3571 (16)	123 (2)
$\text{N5}-\text{H9N}\cdots\text{S1}$	0.85 (2)	2.62 (2)	3.4493 (18)	166 (2)
$\text{N5}-\text{H10N}\cdots\text{Cl2}^{\text{v}}$	0.91 (2)	2.35 (2)	3.2262 (16)	161 (2)
$\text{N6}-\text{H11N}\cdots\text{Cl2}^{\text{vi}}$	0.88 (2)	2.47 (2)	3.3555 (17)	176 (1)
$\text{N6}-\text{H12N}\cdots\text{Cl2}^{\text{v}}$	0.88 (2)	2.65 (2)	3.4328 (15)	149 (2)
$\text{N7}-\text{H13N}\cdots\text{Cl2}$	0.86 (2)	2.37 (2)	3.2219 (15)	177 (1)
$\text{N7}-\text{H14N}\cdots\text{Cl3}^{\text{vii}}$	0.85 (2)	2.42 (2)	3.2066 (17)	154 (2)
$\text{N8}-\text{H15N}\cdots\text{S3}^{\text{viii}}$	0.85 (2)	2.44 (2)	3.2649 (16)	164 (2)
$\text{N8}-\text{H16N}\cdots\text{Cl3}^{\text{vii}}$	0.87 (2)	2.39 (2)	3.2036 (17)	156 (2)
$\text{N9}-\text{H19N}\cdots\text{Cl3}^{\text{i}}$	0.84 (2)	2.38 (2)	3.1674 (17)	157 (2)
$\text{N9}-\text{H20N}\cdots\text{S1}^{\text{vii}}$	0.872 (19)	2.38 (2)	3.2412 (16)	167.7 (19)
$\text{S2}-\text{H2S}\cdots\text{Cl2}$	1.253 (19)	2.315 (19)	3.5612 (6)	172.9 (14)
$\text{S4}-\text{H4S}\cdots\text{S3}^{\text{viii}}$	1.24 (2)	2.69 (2)	3.8755 (8)	159.2 (15)
$\text{S5}-\text{H5S}\cdots\text{S1}^{\text{vii}}$	1.29 (2)	2.64 (2)	3.8691 (7)	159.3 (16)

Symmetry codes: (i)  $-x+1, y-\frac{1}{2}, -z+\frac{1}{2}$ ; (ii)  $-x+1, y+\frac{1}{2}, -z+\frac{1}{2}$ ; (iii)  $x, -y+\frac{1}{2}, z+\frac{1}{2}$ ; (iv)  $x, -y+\frac{3}{2}, z+\frac{1}{2}$ ; (v)  $x, y+1, z$ ; (vi)  $-x, y+\frac{1}{2}, -z+\frac{1}{2}$ ; (vii)  $x, -y+\frac{3}{2}, z-\frac{1}{2}$ ; (viii)  $-x, -y+1, -z$ .

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009); software used to prepare material for publication: publCIF (Westrip, 2010).

The author thanks the Unit of Support for Technical and Scientific Research (UATRS, CNRST) for the X-ray measurements.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5746).

**References**

- Bruker (2005). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.  
 Velsko, S. (1990). Laser Program Annual Report, Lawrence Livermore National Laboratory, Livermore, Canada.  
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.  
 Xing, G., Jiang, M., Shao, Z. & Xu, D. (1987). *Chin. J. Lasers*, **14**, 357–360.

**supplementary materials**

*Acta Cryst.* (2012). E68, o257 [ doi:10.1107/S1600536811052809 ]

## (Diaminomethylidene)sulfonium chloride-thiourea (3/2)

H. Zouihri

### Comment

Various semi-organic NLO materials of thiourea were found to have higher mechanical strength, chemical stability, large nonlinearity, high resistance to laser induced damage, low angular sensitivity and good mechanical hardness [Xing, *et al.* 1987 and Velsko, *et al.* 1990]. Herein we present the crystal structure of the title compound (I).

The asymmetric unit of the title salt compound contains two molecules of thiourea (A and B), three cations of (diaminomethylidene)sulfonium (C, D and E) and three chloride anions.

The dihedral angles between the five molecules of the asymmetric unit are: A/B = 11,96 (9)°, A/C = 75,67 (9)°, A/D = 84,56 (9)°, A/E = 85,02 (9)°, B/C = 71,4 (3)°, B/D = 88,34 (9)° and B/E = 73,21 (9)° (Fig. 1).

In the crystal, the components are linked by a combination of thirteen N—H···Cl, five N—H···S, one S—H···Cl and two S—H···S hydrogen bonds into a three-dimensional structure. (Fig. 2 and Table. 1).

### Experimental

The title compound was synthesized at ambient temperature by a mixture of 5 mmol of thiourea and 5 mmol of HCl in ethanolic solution. The solution was slowly evaporated until solvent completely dried and white crystalline salt was obtained.

Suitable transparent crystals for X-ray measurement were grown from the final product in aqueous solution by slow evaporation method.

### Refinement

All H atoms were located from difference Fourier maps and refined isotropically, with restrained distance N—H = 0.88 (0.02) Å.

The highest residual density was found 0.82 Å from S4 and the deepest hole 0.74 Å from S4.

### Figures




Fig. 1. Molecular view of the title compound showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small spheres of arbitrary radii.





Fig. 2. Projection of the title compound along the *c* axis, H-bonds are represented by dashed lines.

## (Diaminomethylidene)sulfonium chloride–thiourea (3/2)

### Crystal data



$$M_r = 489.98$$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$$a = 16.3469 (6) \text{ \AA}$$

$$b = 8.9579 (3) \text{ \AA}$$

$$c = 16.1505 (5) \text{ \AA}$$

$$\beta = 109.105 (2)^\circ$$

$$V = 2234.72 (13) \text{ \AA}^3$$

$$Z = 4$$

$$F(000) = 1016$$

$$D_x = 1.456 \text{ Mg m}^{-3}$$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 257 reflections

$$\theta = 1.9\text{--}26.7^\circ$$

$$\mu = 0.89 \text{ mm}^{-1}$$

$$T = 100 \text{ K}$$

Prism, colourless

$$0.45 \times 0.32 \times 0.29 \text{ mm}$$

### Data collection

Bruker APEXII CCD detector  
diffractometer

Radiation source: fine-focus sealed tube

graphite

$\omega$  and  $\varphi$  scans

25994 measured reflections

4885 independent reflections

4190 reflections with  $I > 2\sigma(I)$

$$R_{\text{int}} = 0.036$$

$$\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 1.3^\circ$$

$$h = -19 \rightarrow 20$$

$$k = -11 \rightarrow 11$$

$$l = -20 \rightarrow 20$$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.026$$

$$wR(F^2) = 0.068$$

$$S = 1.06$$

4885 reflections

300 parameters

15 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

All H-atom parameters refined

$$w = 1/[\sigma^2(F_o^2) + (0.030P)^2 + 0.6051P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\text{max}} = 0.001$$

$$\Delta\rho_{\text{max}} = 0.45 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\text{min}} = -0.39 \text{ e \AA}^{-3}$$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.38413 (9)	0.58264 (16)	0.27012 (9)	0.0213 (3)
C2	0.25156 (10)	0.52767 (18)	0.50586 (9)	0.0248 (3)
C3	0.11263 (10)	0.86257 (17)	0.22690 (10)	0.0243 (3)
C4	0.07611 (10)	0.32825 (17)	0.03972 (10)	0.0256 (3)
C5	0.56391 (11)	0.66084 (17)	0.01498 (10)	0.0274 (3)
C11	0.36083 (2)	0.16969 (4)	0.21733 (2)	0.02368 (9)
C12	0.14816 (2)	0.26513 (4)	0.30590 (2)	0.02530 (9)
C13	0.25954 (3)	0.95922 (5)	0.52321 (3)	0.03198 (10)
H10N	0.1886 (13)	1.0337 (17)	0.2623 (13)	0.049 (6)*
H11N	-0.0063 (10)	0.887 (2)	0.2194 (12)	0.039 (5)*
H12N	0.0483 (13)	1.0283 (18)	0.2478 (13)	0.047 (6)*
H13N	0.1429 (12)	0.326 (2)	0.1616 (10)	0.038 (5)*
H14N	0.1828 (11)	0.408 (2)	0.1057 (12)	0.039 (5)*
H15N	0.0337 (12)	0.357 (2)	-0.0832 (11)	0.045 (6)*
H16N	0.1202 (11)	0.429 (2)	-0.0372 (12)	0.039 (5)*
H17N	0.6669 (11)	0.561 (2)	0.0740 (13)	0.044 (6)*
H18N	0.6263 (13)	0.628 (2)	0.1350 (11)	0.045 (6)*
H19N	0.6042 (11)	0.582 (2)	-0.0693 (12)	0.042 (6)*
H1N	0.2689 (14)	0.552 (2)	0.2609 (13)	0.048 (6)*
H20N	0.5209 (13)	0.667 (2)	-0.1093 (12)	0.035 (5)*
H2N	0.3116 (12)	0.413 (2)	0.2562 (12)	0.034 (5)*
H2S	0.1772 (14)	0.501 (2)	0.3670 (14)	0.063 (7)*
H3N	0.4498 (11)	0.418 (2)	0.2606 (11)	0.028 (5)*
H4N	0.5001 (12)	0.556 (2)	0.2727 (12)	0.033 (5)*
H4S	-0.0549 (15)	0.234 (2)	-0.0260 (15)	0.065 (7)*
H5N	0.2310 (11)	0.331 (2)	0.4620 (10)	0.033 (5)*
H5S	0.4337 (15)	0.768 (2)	-0.0474 (15)	0.068 (7)*
H6N	0.2823 (13)	0.341 (2)	0.5582 (11)	0.052 (6)*
H7N	0.2821 (12)	0.7027 (17)	0.5739 (12)	0.036 (5)*
H8N	0.3128 (12)	0.567 (2)	0.6237 (10)	0.042 (6)*
H9N	0.2319 (10)	0.890 (2)	0.2464 (12)	0.038 (5)*
N1	0.31308 (10)	0.50667 (17)	0.26422 (10)	0.0286 (3)
N10	0.62646 (10)	0.61272 (18)	0.08312 (9)	0.0349 (3)

## supplementary materials

---

N2	0.45314 (10)	0.50807 (17)	0.26805 (10)	0.0295 (3)
N3	0.25464 (10)	0.38179 (17)	0.50862 (10)	0.0341 (3)
N4	0.28862 (11)	0.60913 (18)	0.57553 (10)	0.0362 (4)
N5	0.18693 (10)	0.93523 (17)	0.24771 (11)	0.0355 (3)
N6	0.04351 (10)	0.93486 (16)	0.23118 (10)	0.0320 (3)
N7	0.14216 (10)	0.35255 (17)	0.11049 (9)	0.0307 (3)
N8	0.07441 (10)	0.37960 (17)	-0.03649 (9)	0.0307 (3)
N9	0.56308 (10)	0.63381 (18)	-0.06456 (9)	0.0317 (3)
S1	0.38657 (3)	0.77282 (4)	0.28160 (3)	0.02625 (10)
S2	0.19881 (3)	0.62081 (5)	0.40844 (3)	0.03084 (11)
S3	0.10697 (3)	0.67952 (4)	0.19506 (3)	0.02933 (10)
S4	-0.00958 (3)	0.22533 (6)	0.05196 (4)	0.04788 (14)
S5	0.48080 (3)	0.76310 (6)	0.03389 (3)	0.04573 (13)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0220 (8)	0.0232 (7)	0.0164 (7)	0.0000 (6)	0.0031 (6)	-0.0008 (5)
C2	0.0248 (8)	0.0286 (8)	0.0211 (8)	0.0015 (6)	0.0077 (6)	0.0016 (6)
C3	0.0261 (8)	0.0229 (7)	0.0215 (7)	0.0010 (6)	0.0045 (6)	0.0007 (6)
C4	0.0254 (8)	0.0247 (8)	0.0275 (8)	-0.0002 (6)	0.0099 (7)	-0.0041 (6)
C5	0.0271 (9)	0.0276 (8)	0.0276 (8)	-0.0040 (7)	0.0093 (7)	-0.0006 (6)
Cl1	0.0267 (2)	0.02205 (18)	0.02225 (18)	-0.00058 (14)	0.00803 (15)	-0.00263 (13)
Cl2	0.0296 (2)	0.02340 (18)	0.02352 (19)	-0.00071 (15)	0.00952 (16)	-0.00253 (14)
Cl3	0.0266 (2)	0.0341 (2)	0.0364 (2)	0.00110 (16)	0.01196 (17)	-0.00136 (17)
N1	0.0235 (8)	0.0219 (7)	0.0418 (8)	-0.0027 (6)	0.0126 (6)	-0.0068 (6)
N10	0.0344 (9)	0.0496 (9)	0.0197 (7)	0.0044 (7)	0.0076 (6)	0.0011 (6)
N2	0.0250 (8)	0.0211 (7)	0.0433 (9)	-0.0008 (6)	0.0122 (6)	-0.0021 (6)
N3	0.0438 (10)	0.0277 (8)	0.0246 (8)	0.0012 (7)	0.0027 (7)	0.0029 (6)
N4	0.0482 (10)	0.0320 (8)	0.0220 (7)	-0.0004 (7)	0.0027 (7)	-0.0004 (6)
N5	0.0243 (8)	0.0257 (7)	0.0560 (10)	-0.0006 (6)	0.0124 (7)	-0.0089 (7)
N6	0.0255 (8)	0.0237 (7)	0.0463 (9)	0.0012 (6)	0.0113 (7)	-0.0020 (6)
N7	0.0317 (8)	0.0394 (8)	0.0207 (7)	-0.0071 (7)	0.0082 (6)	-0.0012 (6)
N8	0.0270 (8)	0.0418 (8)	0.0218 (7)	-0.0056 (7)	0.0060 (6)	-0.0018 (6)
N9	0.0287 (8)	0.0426 (9)	0.0212 (7)	0.0079 (7)	0.0047 (6)	0.0034 (6)
S1	0.0223 (2)	0.01955 (19)	0.0319 (2)	-0.00003 (15)	0.00212 (16)	-0.00217 (15)
S2	0.0373 (2)	0.0297 (2)	0.0220 (2)	0.00586 (17)	0.00489 (17)	0.00139 (16)
S3	0.0255 (2)	0.02236 (19)	0.0326 (2)	0.00159 (15)	-0.00071 (17)	-0.00544 (16)
S4	0.0410 (3)	0.0524 (3)	0.0473 (3)	-0.0191 (2)	0.0104 (2)	0.0089 (2)
S5	0.0439 (3)	0.0502 (3)	0.0446 (3)	0.0125 (2)	0.0165 (2)	-0.0067 (2)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C1—N1	1.322 (2)	N4—H8N	0.841 (15)
C1—N2	1.321 (2)	N5—H10N	0.911 (15)
C2—N4	1.311 (2)	N5—H9N	0.847 (14)
C2—N3	1.308 (2)	N6—H11N	0.884 (14)
C3—N6	1.324 (2)	N6—H12N	0.875 (15)
C3—N5	1.321 (2)	N7—H14N	0.851 (14)

C4—N7	1.308 (2)	N7—H13N	0.855 (14)
C4—N8	1.306 (2)	N8—H16N	0.873 (14)
C5—N10	1.306 (2)	N8—H15N	0.851 (15)
C5—N9	1.303 (2)	N9—H19N	0.840 (15)
N1—H1N	0.81 (2)	N9—H20N	0.87 (2)
N1—H2N	0.84 (2)	S1—C1	1.7127 (15)
N10—H18N	0.850 (15)	S2—H2S	1.25 (2)
N10—H17N	0.857 (15)	S2—C2	1.7398 (15)
N2—H4N	0.861 (19)	S3—C3	1.7120 (16)
N2—H3N	0.811 (19)	S4—H4S	1.24 (2)
N3—H6N	0.862 (15)	S4—C4	1.7414 (16)
N3—H5N	0.855 (14)	S5—H5S	1.29 (2)
N4—H7N	0.844 (15)	S5—C5	1.7464 (17)
C2—S2—H2S	92.4 (10)	C1—N2—H3N	119.1 (13)
C5—S5—H5S	94.6 (10)	C1—N2—H4N	119.4 (12)
C4—S4—H4S	94.9 (10)	H3N—N2—H4N	121.4 (18)
N2—C1—N1	118.37 (15)	C3—N5—H9N	119.5 (13)
N2—C1—S1	121.06 (12)	C3—N5—H10N	119.3 (13)
N1—C1—S1	120.56 (12)	H9N—N5—H10N	121.2 (18)
N8—C4—N7	121.45 (15)	C4—N7—H13N	123.0 (13)
N8—C4—S4	121.58 (13)	C4—N7—H14N	117.6 (13)
N7—C4—S4	116.97 (12)	H13N—N7—H14N	118.9 (18)
N9—C5—N10	121.48 (16)	C1—N1—H2N	120.1 (12)
N9—C5—S5	120.81 (13)	C1—N1—H1N	119.4 (15)
N10—C5—S5	117.71 (13)	H2N—N1—H1N	119.9 (19)
N5—C3—N6	118.38 (15)	C5—N9—H20N	120.2 (12)
N5—C3—S3	120.21 (12)	C5—N9—H19N	116.2 (13)
N6—C3—S3	121.41 (13)	H20N—N9—H19N	123.6 (18)
N3—C2—N4	121.66 (15)	C3—N6—H12N	119.5 (14)
N3—C2—S2	120.80 (12)	C3—N6—H11N	119.5 (12)
N4—C2—S2	117.53 (13)	H12N—N6—H11N	120.9 (18)
C2—N3—H5N	119.8 (13)	C5—N10—H17N	117.8 (13)
C2—N3—H6N	117.5 (14)	C5—N10—H18N	121.4 (14)
H5N—N3—H6N	123 (2)	H17N—N10—H18N	120.7 (19)
C2—N4—H8N	119.2 (14)	C4—N8—H15N	121.5 (14)
C2—N4—H7N	120.5 (13)	C4—N8—H16N	115.9 (13)
H8N—N4—H7N	119.7 (19)	H15N—N8—H16N	122.2 (18)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N1—H1N $\cdots$ S3	0.82 (2)	2.76 (2)	3.5403 (17)	161.9 (18)
N1—H2N $\cdots$ C11	0.848 (18)	2.474 (18)	3.2683 (16)	156.2 (18)
N10—H17N $\cdots$ C13 <sup>i</sup>	0.86 (2)	2.45 (2)	3.2261 (17)	152.(2)
N10—H18N $\cdots$ C11 <sup>ii</sup>	0.85 (2)	2.35 (2)	3.2019 (14)	175 (2)
N2—H3N $\cdots$ C11	0.815 (18)	2.624 (18)	3.3656 (16)	152.0 (17)
N2—H4N $\cdots$ C11 <sup>ii</sup>	0.86 (2)	2.45 (2)	3.3060 (17)	174.5 (17)
N3—H5N $\cdots$ C12	0.86 (2)	2.51 (2)	3.3333 (16)	161.(2)

## supplementary materials

---

N3—H6N···C11 <sup>iii</sup>	0.86 (2)	2.47 (2)	3.2799 (16)	157.(2)
N4—H7N···C13	0.84 (2)	2.43 (2)	3.2422 (17)	163.(2)
N4—H8N···C11 <sup>iii</sup>	0.84 (2)	2.57 (2)	3.3327 (16)	151.(2)
N4—H8N···S1 <sup>iv</sup>	0.84 (2)	2.83 (2)	3.3571 (16)	123.(2)
N5—H9N···S1	0.85 (2)	2.62 (2)	3.4493 (18)	166.(2)
N5—H10N···C12 <sup>v</sup>	0.91 (2)	2.35 (2)	3.2262 (16)	161.(2)
N6—H11N···C12 <sup>vi</sup>	0.88 (2)	2.47 (2)	3.3555 (17)	176.(1)
N6—H12N···C12 <sup>v</sup>	0.88 (2)	2.65 (2)	3.4328 (15)	149.(2)
N7—H13N···C12	0.86 (2)	2.37 (2)	3.2219 (15)	177.(1)
N7—H14N···C13 <sup>vii</sup>	0.85 (2)	2.42 (2)	3.2066 (17)	154.(2)
N8—H15N···S3 <sup>viii</sup>	0.85 (2)	2.44 (2)	3.2649 (16)	164.(2)
N8—H16N···C13 <sup>vii</sup>	0.87 (2)	2.39 (2)	3.2036 (17)	156.(2)
N9—H19N···C13 <sup>i</sup>	0.84 (2)	2.38 (2)	3.1674 (17)	157.(2)
N9—H20N···S1 <sup>vii</sup>	0.872 (19)	2.38 (2)	3.2412 (16)	167.7 (19)
S2—H2S···C12	1.253 (19)	2.315 (19)	3.5612 (6)	172.9 (14)
S4—H4S···S3 <sup>viii</sup>	1.24 (2)	2.69 (2)	3.8755 (8)	159.2 (15)
S5—H5S···S1 <sup>vii</sup>	1.29 (2)	2.64 (2)	3.8691 (7)	159.3 (16)

Symmetry codes: (i)  $-x+1, y-1/2, -z+1/2$ ; (ii)  $-x+1, y+1/2, -z+1/2$ ; (iii)  $x, -y+1/2, z+1/2$ ; (iv)  $x, -y+3/2, z+1/2$ ; (v)  $x, y+1, z$ ; (vi)  $-x, y+1/2, -z+1/2$ ; (vii)  $x, -y+3/2, z-1/2$ ; (viii)  $-x, -y+1, -z$ .



Fig. 1




Fig. 2

