

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368


***N'*-[*(1E)*-(2,6-Difluorophenyl)methylidene]thiophene-2-carbohydrazide**Amer M. Alanazi,<sup>a</sup> Adnan A. Kadi,<sup>a</sup> Ali A. El-Emam<sup>a</sup> and Seik Weng Ng<sup>b,c,\*</sup><sup>a</sup>Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and <sup>c</sup>Chemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia  
Correspondence e-mail: seikweng@um.edu.my

Received 19 December 2011; accepted 29 December 2011

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å; disorder in main residue;  $R$  factor = 0.038;  $wR$  factor = 0.108; data-to-parameter ratio = 14.4.

In the title compound,  $\text{C}_{12}\text{H}_8\text{F}_2\text{N}_2\text{OS}$ , the thienyl ring is disordered over two positions, with the S atom of the major component [occupancy = 75.03 (18)%] oriented away from an *ortho*-F atom of the benzene ring. The molecule is nearly planar, the dihedral angle between the thiophene and benzene rings being 6.19 (18) (in the major component) or 3.5 (6)° (in the minor component). The azomethine  $\text{C}=\text{N}$  double-bond in the molecule is of an *E* configuration. In the crystal, molecules are linked by pairs of  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds, generating inversion dimers.

## Related literature

For a related structure, see: Alanazi *et al.* (2012).

## Experimental

## Crystal data

 $\text{C}_{12}\text{H}_8\text{F}_2\text{N}_2\text{OS}$  $M_r = 266.26$ Triclinic,  $P\bar{1}$   
 $a = 6.5032$  (3) Å  
 $b = 7.7516$  (4) Å  
 $c = 11.5224$  (5) Å  
 $\alpha = 95.184$  (4)°  
 $\beta = 103.344$  (4)°  
 $\gamma = 94.285$  (4)° $V = 560.11$  (5) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.30$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.40 \times 0.30 \times 0.20$  mm

## Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector  
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)  
 $T_{\min} = 0.888$ ,  $T_{\max} = 0.942$ 8250 measured reflections  
2594 independent reflections  
2174 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.035$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.038$   
 $wR(F^2) = 0.108$   
 $S = 1.07$   
2594 reflections  
180 parameters  
24 restraintsH atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.33$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.28$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^i$	0.88 (2)	1.97 (2)	2.856 (2)	174 (2)

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank the Deanship of Scientific Research and the Research Center of the College of Pharmacy, King Saud University, and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5416).

## References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.  
Alanazi, A. M., Lahsasni, S., El-Emam, A. A. & Ng, S. W. (2012). *Acta Cryst.* **E68**, o314.  
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

**supplementary materials**

*Acta Cryst.* (2012). E68, o315 [ doi:10.1107/S160053681105611X ]

## *N'*-[(1*E*)-(2,6-Difluorophenyl)methylidene]thiophene-2-carbohydrazide

A. M. Alanazi, A. A. Kadi, A. A. El-Emam and S. W. Ng

### Comment

2-Thienoylhydrazide forms a large number of Schiff base derivatives with substituted benzaldehydes; among those whose crystal structures have been reported are the 4-chloro and 4-bromo derivatives. The 4-fluoro analog is disordered in respect of the thienyl ring (Alanazi *et al.*, 2012). The azomethine double-bond in the approximately planar C<sub>12</sub>H<sub>8</sub>F<sub>2</sub>N<sub>2</sub>OS molecule (Scheme 1) is of an *E* configuration (Fig. 1). The thienyl ring is disordered over two positions, with the S atom of the major component (75.03 (18)%) oriented away from an *ortho*-F atom of the benzene ring. Two molecules are linked across a center-of-inversion by an N–H···O hydrogen bond to generate a dimer (Table 1).

### Experimental

2-Thienoylhydrazide (1.42 g, 0.01 mol) and 2,6-difluorobenzaldehyde (1.42 g, 0.01 mol) dissolved in ethanol (8 ml) was heated for 1 h. The product was collected and recrystallized from ethanol to yield the Schiff base in 90% yield, m.p. 453–45548 K.

### Refinement

Carbon-bound H-atoms were placed in calculated positions [C–H 0.95 Å,  $U_{\text{iso}}(\text{H}) 1.2U_{\text{eq}}(\text{C})$ ] and were included in the refinement in the riding model approximation.

The amino H-atom was located in a difference Fourier map, and was refined freely.

The thiophene ring is disordered over two positions in respect of four of the five atoms, with major component being 87.1 (2) %. Pairs of C–C and C–S bond distances were restrained to within 0.0 Å of each other. The temperature factors of C3' was set to those of S1 (as were these pairs: C2' to C1, C1' to C2 and S1' to C3). The anisotropic temperature factors of the disordered atoms were tightly restrained to be nearly isotropic.

### Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of C<sub>12</sub>H<sub>8</sub>F<sub>2</sub>N<sub>2</sub>OS at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. The disorder is not shown.

## *N*'-[(1*E*)-(2,6-Difluorophenyl)methylidene]thiophene- 2-carbohydrazide

### Crystal data

$C_{12}H_8F_2N_2OS$	$Z = 2$
$M_r = 266.26$	$F(000) = 272$
Triclinic, <i>PT</i>	$D_x = 1.579 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 6.5032 (3) \text{ \AA}$	Cell parameters from 3787 reflections
$b = 7.7516 (4) \text{ \AA}$	$\theta = 2.7\text{--}27.5^\circ$
$c = 11.5224 (5) \text{ \AA}$	$\mu = 0.30 \text{ mm}^{-1}$
$\alpha = 95.184 (4)^\circ$	$T = 100 \text{ K}$
$\beta = 103.344 (4)^\circ$	Wedge, colorless
$\gamma = 94.285 (4)^\circ$	$0.40 \times 0.30 \times 0.20 \text{ mm}$
$V = 560.11 (5) \text{ \AA}^3$	

### Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector	2594 independent reflections
Radiation source: SuperNova (Mo) X-ray Source	2174 reflections with $I > 2\sigma(I)$
Mirror	$R_{\text{int}} = 0.035$
Detector resolution: $10.4041 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 27.6^\circ$ , $\theta_{\text{min}} = 2.7^\circ$
$\omega$ scan	$h = -8 \rightarrow 8$
Absorption correction: multi-scan (CrysAlis PRO; Agilent, 2010)	$k = -10 \rightarrow 10$
$T_{\text{min}} = 0.888$ , $T_{\text{max}} = 0.942$	$l = -15 \rightarrow 14$
8250 measured reflections	

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.038$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.108$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.0585P)^2 + 0.1004P]$
2594 reflections	where $P = (F_o^2 + 2F_c^2)/3$
180 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
24 restraints	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.61235 (9)	0.40630 (10)	0.12814 (6)	0.01924 (19)	0.7503 (18)
S1'	0.9967 (5)	0.2911 (5)	0.2831 (4)	0.0226 (7)	0.2497 (18)
F1	1.02233 (15)	0.19692 (13)	0.85830 (8)	0.0285 (3)	
F2	1.26733 (15)	0.14995 (13)	0.50356 (8)	0.0270 (3)	
O1	0.47384 (17)	0.48037 (15)	0.34181 (10)	0.0228 (3)	
N1	0.7178 (2)	0.37086 (17)	0.47830 (12)	0.0188 (3)	
N2	0.8961 (2)	0.28597 (16)	0.51113 (11)	0.0178 (3)	
C1	0.8196 (10)	0.3496 (7)	0.0668 (6)	0.0214 (8)	0.7503 (18)
H1A	0.8126	0.3501	-0.0165	0.026*	0.7503 (18)
C2	0.9871 (9)	0.3058 (8)	0.1414 (3)	0.0199 (7)	0.7503 (18)
H2	1.1127	0.2725	0.1205	0.024*	0.7503 (18)
C3	0.9481 (7)	0.3168 (7)	0.2667 (5)	0.0226 (7)	0.7503 (18)
H3	1.0483	0.2915	0.3357	0.027*	0.7503 (18)
C1'	0.6702 (15)	0.4020 (15)	0.1526 (9)	0.01924 (19)	0.25
H1'	0.5350	0.4436	0.1277	0.023*	0.2497 (18)
C2'	0.801 (3)	0.371 (2)	0.074 (2)	0.0214 (8)	0.25
H2'	0.7827	0.3824	-0.0092	0.026*	0.2497 (18)
C3'	0.983 (3)	0.314 (3)	0.1659 (13)	0.0199 (7)	0.25
H3'	1.1048	0.2898	0.1375	0.024*	0.2497 (18)
C4	0.7509 (2)	0.36777 (19)	0.26883 (13)	0.0172 (3)	
C5	0.6407 (2)	0.40831 (19)	0.36486 (13)	0.0176 (3)	
C7	0.9443 (2)	0.26314 (19)	0.62276 (14)	0.0176 (3)	
H7	0.8547	0.3020	0.6721	0.021*	
C8	1.1326 (2)	0.17935 (19)	0.67688 (13)	0.0167 (3)	
C9	1.1710 (2)	0.1525 (2)	0.79783 (13)	0.0190 (3)	
C10	1.3490 (3)	0.0862 (2)	0.86064 (14)	0.0225 (4)	
H10	1.3680	0.0733	0.9435	0.027*	
C11	1.4994 (3)	0.0391 (2)	0.79913 (15)	0.0229 (4)	
H11	1.6238	-0.0069	0.8401	0.027*	
C12	1.4694 (2)	0.0587 (2)	0.67816 (15)	0.0217 (3)	
H12	1.5714	0.0246	0.6356	0.026*	
C13	1.2902 (2)	0.1279 (2)	0.62060 (14)	0.0192 (3)	
H1	0.656 (3)	0.409 (3)	0.535 (2)	0.040 (6)*	

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0198 (4)	0.0239 (3)	0.0153 (3)	0.0069 (3)	0.0043 (2)	0.0045 (2)
S1'	0.0216 (14)	0.0257 (11)	0.0218 (11)	0.0042 (9)	0.0074 (10)	0.0018 (8)
F1	0.0271 (5)	0.0443 (6)	0.0201 (5)	0.0167 (5)	0.0120 (4)	0.0075 (4)
F2	0.0266 (5)	0.0396 (6)	0.0208 (5)	0.0157 (4)	0.0119 (4)	0.0080 (4)
O1	0.0185 (6)	0.0310 (6)	0.0209 (6)	0.0117 (5)	0.0054 (5)	0.0045 (5)
N1	0.0168 (6)	0.0247 (7)	0.0175 (6)	0.0095 (5)	0.0071 (5)	0.0026 (5)
N2	0.0135 (6)	0.0196 (7)	0.0208 (7)	0.0053 (5)	0.0043 (5)	0.0026 (5)

## supplementary materials

---

C1	0.0245 (12)	0.0234 (14)	0.0196 (10)	0.0062 (10)	0.0103 (8)	0.0032 (10)
C2	0.0207 (8)	0.0267 (10)	0.0157 (18)	0.0046 (7)	0.0111 (13)	0.0002 (14)
C3	0.0216 (14)	0.0257 (11)	0.0218 (11)	0.0042 (9)	0.0074 (10)	0.0018 (8)
C1'	0.0198 (4)	0.0239 (3)	0.0153 (3)	0.0069 (3)	0.0043 (2)	0.0045 (2)
C2'	0.0245 (12)	0.0234 (14)	0.0196 (10)	0.0062 (10)	0.0103 (8)	0.0032 (10)
C3'	0.0207 (8)	0.0267 (10)	0.0157 (18)	0.0046 (7)	0.0111 (13)	0.0002 (14)
C4	0.0167 (7)	0.0182 (7)	0.0172 (7)	0.0026 (6)	0.0050 (6)	0.0018 (6)
C5	0.0168 (7)	0.0174 (7)	0.0192 (7)	0.0029 (6)	0.0055 (6)	0.0009 (6)
C7	0.0157 (7)	0.0175 (7)	0.0210 (7)	0.0035 (6)	0.0071 (6)	0.0014 (6)
C8	0.0148 (7)	0.0153 (7)	0.0202 (8)	0.0031 (6)	0.0041 (6)	0.0016 (6)
C9	0.0187 (7)	0.0210 (8)	0.0194 (8)	0.0056 (6)	0.0076 (6)	0.0015 (6)
C10	0.0239 (8)	0.0248 (8)	0.0184 (8)	0.0057 (7)	0.0024 (6)	0.0047 (6)
C11	0.0165 (7)	0.0216 (8)	0.0298 (9)	0.0063 (6)	0.0018 (7)	0.0053 (7)
C12	0.0172 (7)	0.0217 (8)	0.0286 (8)	0.0060 (6)	0.0089 (6)	0.0033 (6)
C13	0.0197 (8)	0.0186 (7)	0.0205 (8)	0.0027 (6)	0.0069 (6)	0.0032 (6)

### *Geometric parameters (Å, °)*

S1—C4	1.7269 (16)	C1'—C2'	1.40 (2)
S1—C1	1.727 (6)	C1'—H1'	0.9500
S1'—C3'	1.360 (16)	C2'—C3'	1.52 (3)
S1'—C4	1.725 (3)	C2'—H2'	0.9500
F1—C9	1.3624 (17)	C3'—H3'	0.9500
F2—C13	1.3500 (18)	C4—C5	1.474 (2)
O1—C5	1.2425 (18)	C7—C8	1.466 (2)
N1—C5	1.354 (2)	C7—H7	0.9500
N1—N2	1.3702 (18)	C8—C9	1.395 (2)
N1—H1	0.88 (2)	C8—C13	1.397 (2)
N2—C7	1.284 (2)	C9—C10	1.377 (2)
C1—C2	1.311 (8)	C10—C11	1.385 (2)
C1—H1A	0.9500	C10—H10	0.9500
C2—C3	1.519 (7)	C11—C12	1.386 (2)
C2—H2	0.9500	C11—H11	0.9500
C3—C4	1.374 (5)	C12—C13	1.374 (2)
C3—H3	0.9500	C12—H12	0.9500
C1'—C4	1.378 (10)		
C4—S1—C1	91.0 (2)	C5—C4—S1'	127.47 (18)
C3'—S1'—C4	88.3 (9)	C3—C4—S1	111.5 (3)
C5—N1—N2	123.04 (13)	C5—C4—S1	114.07 (11)
C5—N1—H1	118.9 (14)	O1—C5—N1	118.90 (13)
N2—N1—H1	117.9 (14)	O1—C5—C4	119.41 (14)
C7—N2—N1	113.74 (13)	N1—C5—C4	121.68 (14)
C2—C1—S1	116.3 (5)	N2—C7—C8	122.42 (14)
C2—C1—H1A	121.9	N2—C7—H7	118.8
S1—C1—H1A	121.9	C8—C7—H7	118.8
C1—C2—C3	109.0 (5)	C9—C8—C13	114.10 (14)
C1—C2—H2	125.5	C9—C8—C7	119.41 (14)
C3—C2—H2	125.5	C13—C8—C7	126.42 (14)
C4—C3—C2	112.2 (4)	F1—C9—C10	117.74 (14)

C4—C3—H3	123.9	F1—C9—C8	117.47 (14)
C2—C3—H3	123.9	C10—C9—C8	124.78 (14)
C4—C1'—C2'	115.3 (11)	C9—C10—C11	117.90 (15)
C4—C1'—H1'	122.3	C9—C10—H10	121.0
C2'—C1'—H1'	122.3	C11—C10—H10	121.0
C1'—C2'—C3'	96.1 (16)	C10—C11—C12	120.42 (15)
C1'—C2'—H2'	131.9	C10—C11—H11	119.8
C3'—C2'—H2'	131.9	C12—C11—H11	119.8
S1'—C3'—C2'	129.6 (16)	C13—C12—C11	119.10 (15)
S1'—C3'—H3'	115.2	C13—C12—H12	120.5
C2'—C3'—H3'	115.2	C11—C12—H12	120.5
C3—C4—C1'	103.4 (5)	F2—C13—C12	117.98 (14)
C3—C4—C5	134.2 (3)	F2—C13—C8	118.33 (14)
C1'—C4—C5	121.8 (4)	C12—C13—C8	123.67 (15)
C1'—C4—S1'	110.5 (5)		
C5—N1—N2—C7	179.73 (14)	C3—C4—C5—O1	167.6 (3)
C4—S1—C1—C2	-0.8 (5)	C1'—C4—C5—O1	-2.3 (6)
S1—C1—C2—C3	0.4 (7)	S1'—C4—C5—O1	172.3 (2)
C1—C2—C3—C4	0.4 (7)	S1—C4—C5—O1	-6.89 (19)
C4—C1'—C2'—C3'	-1.0 (16)	C3—C4—C5—N1	-11.6 (4)
C4—S1'—C3'—C2'	-4.0 (19)	C1'—C4—C5—N1	178.5 (5)
C1'—C2'—C3'—S1'	4(2)	S1'—C4—C5—N1	-6.9 (3)
C2—C3—C4—C1'	-4.4 (7)	S1—C4—C5—N1	173.89 (12)
C2—C3—C4—C5	-175.6 (3)	N1—N2—C7—C8	-178.28 (13)
C2—C3—C4—S1'	155 (3)	N2—C7—C8—C9	-177.48 (14)
C2—C3—C4—S1	-1.0 (5)	N2—C7—C8—C13	5.7 (3)
C2'—C1'—C4—C3	1.9 (12)	C13—C8—C9—F1	-178.95 (13)
C2'—C1'—C4—C5	174.5 (10)	C7—C8—C9—F1	3.9 (2)
C2'—C1'—C4—S1'	-0.9 (13)	C13—C8—C9—C10	1.8 (2)
C2'—C1'—C4—S1	-157 (4)	C7—C8—C9—C10	-175.39 (14)
C3'—S1'—C4—C3	-19 (3)	F1—C9—C10—C11	179.29 (14)
C3'—S1'—C4—C1'	2.6 (10)	C8—C9—C10—C11	-1.4 (2)
C3'—S1'—C4—C5	-172.6 (9)	C9—C10—C11—C12	0.0 (2)
C3'—S1'—C4—S1	6.6 (9)	C10—C11—C12—C13	1.0 (2)
C1—S1—C4—C3	1.0 (3)	C11—C12—C13—F2	178.28 (14)
C1—S1—C4—C1'	23 (3)	C11—C12—C13—C8	-0.6 (2)
C1—S1—C4—C5	176.8 (2)	C9—C8—C13—F2	-179.60 (13)
C1—S1—C4—S1'	-2.4 (3)	C7—C8—C13—F2	-2.7 (2)
N2—N1—C5—O1	177.55 (13)	C9—C8—C13—C12	-0.7 (2)
N2—N1—C5—C4	-3.2 (2)	C7—C8—C13—C12	176.23 (15)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 $\cdots$ O1 <sup>i</sup>	0.88 (2)	1.97 (2)	2.856 (2)	174 (2)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ .

Fig. 1

