

$b = 11.5337(8)$ Å
 $c = 11.8745(9)$ Å
 $\alpha = 63.230(7)^\circ$
 $\beta = 80.596(6)^\circ$
 $\gamma = 79.970(6)^\circ$
 $V = 1091.69(14)$ Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 170$ K
 $0.28 \times 0.25 \times 0.24$ mm

3,3-Dimethyl-1,2,3,4-tetrahydrocyclopenta[b]indole-1,2-dione (bruceolline E)

Jason A. Jordon,^a Jeanese C. Badenock,^a Gordon W. Gribble,^b Jerry P. Jasinski^{c*} and James A. Golen^c

^aDepartment of Biological and Chemical Sciences, University of the West Indies, Cave Hill, Barbados, ^bDepartment Chemistry, Dartmouth College, Hanover, NH 03755-3564, USA, and ^cDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA

Correspondence e-mail: jjasinski@keene.edu


Received 28 December 2011; accepted 5 January 2012

Key indicators: single-crystal X-ray study; $T = 170$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.053; wR factor = 0.157; data-to-parameter ratio = 18.9.

The title compound, $C_{13}H_{11}NO_2$, crystallizes with two molecules in the asymmetric unit. The crystal packing is stabilized by N—H···O hydrogen bonds, which link the molecules into chains along [101], and weak C—H···O interactions.

Related literature

For the first isolation of bruceolline E as yellow needles, see: Ouyang *et al.* (1994). For the first total synthesis of bruceolline E in three steps from the known ethyl indole-1-carboxylate, see: Jordan *et al.* (2011). For examples of similar tandem acylation/Nazarov cyclization with pyrroles, see: Song *et al.* (2006). For examples of Nazarov cyclizations with indoles, see: Bergman & Venemalm (1992); Cheng & Cheung (1996); Ishikura *et al.* (2000); Miki *et al.* (2001); Churruca *et al.* (2010). For examples of α -diketone oxidations using selenium dioxide, see: Gribble *et al.* (1988); Xu *et al.* (2002); Belsey *et al.* (2006). For related cyclopenta[b]indolone alkaloids and their analogues, see: Cheng *et al.* (1991); Garcia-Pichel & Castenholz (1991); Garcia-Pichel *et al.* (1992); Proteau *et al.* (1993); Ekebergh *et al.* (2011); Kobayashi *et al.* (1994); Jacquemard *et al.* (2004); Ploutno & Carmeli (2001). For standard bond lengths, see: Allen *et al.* (1987).



Experimental

Crystal data

$C_{13}H_{11}NO_2$
 $M_r = 213.23$

Triclinic, $P\bar{1}$
 $a = 9.1091(7)$ Å

Data collection

Oxford Diffraction Xcalibur Eos
 Gemini diffractometer
 Absorption correction: multi-scan
 (*CrysAlis RED*; Oxford
 Diffraction, 2010)
 $T_{\min} = 0.976$, $T_{\max} = 0.979$

10062 measured reflections
 5644 independent reflections
 4698 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.012$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.157$
 $S = 1.05$
 5644 reflections
 299 parameters
 2 restraints

H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\max} = 0.36$ e Å⁻³
 $\Delta\rho_{\min} = -0.20$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1NA···O2A	0.89 (1)	1.95 (1)	2.8025 (16)	161 (2)
N1A—H1NB···O2 ⁱ	0.92 (1)	1.87 (1)	2.7686 (16)	164 (2)
C9A—H9AA···O1 ⁱ	0.95	2.51	3.376 (2)	152
C12A—H12B···O2 ⁱ	0.98	2.57	3.430 (2)	146

Symmetry code: (i) $x - 1, y, z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

JCB wishes to thank the School of Graduate Studies and Research, UWI, and the Government of Barbados for the funding of this research. JPJ acknowledges the NSF–MRI program (grant No. CHE1039027) for funds to purchase the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: QM2048).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Belsey, S., Danks, T. N. & Wagner, G. (2006). *Synth. Commun.* **36**, 1019–1024.
- Bergman, J. & Venemalm, L. (1992). *Tetrahedron* **48**, 759–768.
- Cheng, K.-F., Chan, K.-P. & Lai, T.-F. (1991). *J. Chem. Soc. Perkin Trans. 1*, pp. 2461–2465.
- Cheng, K.-F. & Cheung, M.-K. (1996). *J. Chem. Soc., Perkin Trans. 1*, pp. 1213–1218.
- Churruca, F., Fousteris, M., Ishikawa, Y., von Wantoch Rekowski, M., Hounsou, C., Surrey, T. & Giannis, A. (2010). *Org. Lett.* **12**, 2096–2099.
- Ekebergh, A., Karlsson, I., Mete, R., Pan, Y., Börje, A. & Mårtensson, J. (2011). *Org. Lett.* **13**, 4458–4461.
- Garcia-Pichel, F. & Castenholz, R. W. (1991). *J. Phycol.* **27**, 395–409.
- Garcia-Pichel, F., Sherry, N. D. & Castenholz, R. W. (1992). *Photochem. Photobiol.* **56**, 17–23.

- Gribble, G. W., Barden, T. C. & Johnson, D. A. (1988). *Tetrahedron*, **44**, 3195–3202.
- Ishikura, M., Imaizumi, K. & Katagiri, N. (2000). *Heterocycles*, **53**, 2201–2220.
- Jacquemard, U., Bénéteau, V., Lefoix, M., Routier, S., Mérou, J.-Y. & Coudert, G. (2004). *Tetrahedron*, **60**, 10039–10047.
- Jordan, J. A., Gribble, G. W. & Badenock, J. C. (2011). *Tetrahedron Lett.* **52**, 6772–6774.
- Kobayashi, A., Kajiyama, S.-I., Inawaka, K., Kanzaki, H. & Kawazu, K. Z. (1994). *Z. Naturforsch. Teil C*, **49**, 464–470.
- Miki, Y., Hachiken, H., Kawazoe, A., Tsuzaki, Y. & Yanase, N. (2001). *Heterocycles*, **55**, 1291–1299.
- Ouyang, Y., Koike, K. & Ohmoto, T. (1994). *Phytochemistry*, **37**, 575–578.
- Oxford Diffraction (2010). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Ploutno, A. & Carmeli, S. (2001). *J. Nat. Prod.* **64**, 544–545.
- Proteau, P. J., Gerwick, W. H., Garcia-Pichel, F. & Castenholz, R. (1993). *Cell. Mol. Life Sci.* **49**, 825–829.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Song, C., Knight, D. W. & Whatton, M. A. (2006). *Org. Lett.* **8**, 163–166.
- Xu, P.-F., Chen, Y.-S., Lin, S.-I. & Lu, T.-J. (2002). *J. Org. Chem.* **67**, 2309–2314.

supplementary materials

Acta Cryst. (2012). E68, o364-o365 [doi:10.1107/S1600536812000517]

3,3-Dimethyl-1,2,3,4-tetrahydrocyclopenta[*b*]indole-1,2-dione (bruceolline E)

J. A. Jordon, J. C. Badenock, G. W. Gribble, J. P. Jasinski and J. A. Golen

Comment

The fused indole alkaloid bruceolline E was first isolated as yellow needles from the root wood *Brucea mollis* Wall. *Var. tonkinensis* Lecomte and reported by Ohmoto (Ouyang *et al.*, 1994). Our synthesis of this cyclopenta[*b*]indolone natural product included a tandem acylation/Nazarov cyclization sequence (Song *et al.*, 2006) and insertion of the α -diketone functionality using selenium dioxide (Gribble *et al.*, 1988) in 60% yield. Similar Nazarov cyclizations with indoles (Bergman & Venemalm, 1992; Cheng & Cheung, 1996; Ishikura *et al.*, 2000; Miki *et al.*, 2001; Churruga *et al.*, 2010) have been reported. Related α -diketone oxidations using selenium dioxide (Xu *et al.*, 2002; Belsey *et al.*, 2006) and cyclopenta[*b*]indolone natural product alkaloid analogues such as scytomin (Garcia-Pichel & Castenholz 1991; Garcia-Pichel *et al.* 1992; Proteau *et al.* 1993; Ekebergh *et al.* 2011), nostodione (Kobayashi *et al.*, 1994) and prenostodione (Ploutno & Carmeli, 2001) have also been reported. Our efforts have yielded the first total synthesis of bruceolline E (Jordan *et al.*, 2011), in three steps from the known ethyl indole-1-carboxylate.

We now report herein the first crystal structure of the title compound, $C_{13}H_{11}NO_2$, bruceolline E, which confirms the cyclopenta[*b*] indole moiety and the α -diketone functionalities earlier assigned by NMR methods.

In the crystal structure of the title compound, $C_{13}H_{11}NO_2$, two molecules crystallize in the asymmetric unit (Fig. 1). Bond lengths are in normal ranges (Allen *et al.*, 1987). Crystal packing is stabilized by N—H \cdots O hydrogen bonds (Table 1) and weak C—H \cdots O intermolecular interactions linking these 1-D chains along [011] (Fig. 2).

Experimental

A solution of 3,3-dimethyl-1,2-dioxo-2,3-dihydro-1H-cyclopenta[*b*] indole-4-carboxylic acid ethyl ester (0.58 g, 2.03 mmol, 1 eq.), anhydrous THF (30 mL) and TBAF (10.2 mL, 1.0 M in THF, 10.2 mmol, 5.02 eq.) was stirred under argon at reflux for 2 h (Jacquemard *et al.*, 2004). The mixture was allowed to cool before quenching with saturated NH_4Cl . After extraction with CH_2Cl_2 (3 x 40 mL), the organic layers were combined, dried over $MgSO_4$ and concentrated in vacuo to give a yellow solid. Flash column chromatography (100% EtOAc) gave the product as a yellow solid (0.42 g, 97%). Crystals suitable for X-ray diffraction were grown from methanol [m.p. 562–563 K (dec); literature value 562–564 K (dec)].

Refinement

H1NA and H1NB were located by a Fourier map and refined isotropically. All of the remaining H atoms were placed in their calculated positions and then refined using the riding model with Atom—H lengths of 0.95 Å (CH) or 0.98 Å (CH₃). Isotropic displacement parameters for these atoms were set to 1.19–1.20 (CH) or 1.4–1.50 (CH₃) times U_{eq} of the parent atom.

supplementary materials

Figures




Fig. 1. Molecular structure of the title compound with two molecules in the asymmetric unit showing the atom labeling scheme and 50% probability displacement ellipsoids.




Fig. 2. Packing diagram of the title compound viewed along the a axis. Dashed lines indicate N—H···O hydrogen bonds forming infinite 1-D chains along [011]. The remaining H atoms have been removed for clarity.

3,3-Dimethyl-1,2,3,4-tetrahydrocyclopenta[*b*]indole-1,2-dione

Crystal data

C ₁₃ H ₁₁ NO ₂	Z = 4
$M_r = 213.23$	$F(000) = 448$
Triclinic, PT	$D_x = 1.297 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.1091 (7) \text{ \AA}$	Cell parameters from 5810 reflections
$b = 11.5337 (8) \text{ \AA}$	$\theta = 3.3\text{--}32.3^\circ$
$c = 11.8745 (9) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\alpha = 63.230 (7)^\circ$	$T = 170 \text{ K}$
$\beta = 80.596 (6)^\circ$	Block, yellow
$\gamma = 79.970 (6)^\circ$	$0.28 \times 0.25 \times 0.24 \text{ mm}$
$V = 1091.69 (14) \text{ \AA}^3$	

Data collection

Oxford Diffraction Xcalibur Eos Gemini diffractometer	5644 independent reflections
Radiation source: Enhance (Mo) X-ray Source graphite	4698 reflections with $I > 2\sigma(I)$
Detector resolution: 16.1500 pixels mm^{-1}	$R_{\text{int}} = 0.012$
ω scans	$\theta_{\text{max}} = 28.7^\circ, \theta_{\text{min}} = 3.3^\circ$
Absorption correction: multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2010)	$h = -12 \rightarrow 11$
$T_{\text{min}} = 0.976, T_{\text{max}} = 0.979$	$k = -15 \rightarrow 15$
	$l = -15 \rightarrow 16$

10062 measured reflections

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.053$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.157$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0864P)^2 + 0.2278P]$ where $P = (F_o^2 + 2F_c^2)/3$
5644 reflections	$(\Delta/\sigma)_{\max} < 0.001$
299 parameters	$\Delta\rho_{\max} = 0.36 \text{ e \AA}^{-3}$
2 restraints	$\Delta\rho_{\min} = -0.20 \text{ e \AA}^{-3}$

Special details

Experimental. ^1H NMR (600 MHz, DMSO-d6) δ 12.9 (bs, 1H), 7.85 7.84 (d, $J = 8.0$ Hz, 1H), 7.62 7.61 (d, $J = 8.1$ Hz, 1H), 7.42 7.39 (t, $J = 7.4$ Hz, 1H), 7.34–7.31 (t, $J = 7.7$ Hz, 1H), 1.44 (s, 6H). ^{13}C NMR (150 MHz, DMSO-d6) δ 206.6, 175.2, 171.0, 140.0, 125.4, 123.4, 121.5, 121.1, 121.0, 113.6, 41.6, 22.9; IR v(KBr) 3418, 1750, 1665, 1469, 1453, 1210, 1152, 1094, 1013; UV λ_{\max} (95% MeOH) 256, 264, 272, 342 nm. Anal. Calcd for $\text{C}_{13}\text{H}_{11}\text{NO}_2$: C, 73.22; H, 5.20; N 6.57. Found: C, 72.48; H, 5.30; N 6.40.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.75420 (16)	0.86467 (16)	0.18664 (11)	0.0673 (4)
O2	1.05510 (14)	0.75658 (14)	0.25613 (11)	0.0591 (3)
N1	0.76832 (12)	0.67293 (11)	0.63752 (10)	0.0329 (2)
H1NA	0.6811 (16)	0.6811 (16)	0.6806 (15)	0.039*
C1	0.67431 (15)	0.78837 (13)	0.41344 (12)	0.0330 (3)
C2	0.78552 (17)	0.80931 (15)	0.29437 (13)	0.0413 (3)
C3	0.94630 (16)	0.75106 (15)	0.33304 (13)	0.0398 (3)
C4	0.93046 (14)	0.69760 (13)	0.46734 (12)	0.0330 (3)
C5	1.01358 (15)	0.62976 (14)	0.57632 (13)	0.0358 (3)
C6	1.16300 (17)	0.58148 (19)	0.59556 (18)	0.0522 (4)
H6A	1.2361	0.5907	0.5260	0.063*

supplementary materials

C7	1.2022 (2)	0.5200 (2)	0.7181 (2)	0.0653 (5)
H7A	1.3038	0.4862	0.7326	0.078*
C8	1.0970 (2)	0.5060 (2)	0.82122 (19)	0.0618 (5)
H8A	1.1280	0.4626	0.9044	0.074*
C9	0.94890 (19)	0.55398 (18)	0.80488 (15)	0.0496 (4)
H9A	0.8771	0.5453	0.8751	0.060*
C10	0.90872 (15)	0.61552 (14)	0.68187 (13)	0.0351 (3)
C11	0.78265 (14)	0.71996 (12)	0.51124 (11)	0.0289 (3)
C12	0.6013 (2)	0.92013 (15)	0.40795 (16)	0.0497 (4)
H12D	0.5338	0.9061	0.4851	0.074*
H12E	0.5442	0.9664	0.3338	0.074*
H12F	0.6791	0.9724	0.4014	0.074*
C13	0.55503 (18)	0.70394 (18)	0.42566 (17)	0.0495 (4)
H13D	0.4878	0.6899	0.5029	0.074*
H13E	0.6036	0.6194	0.4300	0.074*
H13F	0.4972	0.7484	0.3519	0.074*
N1A	0.10484 (12)	0.81468 (12)	1.00094 (10)	0.0343 (3)
H1NB	0.0732 (19)	0.8074 (16)	1.0812 (13)	0.041*
O1A	0.61629 (13)	0.64860 (15)	0.94255 (14)	0.0651 (4)
O2A	0.46978 (13)	0.73188 (12)	0.71755 (11)	0.0529 (3)
C1A	0.38431 (14)	0.72921 (13)	1.03441 (13)	0.0336 (3)
C2A	0.48880 (15)	0.69922 (14)	0.93239 (15)	0.0390 (3)
C3A	0.40683 (16)	0.74091 (13)	0.81363 (13)	0.0370 (3)
C4A	0.25841 (14)	0.78537 (13)	0.84578 (12)	0.0320 (3)
C5A	0.11410 (15)	0.83713 (13)	0.79856 (12)	0.0333 (3)
C6A	0.05507 (19)	0.87076 (16)	0.68526 (14)	0.0452 (3)
H6AA	0.1159	0.8600	0.6169	0.054*
C7A	-0.0938 (2)	0.92010 (18)	0.67500 (16)	0.0538 (4)
H7AA	-0.1353	0.9443	0.5980	0.065*
C8A	-0.18450 (19)	0.93526 (19)	0.77470 (18)	0.0568 (4)
H8AA	-0.2867	0.9692	0.7644	0.068*
C9A	-0.12946 (17)	0.90208 (18)	0.88854 (16)	0.0490 (4)
H9AA	-0.1916	0.9120	0.9568	0.059*
C10A	0.02018 (15)	0.85378 (13)	0.89824 (12)	0.0345 (3)
C11A	0.24447 (14)	0.77534 (12)	0.96840 (12)	0.0292 (3)
C12A	0.37519 (19)	0.60627 (17)	1.15941 (16)	0.0512 (4)
H12A	0.3434	0.5371	1.1456	0.077*
H12B	0.3025	0.6255	1.2211	0.077*
H12C	0.4738	0.5772	1.1919	0.077*
C13A	0.43957 (19)	0.83897 (17)	1.04967 (18)	0.0518 (4)
H13A	0.3713	0.8599	1.1125	0.078*
H13B	0.4424	0.9168	0.9681	0.078*
H13C	0.5403	0.8104	1.0781	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0641 (8)	0.0963 (10)	0.0291 (6)	0.0029 (7)	-0.0074 (5)	-0.0196 (6)

O2	0.0437 (6)	0.0938 (9)	0.0361 (6)	-0.0042 (6)	0.0117 (5)	-0.0315 (6)
N1	0.0271 (5)	0.0433 (6)	0.0241 (5)	-0.0003 (4)	0.0009 (4)	-0.0134 (4)
C1	0.0307 (6)	0.0383 (6)	0.0268 (6)	0.0014 (5)	-0.0042 (5)	-0.0127 (5)
C2	0.0429 (8)	0.0511 (8)	0.0273 (6)	-0.0021 (6)	-0.0012 (5)	-0.0165 (6)
C3	0.0350 (7)	0.0536 (8)	0.0310 (7)	-0.0033 (6)	0.0050 (5)	-0.0217 (6)
C4	0.0268 (6)	0.0428 (7)	0.0290 (6)	-0.0010 (5)	0.0014 (5)	-0.0175 (5)
C5	0.0284 (6)	0.0448 (7)	0.0356 (7)	0.0001 (5)	-0.0040 (5)	-0.0200 (6)
C6	0.0291 (7)	0.0757 (11)	0.0557 (10)	0.0045 (7)	-0.0059 (6)	-0.0351 (9)
C7	0.0378 (8)	0.0923 (14)	0.0728 (13)	0.0151 (9)	-0.0260 (9)	-0.0432 (11)
C8	0.0568 (11)	0.0806 (13)	0.0488 (10)	0.0086 (9)	-0.0269 (8)	-0.0273 (9)
C9	0.0471 (8)	0.0654 (10)	0.0334 (7)	0.0002 (7)	-0.0100 (6)	-0.0190 (7)
C10	0.0308 (6)	0.0429 (7)	0.0306 (6)	-0.0012 (5)	-0.0042 (5)	-0.0157 (5)
C11	0.0272 (6)	0.0335 (6)	0.0242 (6)	-0.0014 (4)	-0.0002 (4)	-0.0124 (5)
C12	0.0527 (9)	0.0409 (8)	0.0455 (8)	0.0076 (7)	-0.0038 (7)	-0.0149 (7)
C13	0.0394 (8)	0.0611 (9)	0.0523 (9)	-0.0075 (7)	-0.0085 (7)	-0.0263 (8)
N1A	0.0278 (5)	0.0469 (6)	0.0250 (5)	0.0028 (4)	-0.0006 (4)	-0.0159 (5)
O1A	0.0302 (6)	0.0908 (10)	0.0778 (9)	0.0113 (6)	-0.0046 (6)	-0.0463 (8)
O2A	0.0440 (6)	0.0689 (7)	0.0422 (6)	0.0004 (5)	0.0130 (5)	-0.0290 (6)
C1A	0.0280 (6)	0.0371 (6)	0.0338 (6)	0.0001 (5)	-0.0046 (5)	-0.0145 (5)
C2A	0.0272 (6)	0.0417 (7)	0.0461 (8)	-0.0024 (5)	0.0023 (5)	-0.0199 (6)
C3A	0.0342 (7)	0.0394 (7)	0.0345 (7)	-0.0039 (5)	0.0063 (5)	-0.0169 (5)
C4A	0.0311 (6)	0.0372 (6)	0.0257 (6)	-0.0032 (5)	0.0011 (5)	-0.0135 (5)
C5A	0.0327 (6)	0.0379 (6)	0.0271 (6)	-0.0041 (5)	-0.0013 (5)	-0.0126 (5)
C6A	0.0510 (9)	0.0542 (8)	0.0301 (7)	-0.0077 (7)	-0.0054 (6)	-0.0170 (6)
C7A	0.0543 (10)	0.0644 (10)	0.0390 (8)	-0.0065 (8)	-0.0201 (7)	-0.0142 (7)
C8A	0.0380 (8)	0.0717 (11)	0.0517 (10)	0.0019 (7)	-0.0157 (7)	-0.0179 (8)
C9A	0.0313 (7)	0.0662 (10)	0.0415 (8)	0.0043 (7)	-0.0037 (6)	-0.0199 (7)
C10A	0.0309 (6)	0.0406 (7)	0.0273 (6)	-0.0015 (5)	-0.0031 (5)	-0.0115 (5)
C11A	0.0266 (6)	0.0316 (6)	0.0261 (6)	-0.0010 (4)	0.0000 (4)	-0.0112 (5)
C12A	0.0402 (8)	0.0522 (9)	0.0417 (8)	0.0038 (6)	-0.0067 (6)	-0.0054 (7)
C13A	0.0453 (9)	0.0570 (9)	0.0638 (11)	-0.0044 (7)	-0.0134 (8)	-0.0336 (8)

Geometric parameters (\AA , $^\circ$)

O1—C2	1.2023 (18)	N1A—C11A	1.3301 (16)
O2—C3	1.2224 (17)	N1A—C10A	1.4057 (17)
N1—C11	1.3371 (16)	N1A—H1NB	0.919 (14)
N1—C10	1.4009 (17)	O1A—C2A	1.2042 (17)
N1—H1NA	0.887 (14)	O2A—C3A	1.2304 (17)
C1—C11	1.4910 (17)	C1A—C11A	1.4949 (17)
C1—C12	1.5260 (19)	C1A—C12A	1.525 (2)
C1—C13	1.530 (2)	C1A—C13A	1.532 (2)
C1—C2	1.5470 (19)	C1A—C2A	1.5424 (19)
C2—C3	1.551 (2)	C2A—C3A	1.545 (2)
C3—C4	1.4192 (19)	C3A—C4A	1.4151 (18)
C4—C11	1.3863 (17)	C4A—C11A	1.3937 (17)
C4—C5	1.4359 (19)	C4A—C5A	1.4370 (18)
C5—C6	1.3920 (19)	C5A—C6A	1.3939 (19)
C5—C10	1.4100 (19)	C5A—C10A	1.4095 (18)

supplementary materials

C6—C7	1.379 (3)	C6A—C7A	1.378 (2)
C6—H6A	0.9500	C6A—H6AA	0.9500
C7—C8	1.392 (3)	C7A—C8A	1.390 (3)
C7—H7A	0.9500	C7A—H7AA	0.9500
C8—C9	1.376 (2)	C8A—C9A	1.383 (2)
C8—H8A	0.9500	C8A—H8AA	0.9500
C9—C10	1.386 (2)	C9A—C10A	1.3822 (19)
C9—H9A	0.9500	C9A—H9AA	0.9500
C12—H12D	0.9800	C12A—H12A	0.9800
C12—H12E	0.9800	C12A—H12B	0.9800
C12—H12F	0.9800	C12A—H12C	0.9800
C13—H13D	0.9800	C13A—H13A	0.9800
C13—H13E	0.9800	C13A—H13B	0.9800
C13—H13F	0.9800	C13A—H13C	0.9800
C11—N1—C10	108.76 (11)	C11A—N1A—C10A	108.31 (11)
C11—N1—H1NA	121.8 (11)	C11A—N1A—H1NB	122.5 (11)
C10—N1—H1NA	129.4 (11)	C10A—N1A—H1NB	129.0 (11)
C11—C1—C12	113.21 (12)	C11A—C1A—C12A	114.55 (12)
C11—C1—C13	113.17 (12)	C11A—C1A—C13A	111.77 (11)
C12—C1—C13	110.52 (13)	C12A—C1A—C13A	111.60 (14)
C11—C1—C2	98.41 (10)	C11A—C1A—C2A	97.98 (10)
C12—C1—C2	109.97 (12)	C12A—C1A—C2A	110.59 (12)
C13—C1—C2	110.99 (12)	C13A—C1A—C2A	109.51 (12)
O1—C2—C1	125.61 (14)	O1A—C2A—C1A	125.67 (14)
O1—C2—C3	124.13 (14)	O1A—C2A—C3A	123.63 (14)
C1—C2—C3	110.25 (11)	C1A—C2A—C3A	110.69 (11)
O2—C3—C4	132.30 (14)	O2A—C3A—C4A	132.76 (14)
O2—C3—C2	123.08 (13)	O2A—C3A—C2A	122.40 (13)
C4—C3—C2	104.62 (11)	C4A—C3A—C2A	104.84 (11)
C11—C4—C3	110.28 (12)	C11A—C4A—C3A	109.89 (12)
C11—C4—C5	107.02 (11)	C11A—C4A—C5A	107.00 (11)
C3—C4—C5	142.68 (12)	C3A—C4A—C5A	143.11 (13)
C6—C5—C10	119.17 (14)	C6A—C5A—C10A	119.22 (13)
C6—C5—C4	135.02 (13)	C6A—C5A—C4A	135.42 (13)
C10—C5—C4	105.80 (11)	C10A—C5A—C4A	105.36 (11)
C7—C6—C5	118.31 (16)	C7A—C6A—C5A	118.32 (15)
C7—C6—H6A	120.8	C7A—C6A—H6AA	120.8
C5—C6—H6A	120.8	C5A—C6A—H6AA	120.8
C6—C7—C8	121.72 (16)	C6A—C7A—C8A	121.50 (14)
C6—C7—H7A	119.1	C6A—C7A—H7AA	119.3
C8—C7—H7A	119.1	C8A—C7A—H7AA	119.3
C9—C8—C7	121.20 (16)	C9A—C8A—C7A	121.55 (15)
C9—C8—H8A	119.4	C9A—C8A—H8AA	119.2
C7—C8—H8A	119.4	C7A—C8A—H8AA	119.2
C8—C9—C10	117.32 (16)	C10A—C9A—C8A	116.88 (15)
C8—C9—H9A	121.3	C10A—C9A—H9AA	121.6
C10—C9—H9A	121.3	C8A—C9A—H9AA	121.6
C9—C10—N1	129.67 (13)	C9A—C10A—N1A	128.81 (13)
C9—C10—C5	122.27 (13)	C9A—C10A—C5A	122.53 (13)

N1—C10—C5	108.06 (11)	N1A—C10A—C5A	108.66 (11)
N1—C11—C4	110.35 (11)	N1A—C11A—C4A	110.66 (11)
N1—C11—C1	133.24 (11)	N1A—C11A—C1A	132.96 (12)
C4—C11—C1	116.41 (11)	C4A—C11A—C1A	116.33 (11)
C1—C12—H12D	109.5	C1A—C12A—H12A	109.5
C1—C12—H12E	109.5	C1A—C12A—H12B	109.5
H12D—C12—H12E	109.5	H12A—C12A—H12B	109.5
C1—C12—H12F	109.5	C1A—C12A—H12C	109.5
H12D—C12—H12F	109.5	H12A—C12A—H12C	109.5
H12E—C12—H12F	109.5	H12B—C12A—H12C	109.5
C1—C13—H13D	109.5	C1A—C13A—H13A	109.5
C1—C13—H13E	109.5	C1A—C13A—H13B	109.5
H13D—C13—H13E	109.5	H13A—C13A—H13B	109.5
C1—C13—H13F	109.5	C1A—C13A—H13C	109.5
H13D—C13—H13F	109.5	H13A—C13A—H13C	109.5
H13E—C13—H13F	109.5	H13B—C13A—H13C	109.5
C11—C1—C2—O1	−178.28 (17)	C11A—C1A—C2A—O1A	−174.04 (15)
C12—C1—C2—O1	−59.7 (2)	C12A—C1A—C2A—O1A	−54.0 (2)
C13—C1—C2—O1	62.9 (2)	C13A—C1A—C2A—O1A	69.41 (19)
C11—C1—C2—C3	0.43 (15)	C11A—C1A—C2A—C3A	4.93 (13)
C12—C1—C2—C3	118.96 (14)	C12A—C1A—C2A—C3A	124.99 (13)
C13—C1—C2—C3	−118.44 (13)	C13A—C1A—C2A—C3A	−111.62 (13)
O1—C2—C3—O2	−0.2 (3)	O1A—C2A—C3A—O2A	−4.3 (2)
C1—C2—C3—O2	−178.95 (15)	C1A—C2A—C3A—O2A	176.71 (13)
O1—C2—C3—C4	179.30 (17)	O1A—C2A—C3A—C4A	175.40 (15)
C1—C2—C3—C4	0.57 (16)	C1A—C2A—C3A—C4A	−3.60 (15)
O2—C3—C4—C11	178.02 (17)	O2A—C3A—C4A—C11A	−179.98 (16)
C2—C3—C4—C11	−1.44 (16)	C2A—C3A—C4A—C11A	0.37 (15)
O2—C3—C4—C5	−0.2 (3)	O2A—C3A—C4A—C5A	0.6 (3)
C2—C3—C4—C5	−179.69 (18)	C2A—C3A—C4A—C5A	−179.08 (17)
C11—C4—C5—C6	−179.60 (17)	C11A—C4A—C5A—C6A	179.17 (16)
C3—C4—C5—C6	−1.3 (3)	C3A—C4A—C5A—C6A	−1.4 (3)
C11—C4—C5—C10	−0.31 (15)	C11A—C4A—C5A—C10A	−0.72 (15)
C3—C4—C5—C10	177.97 (19)	C3A—C4A—C5A—C10A	178.74 (17)
C10—C5—C6—C7	0.7 (3)	C10A—C5A—C6A—C7A	0.4 (2)
C4—C5—C6—C7	179.91 (18)	C4A—C5A—C6A—C7A	−179.47 (16)
C5—C6—C7—C8	−0.3 (3)	C5A—C6A—C7A—C8A	−0.6 (3)
C6—C7—C8—C9	−0.4 (3)	C6A—C7A—C8A—C9A	0.3 (3)
C7—C8—C9—C10	0.6 (3)	C7A—C8A—C9A—C10A	0.3 (3)
C8—C9—C10—N1	−179.92 (16)	C8A—C9A—C10A—N1A	178.99 (15)
C8—C9—C10—C5	−0.2 (3)	C8A—C9A—C10A—C5A	−0.6 (3)
C11—N1—C10—C9	−179.70 (16)	C11A—N1A—C10A—C9A	−179.67 (15)
C11—N1—C10—C5	0.51 (16)	C11A—N1A—C10A—C5A	−0.06 (16)
C6—C5—C10—C9	−0.5 (2)	C6A—C5A—C10A—C9A	0.2 (2)
C4—C5—C10—C9	−179.92 (15)	C4A—C5A—C10A—C9A	−179.88 (14)
C6—C5—C10—N1	179.31 (14)	C6A—C5A—C10A—N1A	−179.42 (13)
C4—C5—C10—N1	−0.11 (15)	C4A—C5A—C10A—N1A	0.48 (15)
C10—N1—C11—C4	−0.73 (15)	C10A—N1A—C11A—C4A	−0.41 (15)
C10—N1—C11—C1	179.04 (14)	C10A—N1A—C11A—C1A	176.95 (13)

supplementary materials


C3—C4—C11—N1	−178.24 (12)	C3A—C4A—C11A—N1A	−178.94 (11)
C5—C4—C11—N1	0.65 (16)	C5A—C4A—C11A—N1A	0.72 (15)
C3—C4—C11—C1	1.95 (17)	C3A—C4A—C11A—C1A	3.22 (16)
C5—C4—C11—C1	−179.16 (11)	C5A—C4A—C11A—C1A	−177.13 (11)
C12—C1—C11—N1	62.8 (2)	C12A—C1A—C11A—N1A	60.7 (2)
C13—C1—C11—N1	−63.97 (19)	C13A—C1A—C11A—N1A	−67.49 (19)
C2—C1—C11—N1	178.83 (15)	C2A—C1A—C11A—N1A	177.73 (14)
C12—C1—C11—C4	−117.46 (14)	C12A—C1A—C11A—C4A	−122.05 (14)
C13—C1—C11—C4	115.79 (14)	C13A—C1A—C11A—C4A	109.76 (14)
C2—C1—C11—C4	−1.41 (15)	C2A—C1A—C11A—C4A	−5.02 (14)

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1NA…O2A	0.89 (1)	1.95 (1)	2.8025 (16)	161.(2)
N1A—H1NB…O2 ⁱ	0.92 (1)	1.87 (1)	2.7686 (16)	164.(2)
C9A—H9AA…O1 ⁱ	0.95	2.51	3.376 (2)	152.
C12A—H12B…O2 ⁱ	0.98	2.57	3.430 (2)	146.

Symmetry codes: (i) $x-1, y, z+1$.

Fig. 1



supplementary materials

Fig. 2

