

1-(4-Methylphenyl)-1*H*-1,2,3,4-tetrazole

Kwan Baek,^a D. Gayathri,^b Vivek K. Gupta,^c Rajni Kant^c and Yeon Tae Jeong^{a*}

^aDepartment of Image Science and Engineering, Pukyong National University, Busan 608 739, Republic of Korea, ^bDepartment of Physics, Dr. M.G.R Educational and Research Institute, Dr. M.G.R University, Maduravoyal, Chennai 600 095, India, and ^cX-ray Crystallography Laboratory, Post Graduate Department of Physics & Electronics, University of Jammu, Jammu Tawi 180 006, India
Correspondence e-mail: ytjeong@pknu.ac.kr


Received 5 January 2012; accepted 9 January 2012

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.067; wR factor = 0.211; data-to-parameter ratio = 12.9.

In the title compound, $\text{C}_8\text{H}_8\text{N}_4$, the dihedral angle between the tetrazole and benzene rings is $21.6(1)^\circ$. An intermolecular $\text{C}-\text{H}\cdots\pi$ interaction is observed.

Related literature

For background to and applications of tetrazole derivatives, see: Singh *et al.* (1980); Brown (1967); Ostrovskii *et al.* (1999). For the synthesis, see: Aridoss & Laali (2011). For related structures, see: Matsunaga *et al.* (1999); Lyakhov *et al.* (2000).



Experimental

Crystal data

$\text{C}_8\text{H}_8\text{N}_4$	$V = 806.92(17)\text{ \AA}^3$
$M_r = 160.18$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 9.8352(13)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 5.7244(6)\text{ \AA}$	$T = 293\text{ K}$
$c = 14.4190(19)\text{ \AA}$	$0.3 \times 0.2 \times 0.1\text{ mm}$
$\beta = 96.285(12)^\circ$	

Data collection

Oxford Diffraction Xcalibur Sapphire3 diffractometer	14618 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2010)	1419 independent reflections
$T_{\min} = 0.762$, $T_{\max} = 1.000$	936 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.067$	110 parameters
$wR(F^2) = 0.211$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\max} = 0.21\text{ e \AA}^{-3}$
1419 reflections	$\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

C_g is the centroid of the C2–C7 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}6-\text{H}6\cdots Cg^i$	0.93	2.89	3.630 (3)	138
Symmetry code: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$.				

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

KB and YTJ acknowledge the support provided by the second stage of the BK21 Program. RK thanks the DST, New Delhi, India, for the X-ray data collection facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5047).

References

- Aridoss, G. & Laali, K. K. (2011). *Eur. J. Org. Chem.* pp. 2827–2835.
Brown, M. (1967). US Patent No. 3 338 915.
Lyakhov, A. S., Ivashkevich, D. O., Gaponik, P. N., Grigoriev, Y. V. & Ivashkevich, L. S. (2000). *Acta Cryst. C56*, 256–257.
Matsunaga, T., Ohno, Y., Akutsu, Y., Arai, M., Tamura, M. & Iida, M. (1999). *Acta Cryst. C55*, 129–131.
Ostrovskii, V. A., Pevzner, M. S., Kofmna, T. P., Shcherbinin, M. B. & Tselsinskii, I. V. (1999). *Targets Heterocycl. Syst. 3*, 467–526.
Oxford Diffraction (2010). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Singh, H., Chawla, A. S., Kapoor, V. K., Paul, D. & Malhotra, R. K. (1980). *Prog. Med. Chem.* **17**, 151–183.
Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2012). E68, o394 [doi:10.1107/S1600536812000797]

1-(4-Methylphenyl)-1*H*-1,2,3,4-tetrazole

K. Baek, D. Gayathri, V. K. Gupta, R. Kant and Y. T. Jeong

Comment

Tetrazoles are an important functional group with wide range of applications (Aridoss & Laali, 2011). They function as ligands in coordination chemistry, in medicinal chemistry as a metabolically stable surrogate for a carboxylic acid group (Singh *et al.*, 1980) and in materials science applications including propellants (Brown, 1967) and explosives (Ostrovskii *et al.*, 1999).

Bond lengths and bond angles are comparable with the similar crystal structures (Matsunaga *et al.*, 1999; Lyakhov *et al.*, 2000). The tetrazole and benzene rings are planar with maximum deviations of 0.006 (2) and 0.011 (2) Å (r.m.s. deviations of the rings being 0.005 and 0.008 Å), respectively. The two rings are not coplanar with the dihedral angle being 21.6 (1)°. Methyl carbon atom lies 0.014 (5) Å from the plane of the phenyl ring. Bond distances C1—N1 [1.304 (4) Å] and N2—N3 [1.289 (4) Å] indicate the consistence of the formation of double bonds.

Experimental

The title compound was synthesized from the known procedure reported elsewhere (Aridoss & Laali, 2011). Fine white diffraction quality crystals were obtained from the slow evaporation of its solution in ethanol.

Refinement

All H atoms were refined using a riding model, with C—H = 0.93 Å (aromatic) or 0.96 Å (methyl), and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids.

1-(4-Methylphenyl)-1*H*-1,2,3,4-tetrazole

Crystal data

$\text{C}_8\text{H}_8\text{N}_4$	$F(000) = 336$
$M_r = 160.18$	$D_x = 1.319 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 3527 reflections
$a = 9.8352 (13) \text{ \AA}$	$\theta = 3.6\text{--}29.2^\circ$

supplementary materials

$b = 5.7244 (6) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 14.4190 (19) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 96.285 (12)^\circ$	Plate, white
$V = 806.92 (17) \text{ \AA}^3$	$0.3 \times 0.2 \times 0.1 \text{ mm}$
$Z = 4$	

Data collection

Oxford Diffraction Xcalibur Sapphire3 diffractometer	1419 independent reflections
Radiation source: fine-focus sealed tube graphite	936 reflections with $I > 2\sigma(I)$
Detector resolution: 16.1049 pixels mm^{-1}	$R_{\text{int}} = 0.054$
ω scans	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 3.8^\circ$
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2010)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.762, T_{\text{max}} = 1.000$	$k = -6 \rightarrow 6$
14618 measured reflections	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.067$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.211$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.1133P)^2 + 0.1542P]$ where $P = (F_o^2 + 2F_c^2)/3$
1419 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
110 parameters	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

x	y	z	$U_{\text{iso}}^* / U_{\text{eq}}$
-----	-----	-----	------------------------------------

C1	0.7153 (3)	0.0846 (5)	0.8500 (2)	0.0726 (9)
H1	0.7271	-0.0729	0.8372	0.087*
C2	0.4656 (3)	0.0810 (4)	0.87347 (17)	0.0540 (7)
C3	0.3730 (3)	0.1934 (5)	0.92335 (18)	0.0635 (8)
H3	0.3959	0.3336	0.9537	0.076*
C4	0.2465 (3)	0.0950 (5)	0.9274 (2)	0.0672 (9)
H4	0.1838	0.1718	0.9605	0.081*
C5	0.2090 (3)	-0.1143 (5)	0.88414 (19)	0.0622 (8)
C6	0.3039 (3)	-0.2219 (5)	0.8337 (2)	0.0634 (8)
H6	0.2811	-0.3625	0.8036	0.076*
C7	0.4315 (3)	-0.1253 (4)	0.82696 (18)	0.0607 (8)
H7	0.4932	-0.1980	0.7918	0.073*
C8	0.0709 (3)	-0.2228 (6)	0.8897 (2)	0.0850 (10)
H8A	0.0534	-0.2328	0.9538	0.127*
H8B	0.0692	-0.3767	0.8631	0.127*
H8C	0.0017	-0.1283	0.8558	0.127*
N1	0.8114 (3)	0.2424 (5)	0.85255 (19)	0.0841 (9)
N2	0.7502 (3)	0.4443 (5)	0.8748 (2)	0.0874 (9)
N3	0.6234 (3)	0.4089 (4)	0.8848 (2)	0.0826 (9)
N4	0.5985 (2)	0.1809 (4)	0.86824 (14)	0.0580 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.077 (2)	0.0625 (18)	0.081 (2)	0.0000 (16)	0.0255 (17)	-0.0042 (15)
C2	0.0584 (16)	0.0504 (15)	0.0528 (15)	0.0089 (11)	0.0041 (12)	0.0054 (11)
C3	0.079 (2)	0.0557 (16)	0.0568 (17)	0.0051 (14)	0.0129 (14)	-0.0082 (13)
C4	0.071 (2)	0.0697 (19)	0.0636 (18)	0.0125 (15)	0.0181 (15)	-0.0055 (14)
C5	0.0616 (18)	0.0644 (17)	0.0609 (17)	0.0079 (13)	0.0075 (14)	0.0067 (13)
C6	0.0678 (18)	0.0531 (16)	0.0686 (18)	0.0045 (13)	0.0046 (14)	-0.0049 (13)
C7	0.0661 (19)	0.0524 (15)	0.0635 (18)	0.0138 (13)	0.0073 (14)	-0.0052 (12)
C8	0.066 (2)	0.101 (3)	0.088 (2)	-0.0003 (17)	0.0104 (17)	-0.0046 (19)
N1	0.0828 (19)	0.086 (2)	0.087 (2)	-0.0101 (15)	0.0223 (15)	0.0017 (14)
N2	0.085 (2)	0.0721 (18)	0.106 (2)	-0.0069 (15)	0.0121 (16)	0.0022 (15)
N3	0.082 (2)	0.0569 (16)	0.108 (2)	0.0002 (13)	0.0060 (16)	-0.0030 (13)
N4	0.0649 (15)	0.0519 (13)	0.0568 (14)	0.0047 (11)	0.0048 (11)	0.0029 (10)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.304 (4)	C5—C8	1.504 (4)
C1—N4	1.326 (3)	C6—C7	1.385 (4)
C1—H1	0.9300	C6—H6	0.9300
C2—C3	1.380 (4)	C7—H7	0.9300
C2—C7	1.381 (4)	C8—H8A	0.9600
C2—N4	1.436 (3)	C8—H8B	0.9600
C3—C4	1.373 (4)	C8—H8C	0.9600
C3—H3	0.9300	N1—N2	1.357 (4)
C4—C5	1.382 (4)	N2—N3	1.288 (4)
C4—H4	0.9300	N3—N4	1.345 (3)

supplementary materials

C5—C6	1.389 (4)		
N1—C1—N4	110.3 (3)	C5—C6—H6	119.1
N1—C1—H1	124.8	C2—C7—C6	118.8 (3)
N4—C1—H1	124.8	C2—C7—H7	120.6
C3—C2—C7	120.8 (3)	C6—C7—H7	120.6
C3—C2—N4	119.9 (2)	C5—C8—H8A	109.5
C7—C2—N4	119.2 (2)	C5—C8—H8B	109.5
C4—C3—C2	119.0 (3)	H8A—C8—H8B	109.5
C4—C3—H3	120.5	C5—C8—H8C	109.5
C2—C3—H3	120.5	H8A—C8—H8C	109.5
C3—C4—C5	122.3 (2)	H8B—C8—H8C	109.5
C3—C4—H4	118.8	C1—N1—N2	105.0 (3)
C5—C4—H4	118.8	N3—N2—N1	110.6 (2)
C4—C5—C6	117.4 (3)	N2—N3—N4	107.0 (2)
C4—C5—C8	122.1 (3)	C1—N4—N3	107.1 (2)
C6—C5—C8	120.5 (3)	C1—N4—C2	131.2 (2)
C7—C6—C5	121.7 (3)	N3—N4—C2	121.7 (2)
C7—C6—H6	119.1		
C7—C2—C3—C4	1.0 (4)	C1—N1—N2—N3	-0.1 (4)
N4—C2—C3—C4	180.0 (2)	N1—N2—N3—N4	-0.6 (4)
C2—C3—C4—C5	0.7 (4)	N1—C1—N4—N3	-1.2 (3)
C3—C4—C5—C6	-1.4 (4)	N1—C1—N4—C2	180.0 (3)
C3—C4—C5—C8	179.3 (3)	N2—N3—N4—C1	1.1 (3)
C4—C5—C6—C7	0.3 (4)	N2—N3—N4—C2	-180.0 (2)
C8—C5—C6—C7	179.7 (3)	C3—C2—N4—C1	157.9 (3)
C3—C2—C7—C6	-2.0 (4)	C7—C2—N4—C1	-23.2 (4)
N4—C2—C7—C6	179.0 (2)	C3—C2—N4—N3	-20.7 (4)
C5—C6—C7—C2	1.3 (4)	C7—C2—N4—N3	158.2 (2)
N4—C1—N1—N2	0.9 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

Cg is the centroid of the C2—C7 ring.

$D\cdots H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C6—H6 \cdots Cg ⁱ	0.93	2.89	3.630 (3)	138

Symmetry codes: (i) $-x+1/2, y-1/2, -z+3/2$.

Fig. 1

