

N'-(*E*)-2-Hydroxy-5-iodobenzylidene]-furan-2-carbohydrazone monohydrate

Rahman Bikas,^{a‡} Parisa Mahboubi Anarjan,^b Seik Weng Ng^{c,d} and Edward R. T. Tiekink^{c*}

^aYoung Researchers Club, Tabriz Branch, Islamic Azad University, Tabriz, Iran,

^bDepartment of Chemistry, University of Zanjan, 45195-313 Zanjan, Iran,

^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia,

and ^dChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: edward.tiekink@gmail.com


Received 20 December 2011; accepted 27 December 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; disorder in main residue; R factor = 0.059; wR factor = 0.158; data-to-parameter ratio = 14.2.

The organic molecule of the title monohydrate, $\text{C}_{12}\text{H}_9\text{IN}_2\text{O}_3 \cdot \text{H}_2\text{O}$, features a disordered furyl ring with the major component [site occupancy = 0.575 (18)] having the carbonyl O and furyl O atoms *syn*, and the other conformation having these atoms *anti*. The molecule is slightly twisted with the dihedral angle between the benzene and furyl rings being 10.3 (6)° (major component). An intramolecular O—H...N(imine) hydrogen bond is formed. In the crystal, the water molecule accepts a hydrogen bond from an amine H atom, and forms two O—H...O(carbonyl) hydrogen bonds, thereby linking three different carbohydrazone molecules. The result is a supramolecular layer parallel to (001). The closest contacts between layers are of the type I...I, at a distance of 3.6986 (6) Å.

Related literature

For historical background to aroylhydrazones, see: Craliz *et al.* (1955). For the structure of the isomorphous bromido derivative, see: Tai *et al.* (2007). For the structures of related carbohydrazides, see: Abdel-Aziz *et al.* (2011); Bikas *et al.* (2012). For the synthesis of a precursor molecule, see: Nielsen & Gothelf (2001).



[‡] Additional correspondence author, e-mail: bikas_r@yahoo.com.

Experimental

Crystal data

$\text{C}_{12}\text{H}_9\text{IN}_2\text{O}_3 \cdot \text{H}_2\text{O}$

$M_r = 374.13$

Orthorhombic, $P2_12_12_1$

$a = 4.8607$ (2) Å

$b = 12.5873$ (4) Å

$c = 21.1627$ (9) Å

$V = 1294.80$ (9) Å³

$Z = 4$

Cu $K\alpha$ radiation

$\mu = 19.57$ mm⁻¹

$T = 100$ K

$0.20 \times 0.08 \times 0.04$ mm

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector

Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)

$T_{\min} = 0.111$, $T_{\max} = 0.508$

4758 measured reflections

2641 independent reflections

2575 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$

$wR(F^2) = 0.158$

$S = 1.09$

2641 reflections

186 parameters

34 restraints

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 3.22$ e Å⁻³

$\Delta\rho_{\text{min}} = -1.82$ e Å⁻³

Absolute structure: Flack (1983),

1050 Friedel pairs

Flack parameter: -0.020 (12)

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2...O1w	0.88	1.96	2.813 (7)	163
O1—H1...N1	0.84	2.20	2.744 (8)	122
O1w—H11...O2 ⁱ	0.84	1.99	2.815 (8)	167
O1w—H12...O2 ⁱⁱ	0.84	2.00	2.826 (8)	168

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors are grateful to the Islamic Azad University, Tabriz Branch, the University of Zanjan and the University of Malaya for support of this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6573).

References

- Abdel-Aziz, H. A., Ng, S. W. & Tiekink, E. R. T. (2011). *Acta Cryst.* **E67**, o2317–o2318.
- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bikas, R., Anarjan, P. M., Ng, S. W. & Tiekink, E. R. T. (2012). *Acta Cryst.* **E68**, o193.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Craliz, J. C., Rub, J. C., Willis, D. & Edger, J. (1955). *Nature (London)*, **34**, 176.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.

Nielsen, M. & Gothelf, K. V. (2001). *J. Chem. Soc. Perkin Trans. 1*, pp. 2440–2444.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

Tai, X.-S., Yin, J., Hao, M.-Y. & Liang, Z.-P. (2007). *Acta Cryst. E* **63**, o2144–o2145.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o413-o414 [doi:10.1107/S1600536811055826]

N'-[(*E*)-2-Hydroxy-5-iodobenzylidene]furan-2-carbohydrazide monohydrate

R. Bikas, P. M. Anarjan, S. W. Ng and E. R. T. Tiekink

Comment

Hydrazone ligands (carbohydrazides), a class of Schiff base, derived from the condensation of acid hydrazides (R–CO–NH–NH₂) with aromatic 2-hydroxy aldehydes or ketones (Craliz *et al.*, 1955) are potentially important tridentate *O,N,O*-donor ligands. Previous structural studies of carbohydrazide derivatives (Abdel-Aziz *et al.*, 2011; Bikas *et al.*, 2012) have been extended to include the title compound, (I), which is isomorphous with bromido derivative (Tai *et al.*, 2007).

In the molecule of (I), Fig. 1, the furyl ring was found to be disordered over two almost diagonally opposed orientations with the dihedral angle between the two components being 9.5 (10)°. In the major component (site occupancy = 0.575 (18)), the carbonyl-O and furyl-O atoms are *syn*. Overall, the molecule of (I) exhibits a small twist with the dihedral angle between the benzene and furyl rings being 10.3 (6)° for the major component; the comparable angle involving the minor component = 15.9 (8)°. The hydroxyl-H atom forms an intramolecular hydrogen bond to the imine-H atom, Table 1. The conformation about the C7=N1 imine bond [1.283 (10) Å] is *E*.

In the crystal packing, the amine-H atoms forms a hydrogen bond to the water-O, and the water-H atoms form hydrogen bonds to carbonyl-O atoms from two different molecules, Table 1. The result is a supramolecular layers parallel to (001) comprising alternating rows of carbohydrazide molecules and water molecules, Fig. 2. The layers stack along the *c* axis with the closest contacts between them being I...I interactions [3.6986 (6) Å for symmetry operation: $-1/2 + x, 5/2 - y, 2 - z$], Fig. 3.

Experimental

2-Hydroxy-5-iodobenzaldehyde was synthesized according to the reported procedure by Nielsen & Gothelf (2001). For preparing the title compound a methanol (10 ml) solution of 2-hydroxy-5-iodobenzaldehyde (1.5 mmol) was added drop-wise to a methanol solution (10 ml) of 2-furanecarboxylic acid hydrazide (1.5 mmol), and the mixture was refluxed for 3 h. The solution was then evaporated on a steam bath to 5 cm³ and cooled to room temperature. The light-yellow precipitates of the title compound were separated and filtered off, washed with 3 ml of cooled methanol and then dried in air. Colourless crystals were obtained from its methanol:water (98:2 *v/v*) solution by slow solvent evaporation. Yield: 86%. IR (cm⁻¹): 3447 (w, broad, —OH), 3262 (m, N—H); 1668 (*versus*, C=O); 1609 (s, C=N(azomethine)); 952 (m, N—N); 1274, 1351 (*versus*, C—O enolate). ¹H NMR (250.13 MHz; DMSO-*d*₆): δ 12.16 (s, 1H, CO—NH—); 11.15 (s, 1H, —OH); 8.57 (s, 1H); 7.92 (s, 1H); 7.88 (s, 1H); 7.51 (d, 1H, *J* = 8.5 Hz); 7.30 (s, 1H); 7.74 (d, 1H, *J* = 8.75 Hz); 6.67 (s, 1H) p.p.m.. ¹H NMR (250.13 MHz; DMSO-*d*₆ + D₂O): δ 8.85 (s, 1H); 7.83 (s, 1H); 7.382 (s, 1H); 7.24 (s, 1H); 6.72 (d, 1H, *J* = 8.75 Hz); 6.64 (s, 1H) p.p.m.. ¹³C NMR (DMSO; 62.90 MHz): δ 81.72, 112.61, 115.82, 119.48, 122.35, 136.62, 139.76, 145.99, 146.52, 146.63, 154.56 and 157.35 p.p.m..

Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H 0.95 Å, N—H 0.88 Å, O—H 0.84 Å; $U_{\text{iso}}(\text{H})$ $1.2U_{\text{eq}}(\text{C}, \text{N}, \text{O})$] and were included in the refinement in the riding model approximation.

The H-atoms of the water molecule were placed in chemically sensible positions on the basis of hydrogen bonds but were not refined; $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

The furyl ring is disordered over two positions in a 0.575 (18): 0.425 (18) ratio. The C—O distances were restrained to 1.37 ± 0.01 Å, the carbon–carbon single-bond distances to 1.42 ± 0.01 Å and the carbon–carbon double-bond distances to 1.34 ± 0.01 Å. The α -carbon atom is ordered. The U_{iso} of the atoms comprising the minor component were set to U_{eq} of those of the atoms of the major component which were refined anisotropically.

The final difference Fourier map had a peak at approximately 1 Å from I1 and a hole at approximately 1 Å from I1.

Figures




Fig. 1. Molecular structure of (I) with displacement ellipsoids at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. For reasons of clarity, only the major component of the furyl ring is shown.




Fig. 2. Supramolecular layer parallel to (001) in (I). The intermolecular O—H...O and N—H...O hydrogen bonds are shown as orange and blue dashed lines, respectively. For reasons of clarity, only the major component of the furyl ring is shown.




Fig. 3. A view of the unit-cell contents of (I) in projection down the a axis. The intra- and inter-molecular O—H...O and the N—H...O hydrogen bonds are shown as orange and blue dashed lines, respectively. For reasons of clarity, only the major component of the furyl ring is shown.

N'-[*E*]-2-Hydroxy-5-iodobenzylidene]furan-2-carbohydrazide monohydrate

Crystal data

$\text{C}_{12}\text{H}_9\text{IN}_2\text{O}_3 \cdot \text{H}_2\text{O}$

$M_r = 374.13$

Orthorhombic, $P2_12_12_1$

$F(000) = 728$

$D_x = 1.919 \text{ Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å

Hall symbol: P 2ac 2ab
 $a = 4.8607$ (2) Å
 $b = 12.5873$ (4) Å
 $c = 21.1627$ (9) Å
 $V = 1294.80$ (9) Å³
 $Z = 4$

Cell parameters from 2924 reflections
 $\theta = 4.1\text{--}76.3^\circ$
 $\mu = 19.57$ mm⁻¹
 $T = 100$ K
 Prism, colourless
 $0.20 \times 0.08 \times 0.04$ mm

Data collection

Agilent SuperNova Dual
 diffractometer with an Atlas detector
 Radiation source: SuperNova (Cu) X-ray Source
 Mirror
 Detector resolution: 10.4041 pixels mm⁻¹
 ω scan
 Absorption correction: multi-scan
 (*Crys.Alis PRO*; Agilent, 2010)
 $T_{\min} = 0.111$, $T_{\max} = 0.508$
 4758 measured reflections

2641 independent reflections
 2575 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$
 $\theta_{\max} = 76.5^\circ$, $\theta_{\min} = 4.1^\circ$
 $h = -6 \rightarrow 5$
 $k = -15 \rightarrow 10$
 $l = -18 \rightarrow 26$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.158$
 $S = 1.09$
 2641 reflections
 186 parameters
 34 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.127P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 3.22$ e Å⁻³
 $\Delta\rho_{\min} = -1.82$ e Å⁻³
 Absolute structure: Flack (1983), 1050 Friedel pairs
 Flack parameter: -0.020 (12)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
II	1.44774 (10)	1.14557 (3)	0.97807 (2)	0.0213 (2)	
O1	0.7447 (14)	0.7625 (4)	0.8846 (3)	0.0239 (12)	
H1	0.7161	0.7655	0.8455	0.036*	
O2	0.1609 (12)	0.7779 (4)	0.7326 (3)	0.0205 (10)	
O3	-0.214 (2)	0.8642 (7)	0.6509 (4)	0.016 (2)	0.575 (18)
O3'	-0.043 (3)	1.0345 (7)	0.6855 (7)	0.016*	0.425 (18)
O1W	0.3405 (13)	1.1654 (4)	0.7575 (3)	0.0265 (12)	
H11	0.4756	1.2055	0.7641	0.040*	
H12	0.1948	1.1982	0.7664	0.040*	
N1	0.5129 (13)	0.9031 (5)	0.8010 (3)	0.0170 (12)	

supplementary materials

N2	0.3308 (13)	0.9420 (4)	0.7566 (3)	0.0158 (11)	
H2	0.3265	1.0105	0.7483	0.019*	
C1	0.8927 (16)	0.8472 (6)	0.9027 (3)	0.0196 (14)	
C2	1.0829 (17)	0.8331 (6)	0.9520 (4)	0.0222 (14)	
H2A	1.1050	0.7648	0.9704	0.027*	
C3	1.2403 (16)	0.9185 (6)	0.9745 (3)	0.0196 (14)	
H3	1.3645	0.9086	1.0087	0.024*	
C4	1.2132 (15)	1.0178 (6)	0.9464 (3)	0.0168 (13)	
C5	1.0320 (17)	1.0338 (5)	0.8971 (3)	0.0176 (13)	
H5	1.0188	1.1018	0.8779	0.021*	
C6	0.8665 (15)	0.9496 (6)	0.8753 (3)	0.0165 (13)	
C7	0.6706 (16)	0.9737 (6)	0.8255 (3)	0.0168 (13)	
H7	0.6588	1.0447	0.8105	0.020*	
C8	0.1574 (17)	0.8747 (6)	0.7254 (3)	0.0184 (13)	
C9	-0.0322 (15)	0.9270 (5)	0.6827 (3)	0.0166 (13)	
C10	-0.085 (3)	1.0323 (8)	0.6628 (8)	0.024 (3)	0.575 (18)
H10	0.0141	1.0934	0.6761	0.028*	0.575 (18)
C10'	-0.226 (5)	0.8865 (16)	0.6390 (12)	0.024*	0.425 (18)
H10'	-0.2631	0.8141	0.6297	0.028*	0.425 (18)
C11	-0.298 (3)	1.0318 (11)	0.6217 (7)	0.021 (3)	0.575 (18)
H11A	-0.3751	1.0920	0.6012	0.025*	0.575 (18)
C11'	-0.345 (4)	0.9727 (17)	0.6138 (9)	0.021*	0.425 (18)
H11'	-0.4821	0.9720	0.5819	0.025*	0.425 (18)
C12	-0.384 (3)	0.9257 (11)	0.6150 (5)	0.018 (3)	0.575 (18)
H12A	-0.5336	0.9012	0.5901	0.022*	0.575 (18)
C12'	-0.233 (4)	1.0650 (16)	0.6422 (9)	0.018*	0.425 (18)
H12'	-0.2832	1.1362	0.6325	0.022*	0.425 (18)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.0259 (3)	0.0187 (3)	0.0192 (3)	0.00161 (17)	-0.00272 (17)	-0.00437 (15)
O1	0.036 (3)	0.015 (2)	0.021 (3)	-0.005 (2)	-0.005 (2)	0.002 (2)
O2	0.022 (2)	0.013 (2)	0.026 (3)	0.002 (2)	-0.001 (2)	-0.0006 (19)
O3	0.023 (4)	0.018 (4)	0.006 (4)	0.004 (4)	-0.006 (3)	-0.004 (3)
O1W	0.027 (3)	0.013 (2)	0.039 (3)	-0.001 (2)	-0.002 (3)	-0.004 (2)
N1	0.023 (3)	0.017 (2)	0.012 (2)	0.002 (2)	0.001 (2)	-0.001 (2)
N2	0.022 (3)	0.010 (2)	0.015 (2)	0.001 (2)	-0.003 (2)	-0.001 (2)
C1	0.024 (4)	0.019 (3)	0.016 (3)	-0.002 (3)	0.001 (3)	0.000 (3)
C2	0.032 (4)	0.016 (3)	0.019 (3)	0.000 (3)	-0.004 (3)	0.003 (3)
C3	0.023 (3)	0.024 (4)	0.012 (3)	0.000 (3)	-0.004 (3)	0.003 (3)
C4	0.021 (3)	0.014 (3)	0.015 (3)	-0.002 (3)	0.001 (3)	-0.006 (2)
C5	0.026 (3)	0.014 (3)	0.012 (3)	0.001 (3)	0.000 (3)	-0.002 (2)
C6	0.016 (3)	0.017 (3)	0.016 (3)	0.003 (3)	0.000 (3)	0.001 (2)
C7	0.021 (3)	0.016 (3)	0.013 (3)	0.002 (3)	0.001 (3)	0.001 (2)
C8	0.020 (3)	0.020 (3)	0.015 (3)	0.003 (3)	0.001 (3)	-0.002 (3)
C9	0.021 (3)	0.014 (3)	0.015 (3)	0.003 (3)	0.001 (3)	-0.005 (2)
C10	0.031 (7)	0.019 (5)	0.020 (6)	0.000 (5)	-0.007 (6)	-0.003 (5)

C11	0.025 (6)	0.018 (6)	0.021 (6)	0.005 (5)	-0.002 (5)	0.005 (5)
C12	0.029 (6)	0.019 (6)	0.007 (4)	-0.003 (5)	-0.007 (4)	0.000 (4)

Geometric parameters (Å, °)

I1—C4	2.082 (7)	C3—H3	0.9500
O1—C1	1.342 (9)	C4—C5	1.379 (10)
O1—H1	0.8400	C5—C6	1.409 (11)
O2—C8	1.228 (9)	C5—H5	0.9500
O3—C9	1.363 (8)	C6—C7	1.451 (10)
O3—C12	1.363 (9)	C7—H7	0.9500
O3'—C9	1.355 (8)	C8—C9	1.449 (10)
O3'—C12'	1.358 (9)	C9—C10	1.413 (9)
O1W—H11	0.8402	C9—C10'	1.413 (10)
O1W—H12	0.8409	C10—C11	1.353 (9)
N1—C7	1.283 (10)	C10—H10	0.9500
N1—N2	1.381 (8)	C10'—C11'	1.341 (10)
N2—C8	1.365 (10)	C10'—H10'	0.9500
N2—H2	0.8800	C11—C12	1.407 (9)
C1—C2	1.406 (10)	C11—H11A	0.9500
C1—C6	1.418 (10)	C11'—C12'	1.415 (10)
C2—C3	1.402 (10)	C11'—H11'	0.9500
C2—H2A	0.9500	C12—H12A	0.9500
C3—C4	1.391 (9)	C12'—H12'	0.9500
C1—O1—H1	109.5	C6—C7—H7	118.6
C9—O3—C12	109.8 (8)	O2—C8—N2	123.1 (7)
C9—O3'—C12'	106.2 (12)	O2—C8—C9	122.5 (7)
H11—O1W—H12	109.0	N2—C8—C9	114.4 (6)
C7—N1—N2	114.4 (6)	O3—C9—C10	106.3 (8)
C8—N2—N1	120.4 (6)	O3'—C9—C10'	111.3 (11)
C8—N2—H2	119.8	O3'—C9—C8	116.8 (8)
N1—N2—H2	119.8	O3—C9—C8	117.1 (6)
O1—C1—C2	117.6 (6)	C10—C9—C8	136.6 (8)
O1—C1—C6	123.9 (7)	C10'—C9—C8	131.8 (10)
C2—C1—C6	118.5 (7)	C11—C10—C9	109.0 (9)
C3—C2—C1	120.9 (6)	C11—C10—H10	125.5
C3—C2—H2A	119.5	C9—C10—H10	125.5
C1—C2—H2A	119.5	C11'—C10'—C9	104.7 (14)
C4—C3—C2	119.5 (7)	C11'—C10'—H10'	127.6
C4—C3—H3	120.3	C9—C10'—H10'	127.6
C2—C3—H3	120.3	C10—C11—C12	107.2 (10)
C5—C4—C3	121.0 (6)	C10—C11—H11A	126.4
C5—C4—H1	118.7 (5)	C12—C11—H11A	126.4
C3—C4—H1	120.3 (5)	C10'—C11'—C12'	109.3 (16)
C4—C5—C6	120.2 (6)	C10'—C11'—H11'	125.3
C4—C5—H5	119.9	C12'—C11'—H11'	125.3
C6—C5—H5	119.9	O3—C12—C11	107.6 (9)
C5—C6—C1	119.9 (7)	O3—C12—H12A	126.2
C5—C6—C7	117.1 (6)	C11—C12—H12A	126.2

supplementary materials

C1—C6—C7	123.0 (7)	O3'—C12'—C11'	108.3 (15)
N1—C7—C6	122.7 (6)	O3'—C12'—H12'	125.8
N1—C7—H7	118.6	C11'—C12'—H12'	125.8
C7—N1—N2—C8	177.9 (6)	C12—O3—C9—C10'	-21 (6)
O1—C1—C2—C3	-178.5 (7)	C12—O3—C9—C8	177.4 (9)
C6—C1—C2—C3	1.0 (11)	O2—C8—C9—O3'	170.8 (9)
C1—C2—C3—C4	-1.8 (12)	N2—C8—C9—O3'	-9.3 (11)
C2—C3—C4—C5	0.6 (11)	O2—C8—C9—O3	1.1 (12)
C2—C3—C4—H1	-179.3 (6)	N2—C8—C9—O3	-179.0 (8)
C3—C4—C5—C6	1.3 (11)	O2—C8—C9—C10	-178.1 (12)
H1—C4—C5—C6	-178.8 (5)	N2—C8—C9—C10	1.8 (16)
C4—C5—C6—C1	-2.0 (11)	O2—C8—C9—C10'	-5(2)
C4—C5—C6—C7	176.9 (7)	N2—C8—C9—C10'	174.5 (19)
O1—C1—C6—C5	-179.6 (7)	O3'—C9—C10—C11	-151 (3)
C2—C1—C6—C5	0.8 (11)	O3—C9—C10—C11	1.9 (15)
O1—C1—C6—C7	1.6 (12)	C10'—C9—C10—C11	6.6 (18)
C2—C1—C6—C7	-178.0 (7)	C8—C9—C10—C11	-178.8 (10)
N2—N1—C7—C6	176.9 (6)	O3'—C9—C10'—C11'	3(3)
C5—C6—C7—N1	178.6 (7)	O3—C9—C10'—C11'	157 (8)
C1—C6—C7—N1	-2.6 (11)	C10—C9—C10'—C11'	-6(2)
N1—N2—C8—O2	-3.3 (11)	C8—C9—C10'—C11'	179.0 (13)
N1—N2—C8—C9	176.8 (6)	C9—C10—C11—C12	0.0 (17)
C12'—O3'—C9—O3	-10.9 (17)	C9—C10'—C11'—C12'	-1(3)
C12'—O3'—C9—C10	21 (2)	C9—O3—C12—C11	3.2 (14)
C12'—O3'—C9—C10'	-3(2)	C10—C11—C12—O3	-1.9 (16)
C12'—O3'—C9—C8	-179.6 (11)	C9—O3'—C12'—C11'	1.6 (19)
C12—O3—C9—O3'	8.7 (15)	C10'—C11'—C12'—O3'	0(3)
C12—O3—C9—C10	-3.1 (13)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2 \cdots O1w	0.88	1.96	2.813 (7)	163
O1—H1 \cdots N1	0.84	2.20	2.744 (8)	122
O1w—H11 \cdots O2 ⁱ	0.84	1.99	2.815 (8)	167
O1w—H12 \cdots O2 ⁱⁱ	0.84	2.00	2.826 (8)	168

Symmetry codes: (i) $-x+1, y+1/2, -z+3/2$; (ii) $-x, y+1/2, -z+3/2$.

Fig. 1




Fig. 2




Fig. 3

