

3-Chloro-4-(4-chlorophenoxy)-N-[(Z)-(5-nitrothiophen-2-yl)methylidene]aniline

Gonca Özdemir Tari* and Şamil Işık

Department of Physics, Faculty of Arts and Sciences, Ondokuz Mayıs University, Kurupelit, TR-55139 Samsun, Turkey

Correspondence e-mail: gozdemir@omu.edu.tr


Received 9 January 2012; accepted 11 January 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.050; wR factor = 0.124; data-to-parameter ratio = 14.8.

In the title compound, $\text{C}_{17}\text{H}_{10}\text{Cl}_2\text{N}_2\text{O}_3\text{S}$, the thiophene ring and the central benzene ring are almost coplanar [dihedral angle = $8.44(3)^\circ$], while the dihedral angle between the two benzene rings is $77.49(9)^\circ$. The crystal packing is stabilized by intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For background to the properties and uses of Schiff bases, see: Barton & Ollis (1979); Layer (1963); Ingold (1969); Cohen *et al.* (1964). For comparative bond lengths, see: Özdemir Tari *et al.* (2011); Kazak *et al.* (2000); Aygün *et al.* (1998).



Experimental

Crystal data

 $\text{C}_{17}\text{H}_{10}\text{Cl}_2\text{N}_2\text{O}_3\text{S}$ $M_r = 393.23$ Monoclinic, $P2_1/c$ $a = 16.3698(9)$ Å $b = 6.7787(2)$ Å $c = 15.9609(9)$ Å $\beta = 105.284(4)^\circ$ $V = 1708.47(14)$ Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.52$ mm⁻¹ $T = 293$ K $0.45 \times 0.30 \times 0.05$ mm

Data collection

Stoe IPDS II diffractometer

Absorption correction: integration

 $(X\text{-RED32}; \text{Stoe}, 2002)$ $T_{\min} = 0.817$, $T_{\max} = 0.942$

10905 measured reflections

3343 independent reflections

2317 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.062$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.050$ $wR(F^2) = 0.124$ $S = 1.01$

3343 reflections

226 parameters

H-atom parameters constrained

 $\Delta\rho_{\max} = 0.73$ e Å⁻³ $\Delta\rho_{\min} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C3}-\text{H3}\cdots\text{O2}^i$	0.93	2.38	3.278 (3)	162
$\text{C13}-\text{H13}\cdots\text{O1}^{ii}$	0.93	2.53	3.369 (4)	150

Symmetry codes: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *X-AREA* (Stoe, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996), *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDS II diffractometer (purchased under grant No. F279 of the University Research Fund).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5785).

References

- Aygün, M., Işık, Ş., Öcal, N., Tahir, M. N., Kaban, Ş. & Büyükgüngör, O. (1998). *Acta Cryst.* **C54**, 527–529.
- Barton, D. & Ollis, W. D. (1979). *Comprehensive Organic Chemistry*, Vol. 2. Oxford: Pergamon.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Cohen, M. D., Schmidt, G. M. J. & Flavian, S. (1964). *J. Chem. Soc.* pp. 1041–2051.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Ingold, C. K. (1969). *Structure and Mechanism in Organic Chemistry*, 2nd ed. Ithaca: Cornell University Press.
- Kazak, C., Aygün, M., Turgut, G., Odabaşoğlu, M., Özbey, S. & Büyükgüngör, O. (2000). *Acta Cryst.* **C56**, 1044–1045.
- Layer, R. W. (1963). *Chem. Rev.* **63**, 489–510.
- Özdemir Tari, G., Işık, Ş., Özkan, R. & Alaman Açar, A. (2011). *Acta Cryst.* **E67**, o343–o344.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Stoe (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2012). E68, o415 [doi:10.1107/S1600536812001316]

3-Chloro-4-(4-chlorophenoxy)-*N*-[(*Z*)-(5-nitrothiophen-2-yl)methylidene]aniline

G. Özdemir Tari and S. Isik

Comment

Schiff bases are used as starting materials in the synthesis of important drugs, such as antibiotics and antiallergic, antiphlogistic, and antitumor substances (Barton *et al.*, 1979; Layer, 1963; Ingold 1969). They show photochromism and thermochromism (Cohen *et al.*, 1964).

In this paper, the structure of the title compound, is reported. The N2=C13 bond length of 1.269 (3)Å is typical of a double bond, which is similar to the corresponding bond length in 2-[(4-Hydroxyphenyl)iminomethyl]-thiophene [1.282 (2) Å; Kazak *et al.*, 2000]; 3-methoxy-2-[(*E*)-(4-methoxyphenyl)imino)methyl]phenol [1.278 (3) Å, Özdemir Tari *et al.*, 2011] and *N*-(2,4-dinitrophenyl)-*N*-methylhydrozone [1.279 (2) Å, Aygün *et al.*, 1998].

The N2—C10 bond distance is 1.419 (3) Å, which is in good agreement with the corresponding bond lengths in 2-[(4-Hydroxyphenyl)iminomethyl]-thiophene [1.422 (2) Å, Kazak *et al.*, 2000] and 3-methoxy-2-[(*E*)-(4-methoxyphenyl)imino)methyl]phenol [1.419 (2) Å, Özdemir Tari *et al.*, 2011].

The C14—S1 and the C17—S1 distance are 1.713 (3)Å and 1.707 (2) Å, respectively. These distances are in good agreement with a related compound [1.712 (2)Å and 1.705 (3) Å; Kazak *et al.*, 2000]. The S··N2 distance is 3.036Å agree with similar length in related compound [3.135 (2) Å; Kazak *et al.*,2000].

Thermochromism or photochromism depends on the planarity or non-planarity of the molecule, respectively. The title compound might have photochromic properties because of the non-planarity of the molecule. The crystal structure is stabilized by intermolecular C—H··O hydrogen bonds.

Experimental

The compound (*Z*)—*N*-[3-chloro-4-(4-chlorophenoxy)phenyl]-1-(5-nitrothiophen-2-yl) methanimine was prepared by refluxing a mixture of a solution containing 5-nitro-2-thiophene-carboxaldehyde (0.077 g 0.049 mmol) in 20 ml ethanol and a solution containing 3-Chloro-4-(4-Chloro-phenoxy)aniline (0.012 g 0.049 mmol) in 20 ml ethanol. The reaction mixture was stirred for 1 h under reflux. The crystals of (*Z*)—*N*-[3-chloro-4-(4-chlorophenoxy)phenyl]-1-(5-nitrothiophen-2-yl)methanimine suitable for X-ray analysis were obtained from ethylalcohol by slow evaporation (yield % 78; m.p 402–404K).

Figures




Fig. 1. The molecular structure of the title compound, showing the atom-numbering scheme and 30% probability displacement ellipsoids.




Fig. 2. The crystal packing of the title compound. Intermolecular hydrogen bonds are shown as dashed lines.

3-Chloro-4-(4-chlorophenoxy)-N-[(Z)-(5-nitrothiophen-2-yl)methylidene]aniline

Crystal data

$C_{17}H_{10}Cl_2N_2O_3S$

$M_r = 393.23$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 16.3698$ (9) Å

$b = 6.7787$ (2) Å

$c = 15.9609$ (9) Å

$\beta = 105.284$ (4)°

$V = 1708.47$ (14) Å³

$Z = 4$

$F(000) = 800$

$D_x = 1.529$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 14032 reflections

$\theta = 1.3$ – 28.4 °

$\mu = 0.52$ mm⁻¹

$T = 293$ K

PRISM, brown

$0.45 \times 0.30 \times 0.05$ mm

Data collection

Stoe IPDS II
diffractometer

Radiation source: fine-focus sealed tube
graphite

Detector resolution: 6.67 pixels mm⁻¹

φ scan rotation method

Absorption correction: integration
(*X-RED32*; Stoe, 2002)

$T_{\min} = 0.817$, $T_{\max} = 0.942$

10905 measured reflections

3343 independent reflections

2317 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.062$

$\theta_{\max} = 26.0$ °, $\theta_{\min} = 2.6$ °

$h = -20 \rightarrow 20$

$k = -7 \rightarrow 8$

$l = -19 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.124$

$S = 1.01$

3343 reflections

226 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0716P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.73$ e Å⁻³

0 restraints

$$\Delta\rho_{\min} = -0.26 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.58285 (16)	-0.1486 (5)	0.16411 (18)	0.0613 (8)
C2	0.63253 (18)	-0.2004 (5)	0.2450 (2)	0.0652 (8)
H2	0.6545	-0.3273	0.2557	0.078*
C3	0.64914 (17)	-0.0589 (5)	0.31044 (18)	0.0605 (8)
H3	0.6824	-0.0910	0.3655	0.073*
C4	0.61642 (14)	0.1292 (4)	0.29387 (16)	0.0498 (6)
C5	0.56516 (16)	0.1787 (5)	0.21356 (17)	0.0566 (7)
H5	0.5422	0.3047	0.2032	0.068*
C6	0.54833 (17)	0.0374 (5)	0.14815 (18)	0.0622 (8)
H6	0.5137	0.0684	0.0935	0.075*
C7	0.69762 (15)	0.2549 (4)	0.42938 (16)	0.0468 (6)
C8	0.68181 (17)	0.2350 (5)	0.50843 (18)	0.0573 (7)
H8	0.6262	0.2267	0.5122	0.069*
C9	0.74793 (17)	0.2271 (5)	0.58380 (17)	0.0553 (7)
H9	0.7362	0.2155	0.6375	0.066*
C10	0.83122 (15)	0.2364 (4)	0.57930 (15)	0.0439 (6)
C11	0.84728 (16)	0.2591 (4)	0.49868 (16)	0.0484 (6)
H11	0.9028	0.2672	0.4946	0.058*
C12	0.78090 (15)	0.2698 (4)	0.42436 (16)	0.0493 (6)
C13	0.97201 (16)	0.2199 (4)	0.66068 (16)	0.0497 (6)
H13	0.9866	0.2123	0.6083	0.060*
C14	1.03864 (15)	0.2203 (4)	0.74123 (16)	0.0468 (6)
C15	1.12379 (17)	0.2157 (5)	0.74914 (18)	0.0576 (7)
H15	1.1471	0.2089	0.7020	0.069*
C16	1.17241 (17)	0.2224 (4)	0.83561 (18)	0.0553 (7)
H16	1.2313	0.2217	0.8532	0.066*
C17	1.12169 (16)	0.2299 (4)	0.88979 (17)	0.0471 (6)
N1	1.14919 (17)	0.2405 (4)	0.98236 (16)	0.0612 (6)
N2	0.89429 (13)	0.2296 (3)	0.65958 (13)	0.0461 (5)
O1	1.0953 (2)	0.2498 (4)	1.02221 (16)	0.0970 (9)
O2	1.22480 (16)	0.2380 (4)	1.01700 (15)	0.0842 (8)

supplementary materials

O3	0.63071 (11)	0.2755 (3)	0.35573 (12)	0.0565 (5)
S1	1.01578 (4)	0.22986 (10)	0.83989 (4)	0.0476 (2)
Cl1	0.56369 (6)	-0.32087 (18)	0.08067 (6)	0.0923 (3)
Cl2	0.80216 (5)	0.30835 (18)	0.32540 (4)	0.0828 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0445 (14)	0.081 (2)	0.0543 (16)	-0.0028 (14)	0.0062 (12)	-0.0082 (16)
C2	0.0536 (16)	0.071 (2)	0.0631 (18)	0.0057 (14)	0.0019 (13)	-0.0003 (16)
C3	0.0503 (15)	0.072 (2)	0.0481 (15)	0.0038 (13)	-0.0069 (12)	0.0053 (14)
C4	0.0382 (12)	0.0660 (18)	0.0418 (13)	-0.0006 (12)	0.0046 (10)	0.0002 (13)
C5	0.0464 (14)	0.072 (2)	0.0451 (14)	0.0038 (13)	0.0017 (11)	0.0075 (14)
C6	0.0513 (15)	0.084 (2)	0.0438 (14)	0.0023 (14)	0.0001 (11)	0.0061 (15)
C7	0.0418 (12)	0.0553 (17)	0.0380 (12)	0.0032 (11)	0.0009 (10)	0.0002 (11)
C8	0.0408 (13)	0.082 (2)	0.0475 (15)	-0.0002 (12)	0.0088 (11)	-0.0007 (14)
C9	0.0497 (14)	0.079 (2)	0.0362 (13)	-0.0028 (13)	0.0088 (11)	0.0001 (13)
C10	0.0464 (13)	0.0454 (15)	0.0352 (12)	-0.0013 (10)	0.0025 (10)	-0.0007 (10)
C11	0.0405 (12)	0.0636 (18)	0.0378 (13)	-0.0019 (11)	0.0045 (10)	-0.0012 (12)
C12	0.0454 (13)	0.0637 (18)	0.0360 (12)	0.0001 (11)	0.0057 (10)	0.0005 (12)
C13	0.0521 (15)	0.0562 (16)	0.0359 (12)	-0.0013 (11)	0.0029 (11)	-0.0003 (11)
C14	0.0468 (13)	0.0489 (15)	0.0397 (12)	-0.0028 (11)	0.0027 (10)	0.0019 (11)
C15	0.0486 (14)	0.077 (2)	0.0458 (14)	-0.0048 (13)	0.0092 (11)	0.0037 (14)
C16	0.0419 (13)	0.0652 (19)	0.0530 (15)	-0.0026 (12)	0.0020 (11)	0.0035 (13)
C17	0.0466 (13)	0.0444 (15)	0.0427 (13)	-0.0035 (10)	-0.0014 (11)	0.0010 (11)
N1	0.0716 (16)	0.0571 (16)	0.0431 (12)	-0.0040 (12)	-0.0055 (12)	-0.0017 (11)
N2	0.0469 (11)	0.0520 (13)	0.0326 (10)	-0.0034 (9)	-0.0013 (8)	-0.0006 (9)
O1	0.106 (2)	0.138 (3)	0.0482 (12)	0.0076 (16)	0.0216 (13)	-0.0073 (14)
O2	0.0833 (16)	0.0886 (18)	0.0569 (13)	-0.0110 (12)	-0.0239 (11)	-0.0014 (11)
O3	0.0443 (9)	0.0723 (14)	0.0440 (10)	0.0095 (8)	-0.0039 (7)	-0.0035 (9)
S1	0.0430 (3)	0.0539 (4)	0.0422 (3)	-0.0016 (3)	0.0049 (2)	-0.0008 (3)
Cl1	0.0864 (6)	0.1102 (8)	0.0710 (6)	0.0071 (5)	0.0042 (4)	-0.0307 (5)
Cl2	0.0535 (4)	0.1552 (9)	0.0373 (4)	-0.0077 (4)	0.0079 (3)	0.0114 (4)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.378 (5)	C10—C11	1.388 (4)
C1—C2	1.378 (4)	C10—N2	1.419 (3)
C1—Cl1	1.736 (3)	C11—C12	1.383 (3)
C2—C3	1.391 (4)	C11—H11	0.9300
C2—H2	0.9300	C12—Cl2	1.725 (3)
C3—C4	1.381 (4)	C13—N2	1.269 (3)
C3—H3	0.9300	C13—C14	1.450 (3)
C4—O3	1.375 (3)	C13—H13	0.9300
C4—C5	1.376 (4)	C14—C15	1.366 (4)
C5—C6	1.390 (4)	C14—S1	1.713 (3)
C5—H5	0.9300	C15—C16	1.400 (4)
C6—H6	0.9300	C15—H15	0.9300
C7—C8	1.360 (4)	C16—C17	1.348 (4)

C7—O3	1.387 (3)	C16—H16	0.9300
C7—C12	1.390 (4)	C17—N1	1.428 (3)
C8—C9	1.391 (4)	C17—S1	1.707 (2)
C8—H8	0.9300	N1—O2	1.215 (3)
C9—C10	1.386 (4)	N1—O1	1.218 (4)
C9—H9	0.9300		
C6—C1—C2	121.2 (3)	C11—C10—N2	124.8 (2)
C6—C1—C11	119.4 (2)	C12—C11—C10	120.2 (2)
C2—C1—C11	119.3 (3)	C12—C11—H11	119.9
C1—C2—C3	118.5 (3)	C10—C11—H11	119.9
C1—C2—H2	120.7	C11—C12—C7	120.5 (2)
C3—C2—H2	120.7	C11—C12—C12	119.4 (2)
C4—C3—C2	120.3 (3)	C7—C12—C12	120.10 (19)
C4—C3—H3	119.9	N2—C13—C14	121.9 (3)
C2—C3—H3	119.9	N2—C13—H13	119.0
O3—C4—C5	116.1 (3)	C14—C13—H13	119.0
O3—C4—C3	123.0 (2)	C15—C14—C13	126.3 (3)
C5—C4—C3	121.0 (3)	C15—C14—S1	112.38 (19)
C4—C5—C6	118.9 (3)	C13—C14—S1	121.3 (2)
C4—C5—H5	120.6	C14—C15—C16	113.0 (3)
C6—C5—H5	120.6	C14—C15—H15	123.5
C1—C6—C5	120.1 (3)	C16—C15—H15	123.5
C1—C6—H6	119.9	C17—C16—C15	110.3 (2)
C5—C6—H6	119.9	C17—C16—H16	124.8
C8—C7—O3	119.7 (2)	C15—C16—H16	124.8
C8—C7—C12	119.3 (2)	C16—C17—N1	125.9 (2)
O3—C7—C12	120.7 (2)	C16—C17—S1	115.0 (2)
C7—C8—C9	120.7 (2)	N1—C17—S1	119.1 (2)
C7—C8—H8	119.6	O2—N1—O1	123.7 (3)
C9—C8—H8	119.6	O2—N1—C17	118.4 (3)
C10—C9—C8	120.4 (2)	O1—N1—C17	118.0 (3)
C10—C9—H9	119.8	C13—N2—C10	120.2 (2)
C8—C9—H9	119.8	C4—O3—C7	118.9 (2)
C9—C10—C11	118.9 (2)	C17—S1—C14	89.30 (12)
C9—C10—N2	116.3 (2)		
C6—C1—C2—C3	-1.6 (5)	N2—C13—C14—C15	-178.1 (3)
C11—C1—C2—C3	178.2 (2)	N2—C13—C14—S1	1.1 (4)
C1—C2—C3—C4	-0.1 (5)	C13—C14—C15—C16	178.5 (3)
C2—C3—C4—O3	-179.9 (3)	S1—C14—C15—C16	-0.7 (3)
C2—C3—C4—C5	1.8 (4)	C14—C15—C16—C17	0.6 (4)
O3—C4—C5—C6	179.9 (2)	C15—C16—C17—N1	-179.2 (3)
C3—C4—C5—C6	-1.6 (4)	C15—C16—C17—S1	-0.1 (3)
C2—C1—C6—C5	1.8 (5)	C16—C17—N1—O2	-1.9 (4)
C11—C1—C6—C5	-178.1 (2)	S1—C17—N1—O2	179.1 (2)
C4—C5—C6—C1	-0.1 (4)	C16—C17—N1—O1	178.8 (3)
O3—C7—C8—C9	-175.8 (3)	S1—C17—N1—O1	-0.2 (4)
C12—C7—C8—C9	-0.8 (4)	C14—C13—N2—C10	177.6 (2)
C7—C8—C9—C10	-0.9 (5)	C9—C10—N2—C13	173.8 (2)

supplementary materials

C8—C9—C10—C11	1.7 (4)	C11—C10—N2—C13	-8.8 (4)
C8—C9—C10—N2	179.3 (3)	C5—C4—O3—C7	-163.4 (2)
C9—C10—C11—C12	-0.7 (4)	C3—C4—O3—C7	18.2 (4)
N2—C10—C11—C12	-178.1 (3)	C8—C7—O3—C4	-114.7 (3)
C10—C11—C12—C7	-1.0 (4)	C12—C7—O3—C4	70.5 (3)
C10—C11—C12—C12	177.3 (2)	C16—C17—S1—C14	-0.2 (2)
C8—C7—C12—C11	1.8 (4)	N1—C17—S1—C14	178.9 (2)
O3—C7—C12—C11	176.7 (2)	C15—C14—S1—C17	0.5 (2)
C8—C7—C12—C12	-176.5 (2)	C13—C14—S1—C17	-178.8 (2)
O3—C7—C12—C12	-1.6 (4)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3 \cdots O2 ⁱ	0.93	2.38	3.278 (3)	162.
C13—H13 \cdots O1 ⁱⁱ	0.93	2.53	3.369 (4)	150.

Symmetry codes: (i) $-x+2, y-1/2, -z+3/2$; (ii) $x, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

