

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(Thiophen-2-yl)-1-(thiophen-2-ylmethyl)-1H-benzimidazole

David K. Geiger,^{a*} H. Cristina Geiger,^a Leo Williams^a and Bruce C. Noll^b^aDepartment of Chemistry, State University of New York-College at Geneseo, 1 College Circle, Geneseo, NY 14454, USA, and ^bChemical Crystallography, Bruker AXS Inc., 5465 East Cheryl Parkway, Madison, Wisconsin, 53711, USA, and 7517 East Pass, Madison, Wisconsin, 53719, USA

Correspondence e-mail: geiger@geneseo.edu


Received 18 October 2011; accepted 21 December 2011

Key indicators: single-crystal X-ray study; $T = 300$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.040; wR factor = 0.103; data-to-parameter ratio = 12.0.

In the title compound, $\text{C}_{16}\text{H}_{12}\text{N}_2\text{S}_2$, the thiophene groups are rotationally disordered over two sets of sites, by approximately 180° , with occupancy ratios of 0.916 (2):0.084 (2) and 0.903 (2):0.097 (2). The major components of the thiophene and methylene substituted thiophene rings are canted by 24.06 (12) and 85.07 (10) $^\circ$, respectively, from the benzimidazole ring system plane and the dihedral angle between the major component thiophene ring planes is 84.90 (14) $^\circ$. In the crystal, there is a weak $\text{C}-\text{H}\cdots\text{N}$ hydrogen bond which links molecules into chains.

Related literature

For a discussion of the rearrangement of 1,2-diiminobenzene species to form benzimidazoles, see: Smith & Ho (1971). See Varala *et al.* (2007) for examples of proline-catalysed 1,2-disubstituted benzimidazole syntheses. Reich *et al.* (2004) provide examples of intermolecular aldimine coupling. For other syntheses of substituted benzimidazoles, see: Grimmett (1997); Bahrami *et al.* (2007); Du & Wang (2007). For the biological activity of benzimidazole derivatives, see: López-Rodríguez *et al.* (1999); Horton *et al.* (2003).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{12}\text{N}_2\text{S}_2$
 $M_r = 296.40$
 Monoclinic, $P2_1/n$
 $a = 8.9859$ (13) Å
 $b = 9.1601$ (11) Å
 $c = 17.476$ (3) Å
 $\beta = 93.629$ (5) $^\circ$
 $V = 1435.6$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.36$ mm⁻¹
 $T = 300$ K
 $0.80 \times 0.30 \times 0.10$ mm

Data collection

Bruker SMART X2S benchtop diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2008)
 $T_{\min} = 0.761$, $T_{\max} = 0.965$
 8857 measured reflections
 2527 independent reflections
 1942 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.103$
 $S = 1.05$
 2527 reflections
 210 parameters
 26 restraints
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.18$ e Å⁻³
 $\Delta\rho_{\min} = -0.25$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C16}-\text{H16}\cdots\text{N2}^i$	0.93	2.54	3.445 (4)	166

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2010); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XSELL (Bruker, 2004); software used to prepare material for publication: SHELXL97.

This work was supported by a Congressionally directed grant from the US Department of Education for the X-ray diffractometer and a grant from the Geneseo Foundation.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5359).

References

- Bahrami, K., Khodaei, M. M. & Naali, F. (2007). *J. Org. Chem.* **73**, 6835–6837.
 Bruker (2004). XSELL. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2008). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2009). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2010). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.
 Du, L.-H. & Wang, Y.-G. (2007). *Synthesis*, pp. 675–678.
 Grimmett, M. R. (1997). In *Imidazole and Benzimidazole Synthesis*. San Diego: Academic Press.
 Horton, D. A., Bourne, G. T. & Smythe, M. L. (2003). *Chem. Rev.* **103**, 893–930.
 López-Rodríguez, M. L., Benhamú, B., Morcillo, M. J., Tejada, I. D., Orensanz, L., Alfaro, M. J. & Martín, M. I. (1999). *J. Med. Chem.* **42**, 5020–5028.
 Reich, B. J. E., Justice, A. K., Beckstead, B. T., Reibenspies, J. H. & Miller, S. A. (2004). *J. Org. Chem.* **69**, 1357–1359.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Smith, J. G. & Ho, I. (1971). *Tetrahedron Lett.* **38**, 3541–3544.
 Varala, R., Nasreen, A., Enugala, R. & Adapa, S. R. (2007). *Tetrahedron Lett.* **48**, 69–72.

supplementary materials

Acta Cryst. (2012). E68, o420 [doi:10.1107/S1600536811055103]

2-(Thiophen-2-yl)-1-(thiophen-2-ylmethyl)-1*H*-benzimidazole

D. K. Geiger, H. C. Geiger, L. Williams and B. C. Noll

Comment

Benzimidazole derivatives are of interest because of their pharmacological uses. Examples include inhibitors of serotonin activated neurotransmission (López-Rodríguez *et al.*, 1999) and antiviral agents (Varala *et al.*, 2007). They have also found use as antiarrhythmic, antihistamine, antiulcer, anticancer, fungicidal, anthelmintical drugs (Horton *et al.*, 2003).

Numerous methods are available for the synthesis of substituted benzimidazoles (Grimmett, 1997). One pot procedures for the synthesis of 2-aryl substituted benzimidazoles involving the condensation of 1,2-diaminobenzene with aldehydes employing hydrogen peroxide and ceric ammonium nitrate (Bahrami *et al.*, 2007) or hypervalent iodine (Du & Wang, 2007) as oxidizing agents have been reported. Syntheses of 1-arylmethyl-2-aryl substituted benzimidazoles *via* the proline catalyzed condensation of 1,2-diaminobenzene derivatives with aryl aldehydes have also been reported (Varala *et al.*, 2007). Attempts to prepare *N,N'*-dibenzal-*o*-phenylenediamine *via* the condensation of 1,2-diaminobenzene and benzaldehyde lead to presumed rearrangement of the initially formed Schiff base yielding 1-benzyl-2-phenylbenzimidazole (Smith & Ho, 1971).

Our efforts have focused on the preparation of benzimidazole analogues which have substituents capable of binding metals. Toward that end, we have prepared 1-(thiophene-2-methyl)-2-(2-thiophene)benzimidazole, TMTB, from a reaction of 1,2-diaminobenzene with 2-thiophenecarboxaldehyde.

Figure 1 shows a perspective view of TMTB with the atom-labeling scheme. Only the major components of the disordered thiophene groups are displayed. Close C—H16···N2 intermolecular interactions (2.535 Å, 165.8°) result in chains of TMTB as shown in Figure 2. Bifurcated C—H11···C14 and C15 intermolecular contacts (2.843 Å and 2.848 Å) join the chains to form a layer network (see Figure 3). Figure 4 is a perspective view of the compound showing both major and minor components of the disordered thiophene groups.

The structure exhibits the expected planar benzimidazole moiety (maximum deviation 0.0076(0.0014) Å, N1). The major components of the disordered thiophene and methylthiophene rings are canted 24.06 (12) ° and 85.07 (10) °, respectively, from the benzimidazole plane, with a thiophene-methylthiophene dihedral angle of 84.90 (14) °. Together, the thiophene groups provide a structure with the potential to behave as a bidentate ligand employing the sulfur atoms. We are exploring the coordination chemistry of TMTB.

Experimental

The title compound was prepared by the reaction of two equivalents of 2-thiophene with 1,2-diaminobenzene in the presence of a catalytic amount of aluminium trichloride under nitrogen in dichloromethane. The reaction mixture was refluxed for 8 h, filtered and the solvent removed by rotary evaporation. The crude product was purified by column chromatography (silica gel) using 20%(v/v) ethylacetate in hexanes.

supplementary materials

The purified product was characterized *via* ^1H and ^{13}C NMR spectroscopies. ^1H NMR spectrum (DMSO- d_6 , 400 MHz, p.p.m.): 8.14 (2H, m), 7.98 (1 H, m), 7.50 (2 H, m), 7.44 (1 H, m), 7.39 (1 H, m), 7.10 (1 H, m), 6.94 (1 H, m), 6.05 (2 H, s). ^{13}C NMR spectrum (DMSO- d_6 , 100 MHz p.p.m.): 177.15, 150.63, 143.52, 138.59, 135.33, 133.05, 130.91, 129.49, 128.70, 128.01, 188.14, 117.51, 117.13, 115.37, 22.58.

Single crystals were grown *via* vapor diffusion of cyclohexane into a concentrated methanolic solution.

Refinement

During the course of the refinement, alternate positions for the thiophene sulfur atoms were apparent in the residual electron density maps. Refinement of the site occupancy factors led to values of about 8% for the minor components. Based on these values, a model was constructed for the minor component of each of the thiophene rings using the metrics of the major components as a guide. The pivot atoms (C8 and C13) were assumed to have full occupancy and so were not included in the disorder model. The minor four-atom components (S1', C11', C10' and C9; S2', C16', C15', and C14') were constrained to planarity using FLAT. Using *DFIX*, the S—C bond distances were set to 1.70 Å. Corresponding bond distances of the minor component and major component were set equal using SADI and corresponding thermal parameters were held the same using EADP. All atoms were refined anisotropically with hydrogen atoms in calculated positions using a riding model. With these constraints, the site occupancies of the major components of the methylthiophene and the thiophene refined to 91.6% and 90.3%, respectively. Based on this model, the angles between the mean planes of the major and minor components of the methylthiophene and thiophene are 5.7(1.9)° and 6.6 (9)°.

Figures




Fig. 1. Perspective view of the title compound with displacement ellipsoids of non-hydrogen atoms drawn at the 50% probability level. The disorder is not shown.




Fig. 2. View of the C—H...N intermolecular interactions leading to chain formation.




Fig. 3. View of layer formed by joining of chains by bifurcated C—H...C intermolecular contacts.




Fig. 4. The title molecule with displacement ellipsoids of non-hydrogen atoms drawn at the 50% probability level. Major and minor components of the disordered thiophene groups are shown. Hydrogen atoms have been omitted for clarity.

2-(Thiophen-2-yl)-1-(thiophen-2-ylmethyl)-1H-benzimidazole

Crystal data

$C_{16}H_{12}N_2S_2$

$M_r = 296.40$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1/n$

$a = 8.9859\ (13)\ \text{\AA}$

$b = 9.1601\ (11)\ \text{\AA}$

$c = 17.476\ (3)\ \text{\AA}$

$\beta = 93.629\ (5)^\circ$

$V = 1435.6\ (3)\ \text{\AA}^3$

$Z = 4$

$F(000) = 616$

$D_x = 1.371\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2556 reflections

$\theta = 2.3\text{--}24.0^\circ$

$\mu = 0.36\ \text{mm}^{-1}$

$T = 300\ \text{K}$

Plate, clear yellow

$0.80 \times 0.30 \times 0.10\ \text{mm}$

Data collection

Bruker SMART X2S benchtop diffractometer

Radiation source: fine-focus sealed tube graphite

ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2008)

$T_{\min} = 0.761$, $T_{\max} = 0.965$

8857 measured reflections

2527 independent reflections

1942 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.5^\circ$

$h = -10 \rightarrow 10$

$k = -10 \rightarrow 10$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.103$

$S = 1.05$

2527 reflections

210 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0518P)^2 + 0.2347P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.18\ \text{e \AA}^{-3}$

supplementary materials

26 restraints

$$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.21859 (8)	0.28769 (8)	0.08330 (3)	0.0467 (2)	0.916 (2)
C9	0.0913 (5)	0.3386 (5)	0.2054 (2)	0.0620 (12)	0.916 (2)
H9	0.0724	0.3488	0.2568	0.074*	0.916 (2)
C10	-0.0188 (4)	0.3558 (5)	0.1464 (2)	0.0648 (10)	0.916 (2)
H10	-0.1168	0.3819	0.1540	0.078*	0.916 (2)
C11	0.0332 (3)	0.3301 (5)	0.07718 (18)	0.0585 (9)	0.916 (2)
H11	-0.0251	0.3349	0.0313	0.070*	0.916 (2)
S1'	0.0789 (14)	0.3656 (15)	0.2202 (7)	0.0467 (2)	0.084 (2)
C9'	0.208 (3)	0.300 (4)	0.1075 (11)	0.0620 (12)	0.084 (2)
H9'	0.2835	0.2677	0.0773	0.074*	0.084 (2)
C11'	-0.017 (3)	0.384 (6)	0.1320 (15)	0.0585 (9)	0.084 (2)
H11'	-0.1150	0.4159	0.1246	0.070*	0.084 (2)
C10'	0.071 (3)	0.344 (6)	0.0747 (16)	0.0648 (10)	0.084 (2)
H10'	0.0434	0.3457	0.0225	0.078*	0.084 (2)
S2	0.18792 (12)	0.42615 (9)	0.47929 (5)	0.0647 (3)	0.903 (2)
C14	0.2441 (4)	0.1896 (4)	0.41297 (19)	0.0411 (8)	0.903 (2)
H14	0.2767	0.1211	0.3785	0.049*	0.903 (2)
C15	0.1692 (5)	0.1519 (4)	0.4778 (3)	0.0510 (8)	0.903 (2)
H15	0.1469	0.0563	0.4907	0.061*	0.903 (2)
C16	0.1329 (4)	0.2674 (5)	0.51922 (17)	0.0549 (10)	0.903 (2)
H16	0.0835	0.2618	0.5643	0.066*	0.903 (2)
S2'	0.2508 (19)	0.1572 (12)	0.3994 (8)	0.0647 (3)	0.097 (2)
C14'	0.187 (3)	0.394 (3)	0.4642 (13)	0.0411 (8)	0.097 (2)
H14'	0.1810	0.4923	0.4762	0.049*	0.097 (2)
C16'	0.146 (5)	0.146 (3)	0.478 (2)	0.0549 (10)	0.097 (2)
H16'	0.1140	0.0590	0.4989	0.066*	0.097 (2)
C15'	0.117 (5)	0.280 (4)	0.5038 (19)	0.0510 (8)	0.097 (2)
H15'	0.0564	0.2969	0.5442	0.061*	0.097 (2)
N1	0.41824 (18)	0.32345 (17)	0.29374 (9)	0.0385 (4)	
N2	0.47184 (19)	0.18925 (19)	0.19147 (9)	0.0453 (4)	
C1	0.5604 (2)	0.2661 (2)	0.31017 (12)	0.0405 (5)	

C2	0.5918 (2)	0.1839 (2)	0.24593 (12)	0.0450 (5)
C3	0.7282 (3)	0.1124 (3)	0.24430 (15)	0.0606 (7)
H3	0.749 (3)	0.062 (3)	0.2051 (14)	0.073*
C4	0.8285 (3)	0.1262 (3)	0.30724 (16)	0.0731 (8)
H4	0.9203	0.0793	0.3071	0.088*
C5	0.7950 (3)	0.2087 (3)	0.37082 (16)	0.0671 (7)
H5	0.8650	0.2161	0.4122	0.081*
C6	0.6600 (2)	0.2799 (3)	0.37371 (13)	0.0536 (6)
H6	0.6369	0.3345	0.4162	0.064*
C7	0.3719 (2)	0.2722 (2)	0.22212 (11)	0.0380 (5)
C8	0.2290 (2)	0.3058 (2)	0.18189 (11)	0.0398 (5)
C12	0.3420 (2)	0.4205 (2)	0.34473 (11)	0.0422 (5)
H12A	0.2690	0.4782	0.3148	0.051*
H12B	0.4139	0.4870	0.3695	0.051*
C13	0.2649 (2)	0.3368 (2)	0.40501 (10)	0.0374 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0584 (4)	0.0489 (4)	0.0324 (4)	0.0037 (3)	0.0007 (3)	-0.0022 (3)
C9	0.0552 (18)	0.079 (3)	0.053 (2)	0.0069 (17)	0.0129 (16)	-0.0182 (18)
C10	0.0447 (14)	0.078 (3)	0.071 (2)	0.0102 (15)	-0.0007 (15)	-0.0031 (18)
C11	0.0576 (19)	0.0612 (18)	0.0542 (16)	0.0000 (18)	-0.0167 (15)	-0.0041 (14)
S1'	0.0584 (4)	0.0489 (4)	0.0324 (4)	0.0037 (3)	0.0007 (3)	-0.0022 (3)
C9'	0.0552 (18)	0.079 (3)	0.053 (2)	0.0069 (17)	0.0129 (16)	-0.0182 (18)
C11'	0.0576 (19)	0.0612 (18)	0.0542 (16)	0.0000 (18)	-0.0167 (15)	-0.0041 (14)
C10'	0.0447 (14)	0.078 (3)	0.071 (2)	0.0102 (15)	-0.0007 (15)	-0.0031 (18)
S2	0.0900 (6)	0.0564 (6)	0.0515 (5)	-0.0007 (4)	0.0340 (4)	-0.0160 (3)
C14	0.0445 (14)	0.044 (2)	0.0355 (19)	0.0040 (14)	0.0106 (13)	-0.0070 (13)
C15	0.045 (2)	0.0523 (16)	0.0560 (16)	-0.0033 (12)	0.0097 (14)	0.0101 (13)
C16	0.0478 (17)	0.083 (2)	0.0345 (17)	0.0014 (15)	0.0113 (15)	0.0023 (15)
S2'	0.0900 (6)	0.0564 (6)	0.0515 (5)	-0.0007 (4)	0.0340 (4)	-0.0160 (3)
C14'	0.0445 (14)	0.044 (2)	0.0355 (19)	0.0040 (14)	0.0106 (13)	-0.0070 (13)
C16'	0.0478 (17)	0.083 (2)	0.0345 (17)	0.0014 (15)	0.0113 (15)	0.0023 (15)
C15'	0.045 (2)	0.0523 (16)	0.0560 (16)	-0.0033 (12)	0.0097 (14)	0.0101 (13)
N1	0.0465 (9)	0.0396 (9)	0.0300 (9)	-0.0002 (8)	0.0080 (7)	-0.0035 (7)
N2	0.0478 (10)	0.0536 (11)	0.0354 (10)	0.0049 (8)	0.0092 (8)	-0.0070 (8)
C1	0.0421 (11)	0.0424 (12)	0.0375 (12)	-0.0069 (9)	0.0068 (9)	0.0027 (9)
C2	0.0428 (11)	0.0526 (13)	0.0405 (12)	0.0011 (10)	0.0106 (10)	0.0035 (10)
C3	0.0491 (14)	0.0785 (18)	0.0558 (16)	0.0117 (13)	0.0157 (12)	-0.0021 (13)
C4	0.0437 (13)	0.100 (2)	0.076 (2)	0.0125 (14)	0.0076 (14)	0.0171 (17)
C5	0.0483 (14)	0.090 (2)	0.0617 (17)	-0.0087 (13)	-0.0084 (12)	0.0143 (15)
C6	0.0548 (14)	0.0645 (15)	0.0411 (13)	-0.0096 (11)	-0.0009 (11)	0.0010 (11)
C7	0.0458 (11)	0.0380 (11)	0.0309 (11)	-0.0025 (9)	0.0088 (9)	0.0011 (9)
C8	0.0463 (11)	0.0373 (11)	0.0360 (11)	-0.0022 (9)	0.0036 (9)	0.0019 (9)
C12	0.0575 (12)	0.0360 (11)	0.0341 (11)	0.0022 (9)	0.0108 (10)	-0.0061 (9)
C13	0.0398 (10)	0.0423 (12)	0.0303 (10)	0.0048 (9)	0.0046 (9)	-0.0055 (9)

supplementary materials

Geometric parameters (Å, °)

S1—C11	1.708 (3)	C14'—C13	1.388 (17)
S1—C8	1.728 (2)	C14'—C15'	1.42 (2)
C9—C8	1.362 (4)	C14'—H14'	0.9300
C9—C10	1.392 (5)	C16'—C15'	1.345 (19)
C9—H9	0.9300	C16'—H16'	0.9300
C10—C11	1.345 (4)	C15'—H15'	0.9300
C10—H10	0.9300	N1—C7	1.377 (2)
C11—H11	0.9300	N1—C1	1.394 (3)
S1'—C8	1.638 (10)	N1—C12	1.460 (2)
S1'—C11'	1.726 (14)	N2—C7	1.316 (2)
C9'—C8	1.303 (19)	N2—C2	1.393 (3)
C9'—C10'	1.39 (2)	C1—C6	1.387 (3)
C9'—H9'	0.9300	C1—C2	1.395 (3)
C11'—C10'	1.366 (19)	C2—C3	1.392 (3)
C11'—H11'	0.9300	C3—C4	1.383 (4)
C10'—H10'	0.9300	C3—H3	0.86 (2)
S2—C16	1.700 (4)	C4—C5	1.393 (4)
S2—C13	1.7166 (18)	C4—H4	0.9300
C14—C13	1.370 (4)	C5—C6	1.381 (3)
C14—C15	1.397 (4)	C5—H5	0.9300
C14—H14	0.9300	C6—H6	0.9300
C15—C16	1.334 (4)	C7—C8	1.457 (3)
C15—H15	0.9300	C12—C13	1.507 (3)
C16—H16	0.9300	C12—H12A	0.9700
S2'—C13	1.653 (10)	C12—H12B	0.9700
S2'—C16'	1.711 (15)		
C11—S1—C8	91.82 (13)	N1—C1—C2	105.49 (18)
C8—C9—C10	114.7 (3)	C3—C2—N2	130.3 (2)
C8—C9—H9	122.6	C3—C2—C1	119.6 (2)
C10—C9—H9	122.6	N2—C2—C1	110.09 (17)
C11—C10—C9	112.0 (3)	C4—C3—C2	118.1 (2)
C11—C10—H10	124.0	C4—C3—H3	121.3 (18)
C9—C10—H10	124.0	C2—C3—H3	120.6 (18)
C10—C11—S1	112.1 (2)	C3—C4—C5	121.4 (2)
C10—C11—H11	123.9	C3—C4—H4	119.3
S1—C11—H11	123.9	C5—C4—H4	119.3
C8—S1'—C11'	92.6 (11)	C6—C5—C4	121.3 (2)
C8—C9'—C10'	118 (2)	C6—C5—H5	119.3
C8—C9'—H9'	121.1	C4—C5—H5	119.3
C10'—C9'—H9'	121.1	C5—C6—C1	116.8 (2)
C10'—C11'—S1'	110.5 (16)	C5—C6—H6	121.6
C10'—C11'—H11'	124.7	C1—C6—H6	121.6
S1'—C11'—H11'	124.7	N2—C7—N1	113.07 (18)
C11'—C10'—C9'	108 (2)	N2—C7—C8	121.92 (18)
C11'—C10'—H10'	125.8	N1—C7—C8	125.00 (17)
C9'—C10'—H10'	125.8	C9'—C8—C9	103.6 (14)

C16—S2—C13	92.54 (14)	C9'—C8—C7	122.6 (13)
C13—C14—C15	113.6 (3)	C9—C8—C7	133.7 (2)
C13—C14—H14	123.2	C9'—C8—S1'	110.7 (13)
C15—C14—H14	123.2	C9—C8—S1'	10.0 (6)
C16—C15—C14	113.1 (3)	C7—C8—S1'	126.6 (5)
C16—C15—H15	123.5	C9'—C8—S1	6.0 (13)
C14—C15—H15	123.5	C9—C8—S1	109.2 (2)
C15—C16—S2	111.6 (2)	C7—C8—S1	116.84 (14)
C15—C16—H16	124.2	S1'—C8—S1	116.5 (5)
S2—C16—H16	124.2	N1—C12—C13	111.81 (16)
C13—S2'—C16'	93.3 (11)	N1—C12—H12A	109.3
C13—C14'—C15'	110.4 (19)	C13—C12—H12A	109.3
C13—C14'—H14'	124.8	N1—C12—H12B	109.3
C15'—C14'—H14'	124.8	C13—C12—H12B	109.3
C15'—C16'—S2'	110.1 (16)	H12A—C12—H12B	107.9
C15'—C16'—H16'	124.9	C14—C13—C14'	102.4 (12)
S2'—C16'—H16'	124.9	C14—C13—C12	130.06 (19)
C16'—C15'—C14'	114 (2)	C14'—C13—C12	127.4 (11)
C16'—C15'—H15'	123.1	C14—C13—S2'	10.1 (5)
C14'—C15'—H15'	123.1	C14'—C13—S2'	112.2 (12)
C7—N1—C1	106.23 (16)	C12—C13—S2'	120.1 (4)
C7—N1—C12	129.49 (17)	C14—C13—S2	109.18 (17)
C1—N1—C12	124.27 (17)	C14'—C13—S2	8.3 (11)
C7—N2—C2	105.12 (16)	C12—C13—S2	120.76 (15)
C6—C1—N1	131.82 (19)	S2'—C13—S2	119.1 (4)
C6—C1—C2	122.7 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C16—H16 \cdots N2 ⁱ	0.93	2.54	3.445 (4)	166

Symmetry codes: (i) $x-1/2, -y+1/2, z+1/2$.

Fig. 1




Fig. 2




Fig. 3

