

3-Acetyl-1-phenylthiourea

Durre Shahwar,^a M. Nawaz Tahir,^{b*} Muhammad Mansha Chohan,^a Naeem Ahmad^a and Samiullah^a

^aDepartment of Chemistry, Government College University, Lahore, Pakistan, and

^bUniversity of Sargodha, Department of Physics, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 15 January 2012; accepted 19 January 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.045; wR factor = 0.129; data-to-parameter ratio = 20.5.

In the crystal structure of title compound, $\text{C}_9\text{H}_{10}\text{N}_2\text{OS}$, there are two symmetry-independent molecules, each having an intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond generating an $S(6)$ ring motif. The benzene rings and the virtually planar acetylthiourea fragments [r.m.s. deviations = 0.0045 and 0.0341 Å] are oriented at dihedral angles of 50.71 (6) and 62.79 (6)° in the two molecules. In the crystal, $\text{N}-\text{H}\cdots\text{S}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link molecules *via* cyclic $R_2^2(8)$ and $R_2^2(12)$ motifs into a one-dimensional polymeric network extending along [101]. The intra- and intermolecular $\text{N}-\text{H}\cdots\text{O}$ interactions are part of a three-center hydrogen bond. A $\text{C}-\text{H}\cdots\text{S}$ interaction also occurs.

Related literature

For related structures, see: Othman *et al.* (2010). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_9\text{H}_{10}\text{N}_2\text{OS}$
 $M_r = 194.25$
 Monoclinic, $P2_1/c$
 $a = 10.1911$ (2) Å

$b = 22.5480$ (4) Å
 $c = 8.9736$ (2) Å
 $\beta = 112.449$ (1)°
 $V = 1905.77$ (7) Å³

$Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.30$ mm⁻¹

$T = 296$ K
 $0.35 \times 0.25 \times 0.22$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.915$, $T_{\max} = 0.938$

16643 measured reflections
 4679 independent reflections
 3626 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.129$
 $S = 1.05$
 4679 reflections

228 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.30$ e Å⁻³
 $\Delta\rho_{\min} = -0.35$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}$	0.86	1.97	2.662 (2)	136
$\text{N1}-\text{H1}\cdots\text{O2}^{\text{i}}$	0.86	2.46	3.1967 (19)	143
$\text{N2}-\text{H2A}\cdots\text{S2}^{\text{ii}}$	0.86	2.64	3.4931 (17)	170
$\text{N3}-\text{H3A}\cdots\text{O2}$	0.86	1.97	2.6633 (7)	137
$\text{N3}-\text{H3A}\cdots\text{O1}^{\text{i}}$	0.86	2.42	3.1418 (18)	142
$\text{N4}-\text{H4A}\cdots\text{S1}^{\text{ii}}$	0.86	2.57	3.4150 (18)	168
$\text{C18}-\text{H18B}\cdots\text{S1}^{\text{ii}}$	0.96	2.83	3.594 (3)	137

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x, -y, -z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of the diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan. They also acknowledge the technical support provided by Syed Muhammad Hussain Rizvi of Bana International, Karachi, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2451).

References

- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555-1573.
 Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837-838.
 Othman, E. A., Soh, S. K. C. & Yamin, B. M. (2010). *Acta Cryst.* **E66**, o628.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148-155.

supplementary materials

Acta Cryst. (2012). E68, o508 [doi:10.1107/S1600536812002371]

3-Acetyl-1-phenylthiourea

D. Shahwar, M. N. Tahir, M. M. Chohan, N. Ahmad and Samiullah

Comment

The title compound **I** (Fig. 1) has been synthesized in search of new enzyme inhibitors.

The crystal structure of *N*-(3-chloropropionyl)-*N'*-phenylthiourea (Othman *et al.*, 2010) has been published which is related to the title compound (I).

In (I), two molecules in the asymmetric unit are present, which differ from each other geometrically. In one molecule, the benzene ring A (C1–C6) and the acetylthiourea moiety B (N1/C7/S1/N2/C8/O1/C9) are planar with r. m. s. deviation of 0.0045 Å and 0.0341 Å, respectively. The dihedral angle between A/B is 50.71 (6)°. In second molecule, the benzene ring C (C10–C15) and the acetylthiourea moiety D (N3/C16/S2/N4/C17/O2/C18) are also planar with r. m. s. deviation of 0.0037 Å and 0.0453 Å, respectively and the dihedral angle between C/D is 62.79 (6)°. In both molecules, there exist classical intramolecular H-bonding of N—H···O type (Table 1, Fig. 1) with *S*(6) ring motif (Bernstein *et al.*, 1995). Both molecules are interlinked due to strong N—H···O, N—H···S and C—H···S types of H-bondings (Table 1, Fig. 2). The *S*(6) ring motifs of both molecules are connected into a four membered ring (—O···H···O···H···O—). The N—H···S type of bonding completes $R_2^2(8)$ ring motif. $R_2^1(6)$ ring motif is formed due to C—H···S and N—H···S types of intermolecular H-bondings (Fig. 2). The molecules are linked to form of one-dimensional polymeric chains extending along [101] (Fig. 2).

Experimental

The title compound was synthesized by adding 0.1 mol (7.13 ml) of acetyl chloride dropwise to a stirred solution of KSCN (0.11 mol) in dry acetone (50 ml), followed by slow addition of aniline (0.1 mol) in dry acetone (25 ml). The mixture was refluxed for 5–10 min, then poured on ice cooled water, which resulted in crude precipitate. Recrystallization of the precipitate in from ethyl acetate yielded light green prisms (m.p. 365 K).

Refinement

The H atoms were positioned geometrically (C—H = 0.93–0.96 Å, N—H = 0.86 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$, where $x = 1.5$ for methyl groups and $x = 1.2$ for other H atoms.

Figures




Fig. 1. View of the title compound with the displacement ellipsoids drawn at the 50% probability level. H atoms are shown by small circles of arbitrary radii. The dotted lines represent the intramolecular H-bondings.




Fig. 2. The partial packing (*PLATON*; Spek, 2009) which shows that molecules form polymeric chains extending along [1 0 1] direction.

3-Acetyl-1-phenylthiourea

Crystal data

$C_9H_{10}N_2OS$

$M_r = 194.25$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 10.1911$ (2) Å

$b = 22.5480$ (4) Å

$c = 8.9736$ (2) Å

$\beta = 112.449$ (1)°

$V = 1905.77$ (7) Å³

$Z = 8$

$F(000) = 816$

$D_x = 1.354$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3626 reflections

$\theta = 2.2$ – 28.3 °

$\mu = 0.30$ mm⁻¹

$T = 296$ K

Prism, light green

$0.35 \times 0.25 \times 0.22$ mm

Data collection

Bruker Kappa APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

Detector resolution: 8.00 pixels mm⁻¹

ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)

$T_{\min} = 0.915$, $T_{\max} = 0.938$

16643 measured reflections

4679 independent reflections

3626 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.024$

$\theta_{\max} = 28.3$ °, $\theta_{\min} = 2.2$ °

$h = -13 \rightarrow 13$

$k = -23 \rightarrow 29$

$l = -11 \rightarrow 11$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.129$

$S = 1.05$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0574P)^2 + 0.6731P]$

where $P = (F_o^2 + 2F_c^2)/3$

4679 reflections	$(\Delta/\sigma)_{\max} < 0.001$
228 parameters	$\Delta\rho_{\max} = 0.30 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	-0.19196 (6)	0.05015 (3)	0.17599 (8)	0.0666 (2)
O1	0.22186 (16)	-0.05280 (6)	0.3471 (2)	0.0647 (5)
N1	0.08613 (17)	0.04741 (6)	0.35398 (19)	0.0455 (5)
N2	-0.01041 (17)	-0.03499 (6)	0.19702 (19)	0.0463 (5)
C1	0.09330 (18)	0.10632 (7)	0.4158 (2)	0.0388 (5)
C2	0.1603 (2)	0.11483 (8)	0.5792 (2)	0.0455 (6)
C3	0.1800 (2)	0.17188 (9)	0.6419 (3)	0.0548 (7)
C4	0.1317 (2)	0.21994 (8)	0.5415 (3)	0.0546 (7)
C5	0.0635 (3)	0.21096 (8)	0.3790 (3)	0.0575 (7)
C6	0.0449 (2)	0.15463 (8)	0.3146 (2)	0.0532 (7)
C7	-0.0283 (2)	0.02154 (8)	0.2479 (2)	0.0428 (5)
C8	0.1098 (2)	-0.06975 (8)	0.2494 (2)	0.0449 (6)
C9	0.08992 (12)	-0.13043 (2)	0.17563 (12)	0.0596 (7)
S2	0.29361 (6)	0.11538 (2)	0.07535 (6)	0.0571 (2)
O2	0.59666 (10)	-0.02218 (2)	0.39879 (7)	0.0624 (5)
N3	0.52470 (6)	0.09140 (2)	0.33705 (7)	0.0456 (5)
N4	0.41578 (16)	0.01207 (6)	0.1764 (2)	0.0463 (5)
C10	0.5529 (2)	0.15166 (7)	0.3895 (2)	0.0419 (5)
C11	0.4580 (2)	0.18411 (10)	0.4315 (3)	0.0569 (7)
C12	0.4945 (3)	0.24100 (10)	0.4927 (3)	0.0692 (9)
C13	0.6232 (3)	0.26489 (9)	0.5078 (3)	0.0669 (8)
C14	0.7165 (3)	0.23256 (10)	0.4653 (3)	0.0641 (8)
C15	0.6826 (2)	0.17540 (9)	0.4056 (3)	0.0523 (6)
C16	0.41954 (19)	0.07247 (7)	0.2070 (2)	0.0405 (5)
C17	0.5029 (2)	-0.03202 (8)	0.2695 (2)	0.0459 (6)
C18	0.4715 (2)	-0.09258 (9)	0.1962 (3)	0.0590 (7)
H1	0.16310	0.02690	0.38876	0.0545*
H2	0.19244	0.08241	0.64760	0.0546*
H2A	-0.08394	-0.05025	0.12321	0.0555*
H3	0.22598	0.17763	0.75237	0.0658*

supplementary materials

H4	0.14526	0.25818	0.58346	0.0655*
H5	0.02914	0.24337	0.31114	0.0690*
H6	0.00015	0.14915	0.20379	0.0639*
H9A	0.15836	-0.15710	0.24765	0.0894*
H9B	0.10276	-0.12869	0.07516	0.0894*
H9C	-0.00410	-0.14434	0.15685	0.0894*
H3A	0.58190	0.06497	0.39608	0.0548*
H4A	0.35054	0.00054	0.08761	0.0556*
H11	0.37031	0.16814	0.41908	0.0682*
H12	0.43186	0.26299	0.52351	0.0830*
H13	0.64664	0.30324	0.54715	0.0803*
H14	0.80349	0.24893	0.47639	0.0769*
H15	0.74639	0.15337	0.37671	0.0627*
H18A	0.52816	-0.09968	0.13370	0.0886*
H18B	0.37273	-0.09519	0.12774	0.0886*
H18C	0.49338	-0.12174	0.28022	0.0886*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0461 (3)	0.0524 (3)	0.0799 (4)	0.0084 (2)	0.0003 (3)	-0.0220 (3)
O1	0.0495 (8)	0.0449 (8)	0.0774 (10)	0.0053 (6)	-0.0008 (8)	-0.0177 (7)
N1	0.0430 (8)	0.0309 (8)	0.0525 (9)	0.0012 (6)	0.0071 (7)	-0.0056 (6)
N2	0.0436 (8)	0.0339 (8)	0.0500 (9)	-0.0021 (6)	0.0053 (7)	-0.0088 (6)
C1	0.0400 (9)	0.0290 (8)	0.0459 (9)	-0.0014 (7)	0.0147 (8)	-0.0031 (7)
C2	0.0501 (10)	0.0370 (9)	0.0446 (10)	0.0048 (8)	0.0128 (8)	0.0013 (7)
C3	0.0629 (12)	0.0491 (11)	0.0465 (11)	0.0021 (9)	0.0143 (9)	-0.0117 (9)
C4	0.0709 (13)	0.0340 (10)	0.0654 (13)	-0.0038 (9)	0.0332 (11)	-0.0104 (9)
C5	0.0835 (15)	0.0314 (10)	0.0602 (12)	0.0039 (9)	0.0302 (12)	0.0064 (8)
C6	0.0739 (14)	0.0386 (10)	0.0422 (10)	0.0001 (9)	0.0166 (10)	0.0025 (8)
C7	0.0470 (10)	0.0336 (9)	0.0431 (9)	-0.0001 (7)	0.0121 (8)	-0.0024 (7)
C8	0.0472 (10)	0.0342 (9)	0.0475 (10)	0.0001 (7)	0.0115 (8)	-0.0036 (7)
C9	0.0599 (12)	0.0392 (10)	0.0691 (14)	0.0009 (9)	0.0128 (11)	-0.0143 (10)
S2	0.0581 (3)	0.0369 (3)	0.0535 (3)	0.0050 (2)	-0.0041 (2)	0.0012 (2)
O2	0.0595 (9)	0.0436 (8)	0.0599 (9)	0.0107 (7)	-0.0041 (7)	-0.0056 (7)
N3	0.0441 (8)	0.0323 (7)	0.0481 (9)	0.0042 (6)	0.0037 (7)	-0.0030 (6)
N4	0.0443 (8)	0.0328 (8)	0.0485 (9)	-0.0012 (6)	0.0029 (7)	-0.0049 (6)
C10	0.0478 (10)	0.0309 (8)	0.0381 (9)	0.0018 (7)	0.0063 (8)	-0.0013 (7)
C11	0.0540 (12)	0.0503 (12)	0.0620 (13)	0.0037 (9)	0.0174 (10)	-0.0091 (9)
C12	0.0836 (17)	0.0504 (13)	0.0657 (14)	0.0180 (12)	0.0197 (13)	-0.0105 (10)
C13	0.0972 (19)	0.0329 (10)	0.0563 (13)	-0.0030 (11)	0.0134 (12)	-0.0048 (9)
C14	0.0773 (15)	0.0446 (12)	0.0647 (14)	-0.0174 (11)	0.0206 (12)	-0.0023 (10)
C15	0.0573 (12)	0.0424 (10)	0.0543 (11)	-0.0020 (9)	0.0182 (10)	-0.0015 (8)
C16	0.0394 (9)	0.0337 (9)	0.0443 (9)	-0.0012 (7)	0.0114 (8)	-0.0004 (7)
C17	0.0412 (9)	0.0357 (9)	0.0532 (11)	0.0031 (7)	0.0095 (8)	-0.0017 (8)
C18	0.0563 (12)	0.0348 (10)	0.0703 (14)	0.0060 (9)	0.0067 (10)	-0.0063 (9)

Geometric parameters (Å, °)

S1—C7	1.671 (2)	C2—H2	0.9300
S2—C16	1.6796 (18)	C3—H3	0.9300
O1—C8	1.206 (3)	C4—H4	0.9300
O2—C17	1.2090 (19)	C5—H5	0.9300
N1—C7	1.326 (2)	C6—H6	0.9300
N1—C1	1.431 (2)	C9—H9A	0.9600
N2—C7	1.389 (2)	C9—H9B	0.9600
N2—C8	1.377 (3)	C9—H9C	0.9600
N1—H1	0.8600	C10—C15	1.382 (3)
N2—H2A	0.8600	C10—C11	1.375 (3)
N3—C10	1.4307 (17)	C11—C12	1.389 (3)
N3—C16	1.3181 (18)	C12—C13	1.376 (4)
N4—C16	1.387 (2)	C13—C14	1.363 (4)
N4—C17	1.381 (2)	C14—C15	1.388 (3)
N3—H3A	0.8600	C17—C18	1.496 (3)
N4—H4A	0.8600	C11—H11	0.9300
C1—C6	1.383 (2)	C12—H12	0.9300
C1—C2	1.374 (2)	C13—H13	0.9300
C2—C3	1.388 (3)	C14—H14	0.9300
C3—C4	1.375 (3)	C15—H15	0.9300
C4—C5	1.370 (4)	C18—H18A	0.9600
C5—C6	1.378 (3)	C18—H18B	0.9600
C8—C9	1.4996 (19)	C18—H18C	0.9600
C1—N1—C7	125.98 (17)	H9A—C9—H9C	109.00
C7—N2—C8	128.41 (16)	C8—C9—H9A	109.00
C7—N1—H1	117.00	H9B—C9—H9C	109.00
C1—N1—H1	117.00	C8—C9—H9C	109.00
C8—N2—H2A	116.00	H9A—C9—H9B	109.00
C7—N2—H2A	116.00	C8—C9—H9B	109.00
C10—N3—C16	126.30 (11)	C11—C10—C15	120.67 (17)
C16—N4—C17	128.62 (16)	N3—C10—C11	121.43 (18)
C16—N3—H3A	117.00	N3—C10—C15	117.80 (17)
C10—N3—H3A	117.00	C10—C11—C12	119.3 (2)
C17—N4—H4A	116.00	C11—C12—C13	120.1 (2)
C16—N4—H4A	116.00	C12—C13—C14	120.2 (2)
C2—C1—C6	119.87 (15)	C13—C14—C15	120.5 (3)
N1—C1—C2	118.32 (15)	C10—C15—C14	119.2 (2)
N1—C1—C6	121.61 (15)	S2—C16—N3	125.50 (11)
C1—C2—C3	119.88 (17)	S2—C16—N4	118.07 (13)
C2—C3—C4	120.3 (2)	N3—C16—N4	116.42 (14)
C3—C4—C5	119.36 (19)	O2—C17—C18	123.18 (17)
C4—C5—C6	121.01 (18)	N4—C17—C18	114.25 (16)
C1—C6—C5	119.54 (17)	O2—C17—N4	122.57 (16)
N1—C7—N2	116.73 (18)	C10—C11—H11	120.00
S1—C7—N2	117.66 (14)	C12—C11—H11	120.00
S1—C7—N1	125.58 (14)	C11—C12—H12	120.00

supplementary materials

O1—C8—N2	122.67 (17)	C13—C12—H12	120.00
N2—C8—C9	114.58 (15)	C12—C13—H13	120.00
O1—C8—C9	122.74 (17)	C14—C13—H13	120.00
C3—C2—H2	120.00	C13—C14—H14	120.00
C1—C2—H2	120.00	C15—C14—H14	120.00
C4—C3—H3	120.00	C10—C15—H15	120.00
C2—C3—H3	120.00	C14—C15—H15	120.00
C3—C4—H4	120.00	C17—C18—H18A	109.00
C5—C4—H4	120.00	C17—C18—H18B	109.00
C6—C5—H5	120.00	C17—C18—H18C	109.00
C4—C5—H5	119.00	H18A—C18—H18B	109.00
C1—C6—H6	120.00	H18A—C18—H18C	109.00
C5—C6—H6	120.00	H18B—C18—H18C	109.00
C7—N1—C1—C2	133.6 (2)	C6—C1—C2—C3	-0.4 (3)
C7—N1—C1—C6	-51.6 (3)	N1—C1—C6—C5	-175.2 (2)
C1—N1—C7—S1	-5.7 (3)	C2—C1—C6—C5	-0.5 (3)
C1—N1—C7—N2	176.57 (16)	N1—C1—C2—C3	174.48 (19)
C8—N2—C7—S1	-173.67 (16)	C1—C2—C3—C4	0.6 (3)
C8—N2—C7—N1	4.2 (3)	C2—C3—C4—C5	0.2 (4)
C7—N2—C8—O1	-3.3 (3)	C3—C4—C5—C6	-1.2 (4)
C7—N2—C8—C9	176.44 (16)	C4—C5—C6—C1	1.3 (4)
C10—N3—C16—S2	-1.0 (2)	N3—C10—C11—C12	175.39 (18)
C16—N3—C10—C11	62.9 (2)	C15—C10—C11—C12	-0.9 (3)
C16—N3—C10—C15	-120.69 (19)	N3—C10—C15—C14	-176.23 (18)
C10—N3—C16—N4	177.67 (15)	C11—C10—C15—C14	0.2 (3)
C16—N4—C17—C18	-177.93 (19)	C10—C11—C12—C13	1.3 (4)
C17—N4—C16—S2	-176.84 (17)	C11—C12—C13—C14	-1.1 (4)
C17—N4—C16—N3	4.4 (3)	C12—C13—C14—C15	0.4 (4)
C16—N4—C17—O2	1.8 (3)	C13—C14—C15—C10	0.1 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O1	0.86	1.97	2.662 (2)	136
N1—H1 \cdots O2 ⁱ	0.86	2.46	3.1967 (19)	143
N2—H2A \cdots S2 ⁱⁱ	0.86	2.64	3.4931 (17)	170
N3—H3A \cdots O2	0.86	1.97	2.6633 (7)	137
N3—H3A \cdots O1 ⁱ	0.86	2.42	3.1418 (18)	142
N4—H4A \cdots S1 ⁱⁱ	0.86	2.57	3.4150 (18)	168
C18—H18B \cdots S1 ⁱⁱ	0.96	2.83	3.594 (3)	137

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x, -y, -z$.

Fig. 1




Fig. 2

