

2-Chloro-*N*-(3-methylphenyl)benzamideVinola Z. Rodrigues,^a B. Thimme Gowda,^{a*} Viktor Vrábel^b and Jozef Kožíšek^b^aDepartment of Chemistry, Mangalore University, Mangalagangothri-574 199, Mangalore, India, and ^bInstitute of Physical Chemistry and Chemical Physics, Slovak University of Technology, Radlinského 9, SK-812 37 Bratislava, Slovak Republic
Correspondence e-mail: gowdabt@yahoo.com


Received 7 February 2012; accepted 9 February 2012

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.031; wR factor = 0.069; data-to-parameter ratio = 16.2.

In the structure of the title compound, $\text{C}_{14}\text{H}_{12}\text{ClNO}$, the *ortho*-Cl atom in the benzoyl ring is positioned *syn* to the $\text{C}=\text{O}$ bond, while the *meta*-methyl group in the aniline ring is positioned *anti* to the $\text{N}-\text{H}$ bond. The amide group forms dihedral angles of 60.1 (1) and 22.0 (1)°, respectively, with the benzoyl and aniline rings, while the angle between these rings is 38.7 (1)°. The crystal structure is stabilized by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, which give rise to infinite chains running along the c axis.

Related literature

For studies, including ours, on the effects of substituents on the structures and other aspects of *N*-(aryl)-amides, see: Bowes *et al.* (2003); Gowda *et al.* (1999, 2006); Rodrigues *et al.* (2011); Saeed *et al.* (2010); for *N*-(aryl)-methanesulfonamides, see: Gowda *et al.* (2007); for *N*-chloroarylamides, see: Jyothi & Gowda (2004); and for *N*-bromoarylsulfonamides, see: Usha & Gowda (2006).



Experimental

Crystal data

 $\text{C}_{14}\text{H}_{12}\text{ClNO}$
 $M_r = 245.70$
Tetragonal, $P4_3$
 $a = 8.8751$ (3) Å
 $c = 15.9642$ (5) Å
 $V = 1257.45$ (6) Å³ $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.29$ mm⁻¹
 $T = 295$ K
 $0.4 \times 0.3 \times 0.2$ mm

Data collection

Oxford Diffraction Xcalibur System diffractometer
Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2009)
 $T_{\min} = 0.898$, $T_{\max} = 0.942$
8244 measured reflections
2552 independent reflections
1757 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.069$
 $S = 1.04$
2552 reflections
158 parameters
2 restraintsH atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.08$ e Å⁻³
 $\Delta\rho_{\min} = -0.11$ e Å⁻³
Absolute structure: Flack (1983), 1229 Friedel pairs
Flack parameter: 0.00 (6)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^i$	0.86 (1)	2.03 (1)	2.8790 (16)	171 (2)

Symmetry code: (i) $y, -x + 1, z + \frac{1}{4}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2002); software used to prepare material for publication: *SHELXL97*, *PLATON* (Spek, 2009) and *WinGX* (Farrugia, 1999).

VZR thanks the University Grants Commission, Government of India, New Delhi, for the award of an RFSMS research fellowship. VV and JK thank the Grant Agencies for their financial support (VEGA Grant Agency of the Slovak Ministry of Education, grant No. 1/0679/11), the Research and Development Agency of Slovakia (grant No. APVV-0202-10) and Structural Funds, Interreg IIIA, for financial support in purchasing the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5196).

References

- Bowes, K. F., Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2003). *Acta Cryst.* **C59**, o1–o3.
Brandenburg, K. (2002). *DIAMOND*. Crystal Impact GbH, Bonn, Germany.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Gowda, B. T., Bhat, D. K., Fuess, H. & Weiss, A. (1999). *Z. Naturforsch. Teil A*, **54**, 261–267.
Gowda, B. T., Foro, S. & Fuess, H. (2007). *Acta Cryst.* **E63**, o2597.
Gowda, B. T., Kozisek, J. & Fuess, H. (2006). *Z. Naturforsch. Teil A*, **61**, 588–594.
Jyothi, K. & Gowda, B. T. (2004). *Z. Naturforsch. Teil A*, **59**, 64–68.
Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
Rodrigues, V. Z., Kucková, L., Gowda, B. T. & Kožíšek, J. (2011). *Acta Cryst.* **E67**, o3277.
Saeed, A., Arshad, M. & Simpson, J. (2010). *Acta Cryst.* **E66**, o2808–o2809.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Usha, K. M. & Gowda, B. T. (2006). *J. Chem. Sci.* **118**, 351–359.

supplementary materials

Acta Cryst. (2012). E68, o723 [doi:10.1107/S1600536812005739]

2-Chloro-*N*-(3-methylphenyl)benzamide

Vinola Z. Rodrigues, B. Thimme Gowda, Viktor Vrábel and Jozef Kožíšek

Comment

The amide and sulfonamide moieties are the constituents of many biologically important compounds. As part of our studies on the substituent effects on the structures and other aspects of *N*-(aryl)-amides (Bowes *et al.*, 2003; Gowda *et al.*, 1999, 2006; Rodrigues *et al.*, 2011; Saeed *et al.*, 2010), *N*-(aryl)-methanesulfonamides (Gowda *et al.*, 2007), *N*-chloro-arylsulfonamides (Jyothi & Gowda, 2004) and *N*-bromoarylsulfonamides (Usha & Gowda, 2006), in the present work, the crystal structure of 2-chloro-*N*-(3-methylphenyl)benzamide (I) has been determined (Fig. 1).

In (I), the *ortho*-Cl atom in the benzoyl ring is positioned *syn* to the C=O bond, while the *meta*-methyl group in the anilino ring is positioned *anti* to the N—H bond, similar to that observed in 3-chloro-*N*-(3-methylphenyl)benzamide (I) (Rodrigues *et al.*, 2011).

The amide group forms dihedral angles of 60.1 (1) and 22.0 (1)°, respectively, with the benzoyl and aniline rings, while the angle between the benzoyl and aniline rings is 38.7 (1)°, compared to the value of 77.4 (1)° in (II).

In the crystal structure, intermolecular N1—H1⋯O1 hydrogen bonds (Table 1) link the molecules into infinite chains running along the *c*-axis. Part of the crystal structure is shown in Fig. 2.

Experimental

The title compound was prepared by a method similar to the one described by Gowda *et al.* (2006). The purity of the compound was checked by determining its melting point. It was characterized by recording its infrared and NMR spectra.

Plate like colorless single crystals of the title compound used in the X-ray diffraction studies were obtained by slow evaporation of an ethanol solution of the compound (0.5 g in about 30 ml of ethanol) at room temperature.

Refinement

All hydrogen atoms bound to carbon were placed in calculated positions with C—H distances of 0.93 Å (C-aromatic), 0.96 Å (C-methyl) and constrained to ride on their parent atoms. The amide H atom was located in a difference map and refined with the N—H distance restrained to 0.86 (1) Å. $U_{\text{iso}}(\text{H})$ values were set at 1.2 $U_{\text{eq}}(\text{C-aromatic, N})$ and 1.5 $U_{\text{eq}}(\text{C-methyl})$.

Computing details

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis CCD* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED* (Oxford Diffraction, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2002); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008), *PLATON* (Spek, 2009) and *WinGX* (Farrugia, 1999).





Figure 1

Molecular structure of the title compound showing the atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small spheres of arbitrary radius.

**Figure 2**

Packing view of the title compound. Molecular chains along along the *c* axis are generated by N—H...O hydrogen bonds which are shown as dashed lines. H atoms have been omitted.

2-Chloro-*N*-(3-methylphenyl)benzamide*Crystal data* $C_{14}H_{12}ClNO$ $M_r = 245.70$ Tetragonal, $P4_3$ $a = 8.8751(3) \text{ \AA}$ $c = 15.9642(5) \text{ \AA}$ $V = 1257.45(6) \text{ \AA}^3$ $Z = 4$ $F(000) = 512$ $D_x = 1.298 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2552 reflections

 $\theta = 3.4\text{--}29.3^\circ$ $\mu = 0.29 \text{ mm}^{-1}$ $T = 295 \text{ K}$

Plate, colourless

 $0.4 \times 0.3 \times 0.2 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur System diffractometer	8244 measured reflections 2552 independent reflections
Radiation source: fine-focus sealed tube	1757 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.019$
Detector resolution: 0 pixels mm^{-1}	$\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 4.1^\circ$
ω scans with κ offsets	$h = -10 \rightarrow 11$
Absorption correction: multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2009)	$k = -11 \rightarrow 10$
$T_{\text{min}} = 0.898$, $T_{\text{max}} = 0.942$	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.031$	$w = 1/[\sigma^2(F_o^2) + (0.0348P)^2]$
$wR(F^2) = 0.069$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2552 reflections	$\Delta\rho_{\text{max}} = 0.08 \text{ e } \text{\AA}^{-3}$
158 parameters	$\Delta\rho_{\text{min}} = -0.11 \text{ e } \text{\AA}^{-3}$
2 restraints	Absolute structure: Flack (1983), ???? Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.00 (6)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6060 (2)	0.6255 (2)	0.21616 (9)	0.0579 (4)
C2	0.6965 (2)	0.7340 (2)	0.25296 (11)	0.0731 (5)
C3	0.6388 (4)	0.8363 (2)	0.30991 (13)	0.1015 (8)
H3	0.7002	0.9103	0.3330	0.122*
C4	0.4900 (5)	0.8268 (3)	0.33167 (17)	0.1176 (10)
H4	0.4505	0.8952	0.3700	0.141*
C5	0.3986 (3)	0.7188 (4)	0.29813 (18)	0.1124 (9)
H5	0.2979	0.7131	0.3140	0.135*
C6	0.4562 (3)	0.6180 (2)	0.24059 (14)	0.0819 (6)
H6	0.3939	0.5442	0.2180	0.098*
C7	0.66413 (19)	0.52431 (19)	0.14825 (9)	0.0525 (4)
C8	0.69202 (18)	0.25396 (19)	0.11204 (9)	0.0529 (4)
C9	0.7955 (2)	0.2664 (2)	0.04811 (10)	0.0582 (4)
H9	0.8393	0.3594	0.0368	0.070*
C10	0.8357 (2)	0.1411 (2)	-0.00002 (10)	0.0678 (5)

C11	0.7689 (3)	0.0059 (3)	0.01807 (13)	0.0839 (6)
H11	0.7941	-0.0786	-0.0134	0.101*
C12	0.6663 (3)	-0.0075 (2)	0.08115 (15)	0.0927 (7)
H12	0.6228	-0.1007	0.0923	0.111*
C13	0.6264 (2)	0.1167 (2)	0.12884 (12)	0.0732 (5)
H13	0.5560	0.1075	0.1717	0.088*
C14	0.9504 (3)	0.1563 (3)	-0.06817 (13)	0.0986 (7)
H14A	0.9645	0.0605	-0.0950	0.148*
H14B	1.0442	0.1896	-0.0447	0.148*
H14C	0.9160	0.2285	-0.1086	0.148*
N1	0.65297 (16)	0.37678 (15)	0.16483 (7)	0.0541 (3)
H1	0.6216 (16)	0.3554 (18)	0.2143 (4)	0.057 (5)*
O1	0.71348 (15)	0.57708 (12)	0.08283 (6)	0.0695 (3)
Cl1	0.88575 (8)	0.74156 (9)	0.22927 (4)	0.1238 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0768 (13)	0.0532 (10)	0.0435 (9)	0.0070 (9)	0.0009 (8)	0.0051 (8)
C2	0.1011 (15)	0.0678 (12)	0.0505 (10)	-0.0026 (11)	0.0069 (10)	-0.0031 (9)
C3	0.167 (3)	0.0738 (15)	0.0636 (13)	-0.0023 (15)	0.0112 (15)	-0.0163 (12)
C4	0.184 (3)	0.0806 (18)	0.0882 (17)	0.052 (2)	0.024 (2)	-0.0110 (15)
C5	0.106 (2)	0.122 (2)	0.1091 (19)	0.0456 (19)	0.0298 (16)	-0.0028 (18)
C6	0.0827 (15)	0.0798 (14)	0.0831 (13)	0.0154 (11)	0.0071 (12)	-0.0038 (12)
C7	0.0604 (10)	0.0576 (11)	0.0397 (8)	0.0039 (8)	-0.0057 (7)	0.0000 (8)
C8	0.0618 (11)	0.0561 (11)	0.0407 (8)	0.0074 (9)	-0.0086 (8)	0.0017 (7)
C9	0.0647 (11)	0.0597 (11)	0.0503 (9)	0.0074 (8)	-0.0004 (8)	-0.0018 (8)
C10	0.0762 (13)	0.0770 (14)	0.0504 (10)	0.0228 (11)	-0.0066 (9)	-0.0098 (9)
C11	0.1158 (18)	0.0646 (14)	0.0714 (13)	0.0147 (12)	-0.0142 (13)	-0.0211 (10)
C12	0.134 (2)	0.0550 (13)	0.0897 (15)	-0.0091 (12)	-0.0063 (15)	-0.0062 (12)
C13	0.0945 (15)	0.0604 (13)	0.0647 (11)	-0.0058 (10)	0.0061 (10)	0.0001 (10)
C14	0.1021 (18)	0.118 (2)	0.0757 (13)	0.0372 (14)	0.0136 (12)	-0.0137 (13)
N1	0.0747 (10)	0.0516 (9)	0.0361 (7)	0.0009 (7)	0.0059 (6)	0.0023 (6)
O1	0.1070 (10)	0.0619 (8)	0.0397 (6)	-0.0015 (6)	0.0060 (6)	0.0053 (6)
Cl1	0.1097 (5)	0.1673 (7)	0.0943 (4)	-0.0532 (4)	0.0089 (3)	-0.0407 (4)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.384 (3)	C8—C9	1.378 (2)
C1—C6	1.387 (3)	C8—N1	1.421 (2)
C1—C7	1.499 (2)	C9—C10	1.398 (2)
C2—C3	1.383 (3)	C9—H9	0.9300
C2—Cl1	1.723 (2)	C10—C11	1.369 (3)
C3—C4	1.368 (4)	C10—C14	1.496 (3)
C3—H3	0.9300	C11—C12	1.363 (3)
C4—C5	1.366 (4)	C11—H11	0.9300
C4—H4	0.9300	C12—C13	1.386 (3)
C5—C6	1.381 (3)	C12—H12	0.9300
C5—H5	0.9300	C13—H13	0.9300
C6—H6	0.9300	C14—H14A	0.9600

C7—O1	1.2254 (19)	C14—H14B	0.9600
C7—N1	1.340 (2)	C14—H14C	0.9600
C8—C13	1.376 (2)	N1—H1	0.859 (2)
C2—C1—C6	118.06 (17)	C8—C9—C10	120.92 (17)
C2—C1—C7	121.60 (16)	C8—C9—H9	119.5
C6—C1—C7	120.25 (17)	C10—C9—H9	119.5
C3—C2—C1	121.4 (2)	C11—C10—C9	118.07 (18)
C3—C2—C11	118.66 (19)	C11—C10—C14	121.81 (18)
C1—C2—C11	119.96 (14)	C9—C10—C14	120.12 (19)
C4—C3—C2	119.0 (2)	C12—C11—C10	121.44 (18)
C4—C3—H3	120.5	C12—C11—H11	119.3
C2—C3—H3	120.5	C10—C11—H11	119.3
C5—C4—C3	121.1 (2)	C11—C12—C13	120.5 (2)
C5—C4—H4	119.4	C11—C12—H12	119.7
C3—C4—H4	119.4	C13—C12—H12	119.7
C4—C5—C6	119.7 (3)	C8—C13—C12	119.23 (18)
C4—C5—H5	120.2	C8—C13—H13	120.4
C6—C5—H5	120.2	C12—C13—H13	120.4
C5—C6—C1	120.7 (2)	C10—C14—H14A	109.5
C5—C6—H6	119.6	C10—C14—H14B	109.5
C1—C6—H6	119.6	H14A—C14—H14B	109.5
O1—C7—N1	124.64 (14)	C10—C14—H14C	109.5
O1—C7—C1	120.67 (15)	H14A—C14—H14C	109.5
N1—C7—C1	114.67 (14)	H14B—C14—H14C	109.5
C13—C8—C9	119.83 (16)	C7—N1—C8	127.93 (13)
C13—C8—N1	117.39 (15)	C7—N1—H1	115.0 (11)
C9—C8—N1	122.74 (16)	C8—N1—H1	117.1 (11)
C6—C1—C2—C3	-2.6 (3)	C13—C8—C9—C10	-0.3 (2)
C7—C1—C2—C3	173.88 (17)	N1—C8—C9—C10	177.13 (14)
C6—C1—C2—C11	176.38 (15)	C8—C9—C10—C11	0.2 (2)
C7—C1—C2—C11	-7.1 (2)	C8—C9—C10—C14	-178.91 (17)
C1—C2—C3—C4	1.8 (3)	C9—C10—C11—C12	-0.2 (3)
C11—C2—C3—C4	-177.27 (19)	C14—C10—C11—C12	178.9 (2)
C2—C3—C4—C5	-0.1 (4)	C10—C11—C12—C13	0.2 (3)
C3—C4—C5—C6	-0.7 (4)	C9—C8—C13—C12	0.3 (3)
C4—C5—C6—C1	-0.3 (4)	N1—C8—C13—C12	-177.23 (16)
C2—C1—C6—C5	1.9 (3)	C11—C12—C13—C8	-0.3 (3)
C7—C1—C6—C5	-174.69 (19)	O1—C7—N1—C8	-1.0 (3)
C2—C1—C7—O1	-58.9 (2)	C1—C7—N1—C8	177.16 (15)
C6—C1—C7—O1	117.55 (19)	C13—C8—N1—C7	-158.57 (16)
C2—C1—C7—N1	122.85 (17)	C9—C8—N1—C7	24.0 (2)
C6—C1—C7—N1	-60.7 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
---------------	-------	-------------	-------------	---------------

N1—H1 \cdots O1 ⁱ	0.86 (1)	2.03 (1)	2.8790 (16)	171 (2)
--------------------------------	----------	----------	-------------	---------

Symmetry code: (i) $y, -x+1, z+1/4$.