

trans-Dichloridobis(triphenylphosphane- κP)palladium(II) benzene hemisolvate

Frank Meyer-Wegner, Hans-Wolfram Lerner, Tanja Sinke and Michael Bolte*

Institut für Anorganische und Analytische Chemie, Goethe-Universität Frankfurt,
Max-von-Laue-Strasse 7, 60438 Frankfurt am Main, Germany
Correspondence e-mail: bolte@chemie.uni-frankfurt.de


Received 21 March 2012; accepted 22 March 2012

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$;
 R factor = 0.041; wR factor = 0.095; data-to-parameter ratio = 15.9.

The title complex, $[\text{PdCl}_2(\text{C}_{18}\text{H}_{15}\text{P})_2] \cdot 0.5\text{C}_6\text{H}_6$, has the Pd^{II} ion in a square-planar coordination mode (r.m.s. deviation for Pd , P and Cl atoms = 0.024 Å) with the PPh_3 and Cl ligands mutually *trans*. The benzene solvent molecule is located about a crystallographic inversion centre. The title complex is isostructural with *trans*-dichloridobis(triphenylphosphane)-palladium(II) 1,4-dichlorobenzene sesquisolvate [Kitano *et al.* (1983). *Acta Cryst. C* **39**, 1015–1017].

Related literature

For the synthetic background, see: Lerner (2005); Meyer-Wegner *et al.* (2009, 2011). For *trans*-dichlorido-bis(triphenylphosphane)palladium(II) sesqui(*p*-dichlorobenzene), see: Kitano *et al.* (1983).



Experimental

Crystal data

$[\text{PdCl}_2(\text{C}_{18}\text{H}_{15}\text{P})_2] \cdot 0.5\text{C}_6\text{H}_6$	$V = 3361.8 (3)\text{ \AA}^3$
$M_r = 740.89$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 11.4530 (7)\text{ \AA}$	$\mu = 0.83\text{ mm}^{-1}$
$b = 18.4493 (8)\text{ \AA}$	$T = 173\text{ K}$
$c = 16.4696 (10)\text{ \AA}$	$0.26 \times 0.08 \times 0.08\text{ mm}$
$\beta = 104.979 (5)^{\circ}$	

Data collection

Stoe IPDS II two-circle diffractometer	42186 measured reflections
Absorption correction: multi-scan (<i>X-AREA</i> ; Stoe & Cie, 2001)	6302 independent reflections
$T_{\min} = 0.812$, $T_{\max} = 0.936$	5104 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.089$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	4 restraints
$wR(F^2) = 0.095$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.45\text{ e \AA}^{-3}$
6302 reflections	$\Delta\rho_{\text{min}} = -0.76\text{ e \AA}^{-3}$
397 parameters	

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

This work was supported by the Beilstein Institute as part of the NanoBiC research cooperative (project eNet).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5074).

References

- Kitano, Y., Kinoshita, Y., Nakamura, R. & Ashida, T. (1983). *Acta Cryst. C* **39**, 1015–1017.
- Lerner, H.-W. (2005). *Coord. Chem. Rev.* **249**, 781–798.
- Meyer-Wegner, F., Nadj, A., Bolte, M., Auner, N., Wagner, M., Holthausen, M. C. & Lerner, H.-W. (2011). *Chem. Eur. J.* **17**, 4715–4719.
- Meyer-Wegner, F., Scholz, S., Sänger, I., Schödel, F., Bolte, M., Wagner, M. & Lerner, H.-W. (2009). *Organometallics*, **28**, 6835–6837.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Stoe & Cie (2001). *X-AREA*. Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2012). E68, m507 [doi:10.1107/S1600536812012494]

***trans*-Dichloridobis(triphenylphosphane- κP)palladium(II) benzene hemisolvate**

Frank Meyer-Wegner, Hans-Wolfram Lerner, Tanja Sinke and Michael Bolte

Comment

Recently, we have shown that the degradation of $\text{Cl}_3\text{SiSiCl}_3$ in the presence of donors, such as amines NR_3 ($R = \text{Me}, \text{Et}$) and phosphanes PR_3 ($R = t\text{Bu}$) and the isoelectronic silanide $[\text{Si}t\text{Bu}_3]^-$ (Lerner, 2005), in the first step gives dichlorsilylene SiCl_2 and ultimately produces the perchlorinated neopentasilane $\text{Si}(\text{SiCl}_3)_4$ and in the case of the silanide the silatetrahedrane $(t\text{Bu}_3\text{Si})_4\text{Si}_4$, respectively (Meyer-Wegner *et al.*, 2009, 2011). Moreover, we verified that the donor-induced degradation of $\text{Cl}_3\text{SiSiCl}_3$ in the presence of the silylene trapping agent 2,3-dimethyl-1,3-butadiene gives the [4 + 1] cycloadduct (Meyer-Wegner *et al.*, 2011). We are currently interested in dichlorsilylene transition metal complexes. To this end, we thought that such complexes can be prepared using the reaction between $\text{Cl}_3\text{SiSiCl}_3$ and $[\text{Pd}(\text{PPh}_3)_4]$. However, the reaction of $\text{Cl}_3\text{SiSiCl}_3$ with $[\text{Pd}(\text{PPh}_3)_4]$ gives exclusively $[\text{PdCl}_2(\text{PPh}_3)_2]$ (Fig. 1). Apparently no dichlorsilylene transition metal complex was formed thereby. Single crystals composed of one molecule of benzene and one molecule of $[\text{PdCl}_2(\text{PPh}_3)_2]$ could be isolated from the reaction solution.

The title complex (Fig. 2) has the Pd centre in a quadratic planar (r.m.s. deviation for Pd, P and Cl atoms: 0.024 Å) coordination mode with the PPh_3 and Cl ligands mutually *trans*. The solvent benzene molecule is located about a crystallographic inversion centre. The title compound is isomorphous with *trans*-dichlorido-bis(triphenyl-phosphane)palladium(II) sesqui(*p*-dichlorobenzene) (Kitano *et al.*, 1983). The packing diagram (Fig. 3) shows how the benzene molecules fill the empty space between the complexes. There are no unusual features.

Experimental


$\text{Pd}(\text{PPh}_3)_4$ (0.19 g, 0.16 mmol) was suspended in benzene (5 ml) and then treated with one equivalent of Si_2Cl_6 (0.028 ml, 44 mg, 0.16 mmol) in benzene (0.5 ml) at ambient temperature. The solution immediately became clear and turned from yellow to dark red. Crystals of the title compound were obtained by slow evaporation of the solvent at ambient temperature.

Refinement


H atoms were refined using a riding model, with C—H = 0.95 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The C—C distances in the solvent benzene molecule were restrained to 1.39 (1) Å.

Computing details


Data collection: *X-Area* (Stoe & Cie, 2001); cell refinement: *X-Area* (Stoe & Cie, 2001); data reduction: *X-Area* (Stoe & Cie, 2001); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

**Figure 1**

Reaction scheme.

**Figure 2**

A perspective view of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms and the benzene molecule are omitted for clarity.

**Figure 3**

Packing diagram of the title compound. H atoms omitted for clarity.

***trans*-Dichloridobis(triphenylphosphane- κP)palladium(II) benzene hemisolvate**

Crystal data



$M_r = 740.89$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 11.4530 (7)$ Å

$b = 18.4493 (8)$ Å

$c = 16.4696 (10)$ Å

$\beta = 104.979 (5)^\circ$

$V = 3361.8 (3)$ Å³

$Z = 4$

$F(000) = 1508$

$D_x = 1.464 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 28523 reflections

$\theta = 3.3\text{--}26.1^\circ$

$\mu = 0.83 \text{ mm}^{-1}$

$T = 173$ K

Rod, yellow

$0.26 \times 0.08 \times 0.08$ mm

Data collection

Stoe IPDS II two-circle

diffractometer

Radiation source: Genix 3D $I\mu S$ microfocus X-ray source

Genix 3D multilayer optics monochromator

ω scans

Absorption correction: multi-scan
(*X-AREA*; Stoe & Cie, 2001)

$T_{\min} = 0.812$, $T_{\max} = 0.936$

42186 measured reflections

6302 independent reflections

5104 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.089$

$\theta_{\max} = 25.7^\circ$, $\theta_{\min} = 3.3^\circ$

$h = -13 \rightarrow 13$

$k = -22 \rightarrow 22$

$l = -20 \rightarrow 19$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.041$$

$$wR(F^2) = 0.095$$

$$S = 1.03$$

6302 reflections

397 parameters

4 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0537P)^2]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.45 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.76 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Pd1	0.45706 (2)	0.831821 (14)	0.206242 (15)	0.01813 (9)
Cl1	0.42899 (9)	0.70941 (5)	0.19156 (6)	0.0353 (2)
Cl2	0.48343 (8)	0.95572 (5)	0.21552 (6)	0.0292 (2)
P1	0.25111 (7)	0.83867 (5)	0.20060 (5)	0.01825 (18)
P2	0.65875 (7)	0.81933 (5)	0.20311 (5)	0.01921 (18)
C11	0.1927 (3)	0.75972 (18)	0.2434 (2)	0.0210 (7)
C12	0.2494 (3)	0.7383 (2)	0.3252 (2)	0.0255 (7)
H12	0.3193	0.7632	0.3561	0.031*
C13	0.2041 (4)	0.6805 (2)	0.3619 (2)	0.0320 (9)
H13	0.2417	0.6669	0.4182	0.038*
C14	0.1042 (4)	0.6429 (2)	0.3163 (3)	0.0363 (9)
H14	0.0729	0.6035	0.3413	0.044*
C15	0.0500 (4)	0.6625 (2)	0.2346 (3)	0.0339 (8)
H15	-0.0171	0.6357	0.2027	0.041*
C16	0.0930 (3)	0.7213 (2)	0.1988 (2)	0.0284 (8)
H16	0.0536	0.7353	0.1430	0.034*
C21	0.1665 (3)	0.84101 (19)	0.0905 (2)	0.0221 (7)
C22	0.2071 (4)	0.8002 (2)	0.0328 (2)	0.0307 (8)
H22	0.2799	0.7733	0.0505	0.037*
C23	0.1417 (4)	0.7986 (2)	-0.0508 (2)	0.0387 (10)
H23	0.1690	0.7695	-0.0898	0.046*
C24	0.0377 (4)	0.8389 (2)	-0.0778 (2)	0.0363 (9)
H24	-0.0066	0.8378	-0.1351	0.044*
C25	-0.0017 (4)	0.8810 (3)	-0.0206 (2)	0.0375 (10)
H25	-0.0724	0.9098	-0.0392	0.045*
C26	0.0608 (3)	0.8817 (2)	0.0633 (2)	0.0288 (8)

H26	0.0319	0.9097	0.1024	0.035*
C31	0.1919 (3)	0.91423 (19)	0.2491 (2)	0.0208 (7)
C32	0.1352 (3)	0.9029 (2)	0.3138 (2)	0.0276 (8)
H32	0.1272	0.8553	0.3335	0.033*
C33	0.0908 (4)	0.9614 (2)	0.3490 (3)	0.0348 (9)
H33	0.0523	0.9535	0.3929	0.042*
C34	0.1013 (4)	1.0303 (2)	0.3217 (2)	0.0330 (9)
H34	0.0702	1.0699	0.3464	0.040*
C35	0.1580 (3)	1.0425 (2)	0.2574 (2)	0.0282 (8)
H35	0.1657	1.0905	0.2385	0.034*
C36	0.2029 (3)	0.98475 (19)	0.2212 (2)	0.0231 (7)
H36	0.2413	0.9931	0.1773	0.028*
C41	0.7013 (3)	0.73483 (19)	0.1600 (2)	0.0246 (7)
C42	0.6964 (4)	0.6693 (2)	0.2010 (3)	0.0361 (9)
H42	0.6730	0.6691	0.2523	0.043*
C43	0.7250 (4)	0.6044 (2)	0.1681 (3)	0.0427 (10)
H43	0.7216	0.5602	0.1970	0.051*
C44	0.7585 (4)	0.6041 (2)	0.0931 (3)	0.0424 (11)
H44	0.7772	0.5598	0.0699	0.051*
C45	0.7642 (4)	0.6682 (3)	0.0528 (3)	0.0424 (10)
H45	0.7871	0.6681	0.0014	0.051*
C46	0.7371 (4)	0.7336 (2)	0.0858 (2)	0.0312 (8)
H46	0.7432	0.7776	0.0574	0.037*
C51	0.7630 (3)	0.82704 (19)	0.3072 (2)	0.0218 (7)
C52	0.8674 (3)	0.7857 (2)	0.3317 (2)	0.0285 (8)
H52	0.8875	0.7529	0.2930	0.034*
C53	0.9426 (3)	0.7919 (2)	0.4122 (2)	0.0327 (9)
H53	1.0135	0.7630	0.4284	0.039*
C54	0.9154 (3)	0.8397 (2)	0.4689 (2)	0.0329 (9)
H54	0.9670	0.8437	0.5241	0.039*
C55	0.8125 (4)	0.8818 (2)	0.4449 (2)	0.0349 (9)
H55	0.7940	0.9152	0.4837	0.042*
C56	0.7358 (3)	0.8758 (2)	0.3645 (2)	0.0291 (8)
H56	0.6650	0.9048	0.3486	0.035*
C61	0.7014 (3)	0.89120 (18)	0.1407 (2)	0.0211 (7)
C62	0.8040 (3)	0.9323 (2)	0.1700 (2)	0.0288 (8)
H62	0.8577	0.9215	0.2230	0.035*
C63	0.8293 (4)	0.9893 (2)	0.1227 (3)	0.0367 (9)
H63	0.9004	1.0173	0.1433	0.044*
C64	0.7520 (4)	1.0056 (2)	0.0460 (3)	0.0375 (9)
H64	0.7696	1.0449	0.0138	0.045*
C65	0.6488 (4)	0.9648 (2)	0.0157 (2)	0.0318 (8)
H65	0.5958	0.9759	-0.0374	0.038*
C66	0.6223 (3)	0.9076 (2)	0.0625 (2)	0.0261 (7)
H66	0.5511	0.8797	0.0418	0.031*
C1	0.4356 (5)	1.0112 (4)	0.4195 (3)	0.0687 (19)
H1	0.3902	1.0194	0.3633	0.082*
C2	0.4811 (6)	0.9443 (4)	0.4426 (4)	0.077 (2)
H2	0.4682	0.9059	0.4028	0.093*

C3	0.5455 (6)	0.9327 (4)	0.5236 (4)	0.080 (2)
H3	0.5774	0.8859	0.5407	0.096*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pd1	0.01820 (13)	0.01790 (14)	0.01891 (13)	0.00024 (10)	0.00593 (9)	0.00096 (10)
Cl1	0.0365 (5)	0.0263 (5)	0.0458 (6)	-0.0019 (4)	0.0154 (4)	-0.0002 (4)
Cl2	0.0279 (4)	0.0214 (4)	0.0400 (5)	0.0020 (3)	0.0116 (4)	-0.0015 (4)
P1	0.0186 (4)	0.0204 (4)	0.0166 (4)	-0.0005 (3)	0.0061 (3)	-0.0001 (3)
P2	0.0191 (4)	0.0201 (4)	0.0190 (4)	0.0008 (3)	0.0060 (3)	0.0011 (3)
C11	0.0201 (16)	0.0188 (17)	0.0254 (17)	0.0008 (13)	0.0082 (13)	-0.0007 (13)
C12	0.0251 (18)	0.0266 (19)	0.0256 (18)	0.0041 (14)	0.0079 (14)	-0.0004 (14)
C13	0.043 (2)	0.027 (2)	0.0301 (19)	0.0080 (16)	0.0168 (17)	0.0063 (15)
C14	0.043 (2)	0.024 (2)	0.050 (2)	-0.0011 (17)	0.028 (2)	0.0041 (17)
C15	0.0290 (19)	0.030 (2)	0.045 (2)	-0.0075 (16)	0.0129 (16)	-0.0047 (18)
C16	0.0248 (18)	0.031 (2)	0.0299 (19)	-0.0036 (15)	0.0084 (15)	-0.0018 (15)
C21	0.0277 (18)	0.0216 (18)	0.0184 (16)	-0.0039 (14)	0.0087 (13)	0.0007 (13)
C22	0.035 (2)	0.035 (2)	0.0235 (18)	0.0072 (16)	0.0096 (15)	-0.0008 (16)
C23	0.053 (3)	0.041 (2)	0.0216 (18)	0.001 (2)	0.0084 (17)	-0.0061 (17)
C24	0.043 (2)	0.041 (2)	0.0197 (17)	-0.0112 (19)	-0.0011 (16)	-0.0035 (16)
C25	0.027 (2)	0.048 (3)	0.031 (2)	-0.0006 (17)	-0.0037 (16)	0.0011 (18)
C26	0.0220 (17)	0.041 (2)	0.0227 (17)	0.0002 (15)	0.0040 (14)	-0.0057 (15)
C31	0.0177 (16)	0.0265 (18)	0.0176 (15)	0.0009 (13)	0.0034 (12)	-0.0051 (13)
C32	0.0304 (19)	0.029 (2)	0.0265 (18)	0.0022 (15)	0.0134 (15)	0.0000 (15)
C33	0.039 (2)	0.038 (2)	0.034 (2)	0.0046 (17)	0.0219 (17)	-0.0028 (17)
C34	0.033 (2)	0.039 (2)	0.0275 (19)	0.0101 (17)	0.0094 (16)	-0.0057 (16)
C35	0.0285 (18)	0.027 (2)	0.0283 (18)	0.0056 (15)	0.0057 (15)	0.0017 (15)
C36	0.0189 (16)	0.0287 (19)	0.0216 (16)	0.0020 (13)	0.0052 (13)	0.0003 (14)
C41	0.0194 (17)	0.0253 (19)	0.0270 (18)	0.0019 (13)	0.0018 (14)	-0.0003 (14)
C42	0.037 (2)	0.032 (2)	0.043 (2)	0.0045 (17)	0.0184 (18)	0.0009 (18)
C43	0.039 (2)	0.029 (2)	0.060 (3)	0.0024 (17)	0.014 (2)	0.002 (2)
C44	0.034 (2)	0.037 (2)	0.052 (3)	0.0040 (18)	0.0030 (19)	-0.015 (2)
C45	0.045 (2)	0.048 (3)	0.035 (2)	0.012 (2)	0.0112 (18)	-0.006 (2)
C46	0.035 (2)	0.034 (2)	0.0255 (19)	0.0057 (16)	0.0103 (16)	-0.0032 (15)
C51	0.0206 (16)	0.0272 (18)	0.0189 (15)	-0.0007 (14)	0.0077 (12)	0.0061 (14)
C52	0.0262 (18)	0.037 (2)	0.0238 (17)	0.0055 (15)	0.0090 (14)	0.0043 (15)
C53	0.0216 (18)	0.046 (2)	0.0304 (19)	0.0048 (16)	0.0055 (15)	0.0069 (18)
C54	0.0286 (19)	0.049 (3)	0.0196 (17)	-0.0083 (17)	0.0039 (14)	0.0028 (17)
C55	0.037 (2)	0.042 (2)	0.0264 (19)	0.0038 (18)	0.0099 (16)	-0.0045 (17)
C56	0.0265 (18)	0.029 (2)	0.0285 (18)	0.0058 (15)	0.0019 (15)	0.0000 (15)
C61	0.0208 (16)	0.0213 (17)	0.0238 (16)	0.0032 (13)	0.0108 (13)	-0.0002 (13)
C62	0.0252 (18)	0.032 (2)	0.0302 (19)	-0.0013 (15)	0.0090 (15)	0.0051 (16)
C63	0.030 (2)	0.034 (2)	0.048 (2)	-0.0062 (16)	0.0120 (17)	0.0080 (18)
C64	0.046 (2)	0.032 (2)	0.042 (2)	0.0038 (18)	0.0244 (19)	0.0128 (18)
C65	0.040 (2)	0.031 (2)	0.0256 (18)	0.0090 (17)	0.0109 (16)	0.0078 (15)
C66	0.0294 (19)	0.0261 (19)	0.0233 (17)	0.0034 (14)	0.0075 (14)	0.0012 (14)
C1	0.037 (3)	0.137 (6)	0.029 (2)	-0.005 (3)	0.0023 (19)	0.000 (3)
C2	0.069 (4)	0.113 (6)	0.062 (4)	-0.034 (4)	0.039 (3)	-0.031 (4)
C3	0.069 (4)	0.092 (5)	0.096 (5)	0.016 (4)	0.053 (4)	0.028 (4)

Geometric parameters (\AA , $\text{^{\circ}}$)

Pd1—Cl1	2.2851 (10)	C36—H36	0.9500
Pd1—Cl2	2.3056 (9)	C41—C46	1.386 (5)
Pd1—P2	2.3351 (9)	C41—C42	1.392 (6)
Pd1—P1	2.3399 (9)	C42—C43	1.388 (6)
P1—C11	1.819 (3)	C42—H42	0.9500
P1—C31	1.822 (3)	C43—C44	1.384 (7)
P1—C21	1.822 (3)	C43—H43	0.9500
P2—C61	1.820 (4)	C44—C45	1.366 (7)
P2—C51	1.824 (3)	C44—H44	0.9500
P2—C41	1.830 (4)	C45—C46	1.391 (6)
C11—C16	1.383 (5)	C45—H45	0.9500
C11—C12	1.394 (5)	C46—H46	0.9500
C12—C13	1.389 (5)	C51—C52	1.387 (5)
C12—H12	0.9500	C51—C56	1.396 (5)
C13—C14	1.383 (6)	C52—C53	1.386 (5)
C13—H13	0.9500	C52—H52	0.9500
C14—C15	1.377 (6)	C53—C54	1.378 (6)
C14—H14	0.9500	C53—H53	0.9500
C15—C16	1.383 (6)	C54—C55	1.381 (6)
C15—H15	0.9500	C54—H54	0.9500
C16—H16	0.9500	C55—C56	1.392 (5)
C21—C22	1.383 (5)	C55—H55	0.9500
C21—C26	1.395 (5)	C56—H56	0.9500
C22—C23	1.388 (5)	C61—C62	1.377 (5)
C22—H22	0.9500	C61—C66	1.402 (5)
C23—C24	1.376 (6)	C62—C63	1.383 (5)
C23—H23	0.9500	C62—H62	0.9500
C24—C25	1.383 (6)	C63—C64	1.375 (6)
C24—H24	0.9500	C63—H63	0.9500
C25—C26	1.383 (5)	C64—C65	1.381 (6)
C25—H25	0.9500	C64—H64	0.9500
C26—H26	0.9500	C65—C66	1.385 (5)
C31—C36	1.396 (5)	C65—H65	0.9500
C31—C32	1.400 (5)	C66—H66	0.9500
C32—C33	1.382 (5)	C1—C3 ⁱ	1.376 (6)
C32—H32	0.9500	C1—C2	1.355 (7)
C33—C34	1.364 (6)	C1—H1	0.9500
C33—H33	0.9500	C2—C3	1.365 (7)
C34—C35	1.396 (6)	C2—H2	0.9500
C34—H34	0.9500	C3—C1 ⁱ	1.376 (6)
C35—C36	1.383 (5)	C3—H3	0.9500
C35—H35	0.9500		
Cl1—Pd1—Cl2	177.80 (4)	C34—C35—H35	120.0
Cl1—Pd1—P2	90.75 (3)	C35—C36—C31	120.0 (3)
Cl2—Pd1—P2	89.20 (3)	C35—C36—H36	120.0
Cl1—Pd1—P1	86.35 (3)	C31—C36—H36	120.0
Cl2—Pd1—P1	93.58 (3)	C46—C41—C42	118.1 (4)

P2—Pd1—P1	175.72 (3)	C46—C41—P2	121.7 (3)
C11—P1—C31	103.13 (16)	C42—C41—P2	120.2 (3)
C11—P1—C21	104.49 (16)	C43—C42—C41	121.1 (4)
C31—P1—C21	104.77 (16)	C43—C42—H42	119.5
C11—P1—Pd1	114.15 (11)	C41—C42—H42	119.5
C31—P1—Pd1	120.63 (11)	C44—C43—C42	120.0 (4)
C21—P1—Pd1	108.21 (11)	C44—C43—H43	120.0
C61—P2—C51	105.54 (16)	C42—C43—H43	120.0
C61—P2—C41	105.34 (16)	C45—C44—C43	119.3 (4)
C51—P2—C41	104.81 (16)	C45—C44—H44	120.4
C61—P2—Pd1	110.59 (11)	C43—C44—H44	120.4
C51—P2—Pd1	112.49 (11)	C44—C45—C46	121.2 (4)
C41—P2—Pd1	117.17 (12)	C44—C45—H45	119.4
C16—C11—C12	118.8 (3)	C46—C45—H45	119.4
C16—C11—P1	122.7 (3)	C41—C46—C45	120.4 (4)
C12—C11—P1	118.4 (3)	C41—C46—H46	119.8
C13—C12—C11	120.4 (3)	C45—C46—H46	119.8
C13—C12—H12	119.8	C52—C51—C56	118.9 (3)
C11—C12—H12	119.8	C52—C51—P2	122.4 (3)
C14—C13—C12	119.9 (4)	C56—C51—P2	118.6 (3)
C14—C13—H13	120.0	C53—C52—C51	120.6 (4)
C12—C13—H13	120.0	C53—C52—H52	119.7
C15—C14—C13	119.9 (4)	C51—C52—H52	119.7
C15—C14—H14	120.0	C54—C53—C52	120.5 (4)
C13—C14—H14	120.0	C54—C53—H53	119.8
C14—C15—C16	120.2 (4)	C52—C53—H53	119.8
C14—C15—H15	119.9	C53—C54—C55	119.5 (3)
C16—C15—H15	119.9	C53—C54—H54	120.3
C11—C16—C15	120.8 (4)	C55—C54—H54	120.3
C11—C16—H16	119.6	C54—C55—C56	120.7 (4)
C15—C16—H16	119.6	C54—C55—H55	119.7
C22—C21—C26	119.3 (3)	C56—C55—H55	119.7
C22—C21—P1	118.9 (3)	C55—C56—C51	119.9 (3)
C26—C21—P1	121.8 (3)	C55—C56—H56	120.1
C21—C22—C23	120.2 (4)	C51—C56—H56	120.1
C21—C22—H22	119.9	C62—C61—C66	119.5 (3)
C23—C22—H22	119.9	C62—C61—P2	122.2 (3)
C24—C23—C22	120.6 (4)	C66—C61—P2	118.2 (3)
C24—C23—H23	119.7	C61—C62—C63	120.4 (4)
C22—C23—H23	119.7	C61—C62—H62	119.8
C23—C24—C25	119.4 (3)	C63—C62—H62	119.8
C23—C24—H24	120.3	C64—C63—C62	120.3 (4)
C25—C24—H24	120.3	C64—C63—H63	119.8
C24—C25—C26	120.6 (4)	C62—C63—H63	119.8
C24—C25—H25	119.7	C63—C64—C65	120.0 (4)
C26—C25—H25	119.7	C63—C64—H64	120.0
C25—C26—C21	119.9 (4)	C65—C64—H64	120.0
C25—C26—H26	120.0	C64—C65—C66	120.3 (4)
C21—C26—H26	120.0	C64—C65—H65	119.9

C36—C31—C32	119.2 (3)	C66—C65—H65	119.9
C36—C31—P1	119.5 (3)	C65—C66—C61	119.5 (3)
C32—C31—P1	121.2 (3)	C65—C66—H66	120.2
C33—C32—C31	119.8 (4)	C61—C66—H66	120.2
C33—C32—H32	120.1	C3 ⁱ —C1—C2	120.9 (5)
C31—C32—H32	120.1	C3 ⁱ —C1—H1	119.6
C34—C33—C32	121.0 (4)	C2—C1—H1	119.6
C34—C33—H33	119.5	C1—C2—C3	119.3 (6)
C32—C33—H33	119.5	C1—C2—H2	120.4
C33—C34—C35	119.9 (4)	C3—C2—H2	120.4
C33—C34—H34	120.0	C1 ⁱ —C3—C2	119.9 (6)
C35—C34—H34	120.0	C1 ⁱ —C3—H3	120.1
C36—C35—C34	120.1 (4)	C2—C3—H3	120.1
C36—C35—H35	120.0		
C11—Pd1—P1—C11	31.06 (13)	C32—C33—C34—C35	-0.2 (6)
C12—Pd1—P1—C11	-151.14 (12)	C33—C34—C35—C36	0.3 (6)
C11—Pd1—P1—C31	154.76 (13)	C34—C35—C36—C31	-0.2 (5)
C12—Pd1—P1—C31	-27.44 (13)	C32—C31—C36—C35	0.0 (5)
C11—Pd1—P1—C21	-84.79 (12)	P1—C31—C36—C35	179.8 (3)
C12—Pd1—P1—C21	93.00 (12)	C61—P2—C41—C46	-10.0 (3)
C11—Pd1—P2—C61	139.03 (12)	C51—P2—C41—C46	-121.1 (3)
C12—Pd1—P2—C61	-38.77 (12)	Pd1—P2—C41—C46	113.4 (3)
C11—Pd1—P2—C51	-103.24 (13)	C61—P2—C41—C42	171.1 (3)
C12—Pd1—P2—C51	78.96 (13)	C51—P2—C41—C42	60.0 (3)
C11—Pd1—P2—C41	18.33 (13)	Pd1—P2—C41—C42	-65.5 (3)
C12—Pd1—P2—C41	-159.48 (13)	C46—C41—C42—C43	-0.8 (6)
C31—P1—C11—C16	100.7 (3)	P2—C41—C42—C43	178.2 (3)
C21—P1—C11—C16	-8.6 (3)	C41—C42—C43—C44	-0.4 (7)
Pd1—P1—C11—C16	-126.6 (3)	C42—C43—C44—C45	0.7 (7)
C31—P1—C11—C12	-77.8 (3)	C43—C44—C45—C46	0.0 (7)
C21—P1—C11—C12	172.9 (3)	C42—C41—C46—C45	1.6 (6)
Pd1—P1—C11—C12	54.9 (3)	P2—C41—C46—C45	-177.4 (3)
C16—C11—C12—C13	-1.8 (5)	C44—C45—C46—C41	-1.2 (6)
P1—C11—C12—C13	176.7 (3)	C61—P2—C51—C52	-96.0 (3)
C11—C12—C13—C14	1.7 (6)	C41—P2—C51—C52	15.0 (3)
C12—C13—C14—C15	0.2 (6)	Pd1—P2—C51—C52	143.4 (3)
C13—C14—C15—C16	-2.0 (6)	C61—P2—C51—C56	84.8 (3)
C12—C11—C16—C15	0.0 (5)	C41—P2—C51—C56	-164.3 (3)
P1—C11—C16—C15	-178.5 (3)	Pd1—P2—C51—C56	-35.9 (3)
C14—C15—C16—C11	1.9 (6)	C56—C51—C52—C53	0.9 (6)
C11—P1—C21—C22	-85.4 (3)	P2—C51—C52—C53	-178.3 (3)
C31—P1—C21—C22	166.5 (3)	C51—C52—C53—C54	-0.6 (6)
Pd1—P1—C21—C22	36.6 (3)	C52—C53—C54—C55	-0.3 (6)
C11—P1—C21—C26	93.3 (3)	C53—C54—C55—C56	0.7 (6)
C31—P1—C21—C26	-14.8 (3)	C54—C55—C56—C51	-0.3 (6)
Pd1—P1—C21—C26	-144.7 (3)	C52—C51—C56—C55	-0.5 (6)
C26—C21—C22—C23	-1.4 (6)	P2—C51—C56—C55	178.8 (3)
P1—C21—C22—C23	177.3 (3)	C51—P2—C61—C62	8.2 (3)

supplementary materials

C21—C22—C23—C24	1.7 (7)	C41—P2—C61—C62	-102.4 (3)
C22—C23—C24—C25	-0.2 (7)	Pd1—P2—C61—C62	130.1 (3)
C23—C24—C25—C26	-1.5 (7)	C51—P2—C61—C66	-167.8 (3)
C24—C25—C26—C21	1.7 (6)	C41—P2—C61—C66	81.6 (3)
C22—C21—C26—C25	-0.3 (6)	Pd1—P2—C61—C66	-45.9 (3)
P1—C21—C26—C25	-179.0 (3)	C66—C61—C62—C63	-0.2 (6)
C11—P1—C31—C36	-170.3 (3)	P2—C61—C62—C63	-176.1 (3)
C21—P1—C31—C36	-61.2 (3)	C61—C62—C63—C64	0.2 (6)
Pd1—P1—C31—C36	60.9 (3)	C62—C63—C64—C65	-0.3 (6)
C11—P1—C31—C32	9.6 (3)	C63—C64—C65—C66	0.3 (6)
C21—P1—C31—C32	118.7 (3)	C64—C65—C66—C61	-0.3 (6)
Pd1—P1—C31—C32	-119.2 (3)	C62—C61—C66—C65	0.2 (5)
C36—C31—C32—C33	0.1 (5)	P2—C61—C66—C65	176.3 (3)
P1—C31—C32—C33	-179.7 (3)	C3 ⁱ —C1—C2—C3	0.5 (10)
C31—C32—C33—C34	0.0 (6)	C1—C2—C3—C1 ⁱ	-0.5 (10)

Symmetry code: (i) $-x+1, -y+2, -z+1$.