

Aqua(2,2'-bipyridine- $\kappa^2 N,N'$){(E)-[(5-chloro-2-oxidobenzylidene)amino- $\kappa^2 N,O$]methanesulfonato- κO }zinc

Yi-Fang Deng,* Xue Nie and Chun-Hua Zhang*

Hengyang Normal University, Department of Chemistry and Materials Science,
Hengyang, Hunan 421008, People's Republic of China
Correspondence e-mail: yifang7124@163.com, zhangchunhua668@163.com


Received 10 March 2012; accepted 25 March 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$;
 R factor = 0.028; wR factor = 0.077; data-to-parameter ratio = 13.9.

In the title compound, $[\text{Zn}(\text{C}_8\text{H}_6\text{ClNO}_4\text{S})(\text{C}_{10}\text{H}_8\text{N}_2)(\text{H}_2\text{O})]$, the Zn^{II} atom is six-coordinated by two O atoms and one N atom from a tridentate Schiff base ligand and two N atoms from a chelating 2,2'-bipyridine ligand and one water molecule, forming a slightly distorted octahedral geometry. In the crystal, $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link pairs of complex molecules into dimers. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond is also present.

Related literature

For related complexes, see: He *et al.* (2007); Xu *et al.* (2007).



Experimental

Crystal data

$[\text{Zn}(\text{C}_8\text{H}_6\text{ClNO}_4\text{S})(\text{C}_{10}\text{H}_8\text{N}_2)(\text{H}_2\text{O})]$	$\beta = 90.664 (2)^\circ$
$M_r = 487.22$	$\gamma = 98.993 (1)^\circ$
Triclinic, $P\bar{1}$	$V = 962.08 (13)\text{ \AA}^3$
$a = 7.7332 (6)\text{ \AA}$	$Z = 2$
$b = 10.8948 (8)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 11.9112 (9)\text{ \AA}$	$\mu = 1.56\text{ mm}^{-1}$
$\alpha = 103.625 (1)^\circ$	$T = 296\text{ K}$
	$0.18 \times 0.14 \times 0.10\text{ mm}$

Data collection

Bruker APEX CCD diffractometer	7155 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	3719 independent reflections
$T_{\min} = 0.766$, $T_{\max} = 0.860$	3345 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.077$	$\Delta\rho_{\text{max}} = 0.27\text{ e \AA}^{-3}$
$S = 1.05$	$\Delta\rho_{\text{min}} = -0.52\text{ e \AA}^{-3}$
3719 reflections	
267 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W—H1WA…O4 ⁱ	0.84 (3)	1.96 (3)	2.791 (2)	172 (3)
O1W—H1WB…O3	0.82	2.10	2.855 (2)	154

Symmetry code: (i) $-x + 1, -y + 1, -z + 2$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the Open Fund Project of Key Laboratories in Hunan Universities (11 K009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2525).

References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- He, K.-H., Li, J.-M. & Jiang, Y.-M. (2007). *Acta Cryst. E63*, m2992–m2993.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Xu, J.-S., Zhang, C.-H., Kuang, D.-Z., Feng, Y.-L. & Peng, Y.-L. (2007). *Acta Cryst. E63*, m2250.

supplementary materials

Acta Cryst. (2012). E68, m517 [doi:10.1107/S1600536812013050]

Aqua(2,2'-bipyridine- κ^2N,N'){(E)-[(5-chloro-2-oxidobenzylidene)amino- κ^2N,O]methanesulfonato- κO }zinc

Yi-Fang Deng, Xue Nie and Chun-Hua Zhang

Comment

Schiff base complexes containing amino acids have been studied for many years, arising interest because of their antiviral, anticancer and antibacterial activity. Herein, we choose amino-methanesulfonic acid-Schiff base to react with $Zn(CH_3CO_2)_2 \cdot 4H_2O$ as well as 2,2'-bipyridine. In the title compound, the Zn^{II} atom forms one five-membered and one six-membered chelating rings with the Schiff base ligand (Fig. 1). The coordinating environment of the Zn^{II} atom is different from that reported by He *et al.* (2007) and Xu *et al.* (2007). The bond length of $Zn—O(\text{sulfonate})$ is longer than that of $Zn—O(\text{aqua})$ and also longer than that of $Zn—N(\text{imine})$. It indicates that the coordinating capability of the sulfonate group is weaker than that of water and imine group. In the crystal, $O—H\cdots O$ hydrogen bonds link two complex molecules into a dimer (Table 1). An intramolecular $O—H\cdots O$ hydrogen bond is also present.

Experimental


The complex was prepared by mixing a methanol-water solution of 5-chlorosalicylaldehyde (1.0 mmol), aminomethane-sulfonic acid (1.0 mmol) and potassium hydrate (1.0 mmol) with heating and stirring. After 2 h, an aqueous solution containing zinc acetate (1 mmol) was added dropwise under stirring. The pH value of the mixture was adjusted to 6 with 0.5 mol/L HCl solution, followed by the dropwise addition of a methanol solution containing 2,2'-bipyridine (1 mmol) with stirring. The resulting yellow filtrate was allowed to stand at room temperature and slowly evaporate for one month to afford yellow block-shaped crystals.

Refinement

H atoms attached to C atoms were positioned geometrically and refined as riding atoms, with $C—H = 0.93$ (CH) and 0.97 (CH_2) Å and with $U_{\text{iso}}(H) = 1.2U_{\text{eq}}(C)$. The water H atoms were located from a difference Fourier map, one of them was refined isotropically and the other was refined as riding, with $U_{\text{iso}}(H) = 1.5U_{\text{eq}}(O)$.

Computing details

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

**Figure 1**

The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity.

Aqua(2,2'-bipyridine- κ^2N,N') $\{(E)$ -[(5-chloro-2-oxidobenzylidene)amino- κ^2N,O]methanesulfonato- κO }zinc

Crystal data

$[Zn(C_8H_6ClNO_4S)(C_{10}H_8N_2)(H_2O)]$
 $M_r = 487.22$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 7.7332 (6)$ Å
 $b = 10.8948 (8)$ Å
 $c = 11.9112 (9)$ Å
 $\alpha = 103.625 (1)^\circ$
 $\beta = 90.664 (2)^\circ$
 $\gamma = 98.993 (1)^\circ$
 $V = 962.08 (13)$ Å³

$Z = 2$
 $F(000) = 496$
 $D_x = 1.682$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 4928 reflections
 $\theta = 2.3\text{--}27.5^\circ$
 $\mu = 1.56$ mm⁻¹
 $T = 296$ K
Block, yellow
 $0.18 \times 0.14 \times 0.10$ mm

Data collection

Bruker APEX CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.766$, $T_{\max} = 0.860$

7155 measured reflections
3719 independent reflections
3345 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$
 $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.8^\circ$
 $h = -5 \rightarrow 9$
 $k = -13 \rightarrow 13$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.077$
 $S = 1.05$
3719 reflections
267 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.039P)^2 + 0.2378P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.27 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.52 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.62092 (3)	0.62060 (2)	0.841029 (19)	0.03292 (9)
C1	0.8756 (3)	0.68019 (19)	1.04504 (17)	0.0312 (4)
C2	0.9837 (3)	0.6387 (2)	1.11991 (18)	0.0382 (5)
H2	0.9938	0.5524	1.1046	0.046*
C3	1.0741 (3)	0.7212 (2)	1.21429 (18)	0.0382 (5)
H3	1.1456	0.6911	1.2614	0.046*
C4	1.0588 (3)	0.8500 (2)	1.23969 (18)	0.0367 (5)
C5	0.9626 (3)	0.8967 (2)	1.16761 (18)	0.0343 (4)
H5	0.9573	0.9838	1.1839	0.041*
C6	0.8707 (2)	0.81353 (18)	1.06824 (17)	0.0298 (4)
C7	0.7749 (3)	0.87525 (18)	0.99868 (17)	0.0305 (4)
H7	0.7835	0.9637	1.0229	0.037*
C8	0.5976 (3)	0.90082 (19)	0.84771 (18)	0.0358 (5)
H8A	0.6656	0.9161	0.7830	0.043*
H8B	0.5928	0.9827	0.9007	0.043*
C9	0.8962 (3)	0.7207 (2)	0.6752 (2)	0.0458 (6)
H9	0.9395	0.7860	0.7392	0.055*
C10	0.9547 (3)	0.7293 (2)	0.5682 (2)	0.0515 (6)
H10	1.0374	0.7980	0.5600	0.062*
C11	0.8876 (3)	0.6336 (3)	0.4738 (2)	0.0500 (6)
H11	0.9206	0.6389	0.4001	0.060*
C12	0.7714 (3)	0.5298 (2)	0.48847 (19)	0.0407 (5)
H12	0.7273	0.4634	0.4254	0.049*
C13	0.7217 (2)	0.52624 (19)	0.59897 (17)	0.0298 (4)
C14	0.6071 (2)	0.41548 (18)	0.62564 (17)	0.0299 (4)
C15	0.5698 (3)	0.2974 (2)	0.54736 (19)	0.0393 (5)
H15	0.6097	0.2864	0.4729	0.047*
C16	0.4727 (3)	0.1964 (2)	0.5817 (2)	0.0467 (6)
H16	0.4458	0.1166	0.5302	0.056*
C17	0.4161 (3)	0.2143 (2)	0.6919 (2)	0.0495 (6)
H17	0.3520	0.1469	0.7167	0.059*

C18	0.4562 (3)	0.3343 (2)	0.7653 (2)	0.0419 (5)
H18	0.4170	0.3465	0.8400	0.050*
Cl1	1.15819 (9)	0.95211 (6)	1.36830 (5)	0.05451 (17)
H1WA	0.340 (4)	0.550 (3)	0.984 (2)	0.050 (8)*
N1	0.7798 (2)	0.62241 (16)	0.69110 (15)	0.0341 (4)
N2	0.5490 (2)	0.43414 (15)	0.73402 (14)	0.0315 (4)
N3	0.6788 (2)	0.81919 (15)	0.90636 (14)	0.0315 (4)
O1	0.4109 (2)	0.69189 (14)	0.73855 (13)	0.0399 (4)
O2	0.3186 (2)	0.89098 (16)	0.72147 (16)	0.0539 (4)
O3	0.2802 (2)	0.81929 (16)	0.89951 (15)	0.0504 (4)
O4	0.7824 (2)	0.59406 (13)	0.96069 (12)	0.0381 (3)
O1W	0.3995 (2)	0.60387 (17)	0.95374 (16)	0.0495 (4)
H1WB	0.3365	0.6563	0.9481	0.074*
S1	0.38106 (7)	0.82061 (5)	0.79767 (5)	0.03552 (13)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.04229 (15)	0.02405 (14)	0.02831 (14)	0.00326 (10)	-0.00157 (10)	-0.00023 (10)
C1	0.0363 (10)	0.0297 (10)	0.0269 (10)	0.0079 (8)	0.0033 (8)	0.0039 (8)
C2	0.0467 (12)	0.0351 (11)	0.0361 (11)	0.0177 (9)	0.0029 (9)	0.0078 (9)
C3	0.0365 (11)	0.0483 (13)	0.0330 (11)	0.0121 (9)	-0.0015 (9)	0.0127 (10)
C4	0.0355 (11)	0.0407 (12)	0.0302 (11)	-0.0001 (9)	-0.0058 (8)	0.0060 (9)
C5	0.0378 (11)	0.0279 (10)	0.0346 (11)	0.0012 (8)	-0.0033 (8)	0.0052 (8)
C6	0.0316 (10)	0.0268 (10)	0.0293 (10)	0.0033 (8)	-0.0009 (8)	0.0044 (8)
C7	0.0344 (10)	0.0221 (9)	0.0320 (10)	0.0023 (8)	-0.0019 (8)	0.0022 (8)
C8	0.0432 (11)	0.0285 (10)	0.0351 (11)	0.0029 (9)	-0.0066 (9)	0.0089 (9)
C9	0.0433 (12)	0.0347 (12)	0.0544 (15)	-0.0009 (10)	0.0061 (11)	0.0053 (10)
C10	0.0452 (13)	0.0442 (14)	0.0691 (17)	0.0052 (11)	0.0157 (12)	0.0223 (13)
C11	0.0506 (14)	0.0620 (16)	0.0459 (14)	0.0147 (12)	0.0137 (11)	0.0255 (12)
C12	0.0427 (12)	0.0491 (13)	0.0316 (11)	0.0126 (10)	-0.0002 (9)	0.0091 (10)
C13	0.0291 (9)	0.0311 (10)	0.0297 (10)	0.0089 (8)	-0.0020 (8)	0.0057 (8)
C14	0.0279 (9)	0.0298 (10)	0.0298 (10)	0.0058 (8)	-0.0046 (8)	0.0022 (8)
C15	0.0435 (12)	0.0357 (12)	0.0330 (11)	0.0064 (9)	-0.0040 (9)	-0.0025 (9)
C16	0.0468 (13)	0.0305 (12)	0.0524 (14)	0.0001 (10)	-0.0051 (11)	-0.0061 (10)
C17	0.0467 (13)	0.0311 (12)	0.0643 (16)	-0.0058 (10)	0.0063 (12)	0.0069 (11)
C18	0.0437 (12)	0.0353 (12)	0.0439 (13)	0.0006 (9)	0.0102 (10)	0.0076 (10)
Cl1	0.0662 (4)	0.0497 (4)	0.0402 (3)	-0.0031 (3)	-0.0221 (3)	0.0057 (3)
N1	0.0362 (9)	0.0283 (9)	0.0346 (9)	0.0031 (7)	0.0010 (7)	0.0028 (7)
N2	0.0331 (8)	0.0270 (9)	0.0313 (9)	0.0035 (7)	0.0022 (7)	0.0019 (7)
N3	0.0369 (9)	0.0246 (8)	0.0321 (9)	0.0045 (7)	-0.0043 (7)	0.0058 (7)
O1	0.0480 (9)	0.0319 (8)	0.0350 (8)	0.0072 (6)	-0.0069 (7)	-0.0015 (6)
O2	0.0584 (10)	0.0472 (10)	0.0587 (11)	0.0110 (8)	-0.0202 (8)	0.0173 (8)
O3	0.0540 (10)	0.0400 (9)	0.0562 (11)	0.0119 (8)	0.0149 (8)	0.0062 (8)
O4	0.0517 (9)	0.0242 (7)	0.0351 (8)	0.0062 (6)	-0.0060 (7)	0.0010 (6)
O1W	0.0585 (11)	0.0410 (10)	0.0561 (11)	0.0136 (8)	0.0200 (9)	0.0211 (8)
S1	0.0397 (3)	0.0292 (3)	0.0357 (3)	0.0067 (2)	-0.0064 (2)	0.0036 (2)

Geometric parameters (\AA , $^\circ$)

Zn1—O4	1.9851 (15)	C9—C10	1.376 (3)
Zn1—N3	2.0936 (16)	C9—H9	0.9300
Zn1—N2	2.1144 (16)	C10—C11	1.372 (4)
Zn1—N1	2.1817 (18)	C10—H10	0.9300
Zn1—O1W	2.1999 (17)	C11—C12	1.377 (3)
Zn1—O1	2.3507 (15)	C11—H11	0.9300
C1—O4	1.318 (2)	C12—C13	1.383 (3)
C1—C2	1.411 (3)	C12—H12	0.9300
C1—C6	1.420 (3)	C13—N1	1.343 (3)
C2—C3	1.367 (3)	C13—C14	1.482 (3)
C2—H2	0.9300	C14—N2	1.352 (3)
C3—C4	1.388 (3)	C14—C15	1.386 (3)
C3—H3	0.9300	C15—C16	1.379 (3)
C4—C5	1.364 (3)	C15—H15	0.9300
C4—C11	1.754 (2)	C16—C17	1.368 (3)
C5—C6	1.414 (3)	C16—H16	0.9300
C5—H5	0.9300	C17—C18	1.378 (3)
C6—C7	1.446 (3)	C17—H17	0.9300
C7—N3	1.287 (2)	C18—N2	1.338 (3)
C7—H7	0.9300	C18—H18	0.9300
C8—N3	1.459 (3)	O1—S1	1.4682 (15)
C8—S1	1.788 (2)	O2—S1	1.4416 (17)
C8—H8A	0.9700	O3—S1	1.4513 (17)
C8—H8B	0.9700	O1W—H1WA	0.84 (3)
C9—N1	1.338 (3)	O1W—H1WB	0.8200
O4—Zn1—N3	90.71 (6)	C11—C10—H10	120.9
O4—Zn1—N2	102.79 (6)	C9—C10—H10	120.9
N3—Zn1—N2	164.81 (7)	C10—C11—C12	120.0 (2)
O4—Zn1—N1	104.54 (7)	C10—C11—H11	120.0
N3—Zn1—N1	93.88 (6)	C12—C11—H11	120.0
N2—Zn1—N1	76.17 (6)	C11—C12—C13	118.6 (2)
O4—Zn1—O1W	90.69 (7)	C11—C12—H12	120.7
N3—Zn1—O1W	92.08 (7)	C13—C12—H12	120.7
N2—Zn1—O1W	94.66 (7)	N1—C13—C12	121.86 (19)
N1—Zn1—O1W	163.55 (7)	N1—C13—C14	115.00 (17)
O4—Zn1—O1	165.47 (5)	C12—C13—C14	123.10 (18)
N3—Zn1—O1	78.29 (6)	N2—C14—C15	121.59 (19)
N2—Zn1—O1	89.43 (6)	N2—C14—C13	115.84 (16)
N1—Zn1—O1	85.85 (6)	C15—C14—C13	122.45 (19)
O1W—Zn1—O1	80.36 (6)	C16—C15—C14	119.0 (2)
O4—C1—C2	118.78 (18)	C16—C15—H15	120.5
O4—C1—C6	124.10 (18)	C14—C15—H15	120.5
C2—C1—C6	117.09 (18)	C17—C16—C15	119.6 (2)
C3—C2—C1	122.1 (2)	C17—C16—H16	120.2
C3—C2—H2	118.9	C15—C16—H16	120.2
C1—C2—H2	118.9	C16—C17—C18	118.6 (2)
C2—C3—C4	119.88 (19)	C16—C17—H17	120.7

C2—C3—H3	120.1	C18—C17—H17	120.7
C4—C3—H3	120.1	N2—C18—C17	123.0 (2)
C5—C4—C3	120.61 (19)	N2—C18—H18	118.5
C5—C4—Cl1	119.85 (17)	C17—C18—H18	118.5
C3—C4—Cl1	119.50 (16)	C9—N1—C13	118.41 (19)
C4—C5—C6	120.4 (2)	C9—N1—Zn1	126.34 (15)
C4—C5—H5	119.8	C13—N1—Zn1	113.33 (13)
C6—C5—H5	119.8	C18—N2—C14	118.20 (17)
C5—C6—C1	119.71 (18)	C18—N2—Zn1	125.78 (14)
C5—C6—C7	114.88 (17)	C14—N2—Zn1	116.00 (13)
C1—C6—C7	125.40 (17)	C7—N3—C8	116.93 (17)
N3—C7—C6	126.09 (18)	C7—N3—Zn1	124.61 (14)
N3—C7—H7	117.0	C8—N3—Zn1	118.37 (13)
C6—C7—H7	117.0	S1—O1—Zn1	111.64 (8)
N3—C8—S1	107.87 (14)	C1—O4—Zn1	128.73 (13)
N3—C8—H8A	110.1	Zn1—O1W—H1WA	140.4 (18)
S1—C8—H8A	110.1	Zn1—O1W—H1WB	109.5
N3—C8—H8B	110.1	H1WA—O1W—H1WB	106.3
S1—C8—H8B	110.1	O2—S1—O3	114.48 (11)
H8A—C8—H8B	108.4	O2—S1—O1	113.83 (10)
N1—C9—C10	122.9 (2)	O3—S1—O1	111.93 (10)
N1—C9—H9	118.6	O2—S1—C8	106.08 (10)
C10—C9—H9	118.6	O3—S1—C8	106.82 (10)
C11—C10—C9	118.2 (2)	O1—S1—C8	102.52 (9)
O4—C1—C2—C3	-175.1 (2)	C15—C14—N2—C18	1.2 (3)
C6—C1—C2—C3	3.1 (3)	C13—C14—N2—C18	-174.98 (18)
C1—C2—C3—C4	0.9 (3)	C15—C14—N2—Zn1	179.59 (15)
C2—C3—C4—C5	-3.9 (3)	C13—C14—N2—Zn1	3.4 (2)
C2—C3—C4—Cl1	173.90 (17)	O4—Zn1—N2—C18	65.54 (19)
C3—C4—C5—C6	2.6 (3)	N3—Zn1—N2—C18	-142.3 (2)
Cl1—C4—C5—C6	-175.13 (16)	N1—Zn1—N2—C18	167.61 (19)
C4—C5—C6—C1	1.5 (3)	O1W—Zn1—N2—C18	-26.24 (19)
C4—C5—C6—C7	-179.6 (2)	O1—Zn1—N2—C18	-106.52 (18)
O4—C1—C6—C5	173.83 (19)	O4—Zn1—N2—C14	-112.71 (14)
C2—C1—C6—C5	-4.3 (3)	N3—Zn1—N2—C14	39.5 (3)
O4—C1—C6—C7	-5.0 (3)	N1—Zn1—N2—C14	-10.64 (13)
C2—C1—C6—C7	176.93 (19)	O1W—Zn1—N2—C14	155.51 (14)
C5—C6—C7—N3	-178.2 (2)	O1—Zn1—N2—C14	75.23 (14)
C1—C6—C7—N3	0.6 (3)	C6—C7—N3—C8	-178.30 (19)
N1—C9—C10—C11	1.2 (4)	C6—C7—N3—Zn1	5.2 (3)
C9—C10—C11—C12	-3.0 (4)	S1—C8—N3—C7	-140.80 (16)
C10—C11—C12—C13	1.6 (4)	S1—C8—N3—Zn1	35.87 (18)
C11—C12—C13—N1	1.8 (3)	O4—Zn1—N3—C7	-5.63 (17)
C11—C12—C13—C14	-175.8 (2)	N2—Zn1—N3—C7	-158.6 (2)
N1—C13—C14—N2	11.9 (2)	N1—Zn1—N3—C7	-110.26 (17)
C12—C13—C14—N2	-170.33 (19)	O1W—Zn1—N3—C7	85.09 (17)
N1—C13—C14—C15	-164.25 (19)	O1—Zn1—N3—C7	164.80 (18)
C12—C13—C14—C15	13.5 (3)	O4—Zn1—N3—C8	177.97 (15)

N2—C14—C15—C16	−0.7 (3)	N2—Zn1—N3—C8	25.0 (3)
C13—C14—C15—C16	175.22 (19)	N1—Zn1—N3—C8	73.34 (15)
C14—C15—C16—C17	−0.4 (4)	O1W—Zn1—N3—C8	−91.31 (15)
C15—C16—C17—C18	0.9 (4)	O1—Zn1—N3—C8	−11.60 (14)
C16—C17—C18—N2	−0.4 (4)	O4—Zn1—O1—S1	20.4 (3)
C10—C9—N1—C13	2.1 (3)	N3—Zn1—O1—S1	−21.04 (9)
C10—C9—N1—Zn1	−161.05 (18)	N2—Zn1—O1—S1	167.95 (10)
C12—C13—N1—C9	−3.6 (3)	N1—Zn1—O1—S1	−115.88 (10)
C14—C13—N1—C9	174.19 (19)	O1W—Zn1—O1—S1	73.13 (10)
C12—C13—N1—Zn1	161.63 (16)	C2—C1—O4—Zn1	−179.41 (14)
C14—C13—N1—Zn1	−20.6 (2)	C6—C1—O4—Zn1	2.5 (3)
O4—Zn1—N1—C9	−79.2 (2)	N3—Zn1—O4—C1	1.91 (17)
N3—Zn1—N1—C9	12.5 (2)	N2—Zn1—O4—C1	174.89 (17)
N2—Zn1—N1—C9	−179.1 (2)	N1—Zn1—O4—C1	96.10 (17)
O1W—Zn1—N1—C9	123.5 (2)	O1W—Zn1—O4—C1	−90.18 (17)
O1—Zn1—N1—C9	90.44 (19)	O1—Zn1—O4—C1	−38.5 (3)
O4—Zn1—N1—C13	116.92 (14)	Zn1—O1—S1—O2	154.40 (10)
N3—Zn1—N1—C13	−151.34 (14)	Zn1—O1—S1—O3	−73.84 (11)
N2—Zn1—N1—C13	17.03 (13)	Zn1—O1—S1—C8	40.30 (11)
O1W—Zn1—N1—C13	−40.4 (3)	N3—C8—S1—O2	−168.93 (14)
O1—Zn1—N1—C13	−73.40 (14)	N3—C8—S1—O3	68.55 (16)
C17—C18—N2—C14	−0.6 (3)	N3—C8—S1—O1	−49.28 (16)
C17—C18—N2—Zn1	−178.84 (18)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1W—H1WA···O4 ⁱ	0.84 (3)	1.96 (3)	2.791 (2)	172 (3)
O1W—H1WB···O3	0.82	2.10	2.855 (2)	154

Symmetry code: (i) $-x+1, -y+1, -z+2$.