

(E)-2-(2-Hydroxy-5-iodobenzylidene)-hydrazinecarboxamide

Rahman Bikas,<sup>a\*</sup> Samra Nikbakht Sardari,<sup>b</sup> Seyed Sajjad Hosseini,<sup>b</sup> Gholam Hossein Shahverdizadeh<sup>c</sup> and Behrouz Notash<sup>d</sup>

<sup>a</sup>Young Researchers Club, Tabriz Branch, Islamic Azad University, Tabriz, Iran, <sup>b</sup>Department of Chemistry, Ardabil Branch, Islamic Azad University, Ardabil, Iran, <sup>c</sup>Department of Chemistry, Faculty of Science, Tabriz Branch, Islamic Azad University, PO Box 1655, Tabriz, Iran, and <sup>d</sup>Department of Chemistry, Shahid Beheshti University, G. C., Evin, Tehran, 1983963113, Iran  
Correspondence e-mail: bikas\_r@yahoo.com


Received 10 February 2012; accepted 7 March 2012

Key indicators: single-crystal X-ray study; *T* = 120 K; mean  $\sigma(\text{C}-\text{C})$  = 0.004 Å; *R* factor = 0.029; *wR* factor = 0.056; data-to-parameter ratio = 18.8.

In the title molecule, C<sub>8</sub>H<sub>8</sub>IN<sub>3</sub>O<sub>2</sub>, there is an intramolecular O—H···N hydrogen bond between the hydroxy group and the imine N atom, which generates an *S*(6) ring. In the crystal, the carbonyl O atom accepts two different N—H···O hydrogen bonds, which connect molecules with two *R*<sub>2</sub><sup>2</sup>(8) motifs.

Related literature

For historical background to semicarbazones, see: Arapov *et al.* (1987); Pickart *et al.* (1983). For related structures see: Bikas *et al.* (2010, 2012a,b); Monfared *et al.* (2010a). For background to the development of hydrazide derivatives for biological evaluation, see: Carvalho *et al.* (2008). For catalytic applications of aroylhydrazones, see: Monfared *et al.* (2010b). For a similar structure, see: Abboud *et al.* (1995).



Experimental

Crystal data

C<sub>8</sub>H<sub>8</sub>IN<sub>3</sub>O<sub>2</sub> *V* = 994.3 (3) Å<sup>3</sup>  
*M<sub>r</sub>* = 305.07 *Z* = 4  
 Monoclinic, *P*2<sub>1</sub>/*c* Mo *K*α radiation  
*a* = 9.1066 (18) Å  $\mu$  = 3.20 mm<sup>-1</sup>  
*b* = 7.6277 (15) Å *T* = 120 K  
*c* = 14.375 (3) Å 0.25 × 0.13 × 0.12 mm  
 $\beta$  = 95.31 (3)°

Data collection

Stoe IPDS 2T diffractometer 10438 measured reflections  
 Absorption correction: numerical 2362 independent reflections  
 (shape of crystal determined 2362 reflections with *I* > 2σ(*I*)  
 optically; *X-RED32* and *R<sub>int</sub>* = 0.041  
*X-SHAPE*, Stoe & Cie, 2005)  
*T<sub>min</sub>* = 0.502, *T<sub>max</sub>* = 0.700

Refinement

*R*[*F*<sup>2</sup> > 2σ(*F*<sup>2</sup>)] = 0.029 *H* atoms treated by a mixture of  
*wR*(*F*<sup>2</sup>) = 0.056 independent and constrained  
*S* = 1.13 refinement  
 2686 reflections  $\Delta\rho_{\text{max}}$  = 0.75 e Å<sup>-3</sup>  
 143 parameters  $\Delta\rho_{\text{min}}$  = -0.71 e Å<sup>-3</sup>  
 1 restraint

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> — <i>H</i> ··· <i>A</i>	<i>D</i> — <i>H</i>	<i>H</i> ··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> — <i>H</i> ··· <i>A</i>
N3—H3B···O2 <sup>i</sup>	0.81 (4)	2.13 (4)	2.920 (3)	163 (3)
N2—H2···O2 <sup>ii</sup>	0.80 (4)	2.00 (4)	2.800 (3)	176 (3)
O1—H1···N1	0.84 (2)	1.88 (3)	2.628 (3)	147 (4)

Symmetry codes: (i)  $-x + 3, y, -z + \frac{3}{2}$ ; (ii)  $-x + 3, -y + 2, -z + 2$ .

Data collection: *X-AREA* (Stoe & Cie, 2005); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors are grateful to the Islamic Azad University (Tabriz Branch) and the Islamic Azad University (Ardabil Branch) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VM2154).

References

Abboud, K. A., Summers, S. P. & Palenik, G. J. (1995). *Acta Cryst.* **C51**, 1707–1709.  
 Arapov, O. V., Alferva, O. F., Levocheskaya, E. I. & Krasilnikov, I. (1987). *Radiobiologiya*, **27**, 843–846.  
 Bikas, R., Anarjan, P. M., Ng, S. W. & Tiekink, E. R. T. (2012a). *Acta Cryst.* **E68**, o193.  
 Bikas, R., Anarjan, P. M., Ng, S. W. & Tiekink, E. R. T. (2012b). *Acta Cryst.* **E68**, o413–o414.  
 Bikas, R., Hosseini Monfared, H., Kazak, C., Arslan, N. B. & Bijanzad, K. (2010). *Acta Cryst.* **E66**, o2015.  
 Carvalho, S. R., da Silva, E. F., de Souza, M. V. N., Lourenco, M. C. S. & Vicente, F. R. (2008). *Bioorg. Med. Chem. Lett.* **18**, 538–541.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.  
 Monfared, H. H., Bikas, R. & Mayer, P. (2010a). *Acta Cryst.* **E66**, o236–o237.  
 Monfared, H. H., Bikas, R. & Mayer, P. (2010b). *Inorg. Chim. Acta*, **363**, 574–2583.  
 Pickart, L., Goodwin, W. H., Burgua, W., Murphy, T. B. & Johnson, D. K. (1983). *Biochem. Pharmacol.* **32**, 3868–3871.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Stoe & Cie (2005). *X-AREA*, *X-RED32* and *X-SHAPE*. Stoe & Cie, Darmstadt, Germany.

## supplementary materials

*Acta Cryst.* (2012). E68, o1090 [doi:10.1107/S1600536812010197]

**(*E*)-2-(2-Hydroxy-5-iodobenzylidene)hydrazinecarboxamide**

**Rahman Bikas, Samra Nikbakht Sardari, Seyed Sajjad Hosseini, Gholam Hossein Shahverdizadeh and Behrouz Notash**

**Comment**

Semicarbazone compounds are derived from the condensation of carbonyl compounds and semicarbazides. This class are important tridentate O, N, O-donor ligands. As biologically active compounds, semicarbazones find application in the treatment of diseases such as anti-tumor, tuberculosis, leprosy and mental disorder. Furthermore, semicarbazone have wide spread applications in fields such as coordination chemistry, bioinorganic chemistry, and in magnetic, electronic, nonlinear optically active and fluorescent compounds. Also semicarbazone metal complexes seem to be a good candidate for catalytic oxidation studies because of their resist to oxidation (Monfared *et al.*, 2010b).

As part of our studies on the synthesis and characterization of hydrazone derivatives (Bikas *et al.*, 2010; Bikas *et al.*, 2012*a,b*), we report here the crystal structure of (*E*)-2-(2-hydroxy-5-iodobenzylidene)hydrazinecarboxamide (Fig. 1). Bond distances are in the normal range for similar hydrazone compounds (Abboud *et al.*, 1995). The molecule is approximately planar, with an r.m.s. deviation from the mean plane through all 14 non-H atoms of 0.181 (2) Å. The dihedral angle between the phenyl ring plane and the least-squares plane through the N3—C8—O2—N2 unit is 14.00 (13)°. In the crystal structure of the title compound, the molecule adopts an *E* configuration with respect to the C7=N1 bond. In the crystal structure of the title compound, there is an intramolecular O—H···N hydrogen bonding between the hydroxyl group and imine nitrogen atom. The carbonyl group forms two different intermolecular N—H···O hydrogen bonds parallel to *ac*- plane which connects molecules with two  $R_2^2(8)$  motifs (Table 1, Fig. 2).

**Experimental**


For preparing the title compound a methanol (10 ml) solution of 2-hydroxy-5-iodobenzaldehyde (1.5 mmol) was added drop-wise to a methanol solution (10 ml) of semicarbazide (1.5 mmol), and the mixture was refluxed for 3 h. The solution was then evaporated on a steam bath to 5 ml and cooled to room temperature. The light-yellow precipitates of the title compound were separated and filtered off, washed with 3 ml of cooled methanol and then dried in air. Colorless crystals were obtained from its methanol solution by slow solvent evaporation. Yield: 92%. IR (cm<sup>-1</sup>): 3464 (m, O—H), 3176 (m, broad, N—H), 1699 (vs, C=O), 1594 (s, C=N), 1463 (s), 1340 (m), 1259 (vs), 1187 (s), 1072 (m), 942 (vs), 893 (m), 818 (m), 769 (vs), 682 (m), 613 (m), 572 (vs), 517 (s), 522 (m), 472 (vs), 427 (vs).

**Refinement**


The hydrogen atoms of the N—H and O—H groups were found in a difference Fourier map and refined isotropically with a distance restraint to 0.84 Å for the O—H group. All other H atoms were positioned geometrically and refined as riding atoms with C—H = 0.95 Å,  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  aromatic and imine H atoms.

**Computing details**

Data collection: *X-AREA* (Stoe & Cie, 2005); cell refinement: *X-AREA* (Stoe & Cie, 2005); data reduction: *X-AREA* (Stoe & Cie, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

**Figure 1**

The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

The packing diagram of the title compound showing intermolecular hydrogen bonds as blue dashed lines.

(E)-2-(2-Hydroxy-5-iodobenzylidene)hydrazinecarboxamide

Crystal data

$C_8H_8IN_3O_2$	$F(000) = 584$
$M_r = 305.07$	$D_x = 2.038 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-P 2_1c$	Cell parameters from 2686 reflections
$a = 9.1066 (18) \text{ \AA}$	$\theta = 2.7\text{--}29.2^\circ$
$b = 7.6277 (15) \text{ \AA}$	$\mu = 3.20 \text{ mm}^{-1}$
$c = 14.375 (3) \text{ \AA}$	$T = 120 \text{ K}$
$\beta = 95.31 (3)^\circ$	Block, colorless
$V = 994.3 (3) \text{ \AA}^3$	$0.25 \times 0.13 \times 0.12 \text{ mm}$
$Z = 4$	

Data collection

Stoe IPDS 2T	$T_{\min} = 0.502, T_{\max} = 0.700$
diffractometer	10438 measured reflections
Radiation source: fine-focus sealed tube	2686 independent reflections
Graphite monochromator	2362 reflections with $I > 2\sigma(I)$
Detector resolution: $0.15 \text{ mm pixels mm}^{-1}$	$R_{\text{int}} = 0.041$
rotation method scans	$\theta_{\max} = 29.2^\circ, \theta_{\min} = 2.7^\circ$
Absorption correction: numerical	$h = -12 \rightarrow 12$
(shape of crystal determined optically; $X$ - $RED32$ and $X$ - $SHAPE$ , Stoe & Cie, 2005)	$k = -10 \rightarrow 10$
	$l = -18 \rightarrow 19$

Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.029$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.056$	$w = 1/[\sigma^2(F_o^2) + (0.0208P)^2 + 1.0932P]$
$S = 1.13$	where $P = (F_o^2 + 2F_c^2)/3$
2686 reflections	$(\Delta/\sigma)_{\max} = 0.001$
143 parameters	$\Delta\rho_{\max} = 0.75 \text{ e \AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.71 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	0.71642 (2)	0.52615 (2)	1.210107 (13)	0.02620 (6)
O2	1.52011 (18)	1.0591 (2)	0.88088 (11)	0.0160 (3)
N2	1.3349 (2)	0.9116 (3)	0.93933 (14)	0.0149 (4)

C8	1.3955 (2)	0.9896 (3)	0.86612 (16)	0.0137 (4)
N3	1.3205 (2)	0.9864 (3)	0.78198 (15)	0.0166 (4)
N1	1.1902 (2)	0.8602 (3)	0.93061 (14)	0.0134 (4)
O1	0.9242 (2)	0.8354 (3)	0.84348 (13)	0.0217 (4)
C1	0.9856 (3)	0.7452 (3)	1.00333 (17)	0.0139 (4)
C2	0.8840 (3)	0.7661 (3)	0.92413 (17)	0.0150 (4)
C7	1.1398 (3)	0.7979 (3)	1.00415 (16)	0.0137 (4)
H7	1.2036	0.7860	1.0599	0.016*
C6	0.9357 (3)	0.6755 (3)	1.08517 (17)	0.0165 (5)
H6	1.0028	0.6609	1.1393	0.020*
C4	0.6899 (3)	0.6494 (3)	1.00927 (19)	0.0191 (5)
H4	0.5895	0.6172	1.0115	0.023*
C3	0.7373 (3)	0.7178 (3)	0.92780 (18)	0.0187 (5)
H3	0.6691	0.7319	0.8742	0.022*
C5	0.7893 (3)	0.6278 (3)	1.08752 (18)	0.0173 (5)
H2	1.380 (4)	0.917 (4)	0.990 (3)	0.022 (8)*
H3A	1.244 (4)	0.929 (5)	0.773 (3)	0.029 (9)*
H3B	1.359 (4)	1.028 (5)	0.738 (3)	0.028 (9)*
H1	1.016 (2)	0.849 (5)	0.849 (3)	0.037 (10)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
I1	0.03341 (10)	0.02256 (9)	0.02507 (9)	-0.00405 (7)	0.01569 (7)	0.00293 (7)
O2	0.0139 (7)	0.0241 (10)	0.0102 (8)	-0.0040 (6)	0.0015 (6)	-0.0006 (6)
N2	0.0121 (9)	0.0243 (10)	0.0081 (9)	-0.0052 (8)	0.0003 (7)	0.0002 (8)
C8	0.0143 (9)	0.0174 (12)	0.0095 (9)	0.0005 (8)	0.0019 (8)	-0.0004 (8)
N3	0.0153 (9)	0.0243 (11)	0.0101 (9)	-0.0040 (8)	0.0007 (7)	0.0005 (8)
N1	0.0112 (9)	0.0159 (9)	0.0132 (9)	-0.0012 (7)	0.0011 (7)	-0.0023 (7)
O1	0.0150 (9)	0.0375 (11)	0.0121 (8)	-0.0015 (7)	-0.0009 (7)	0.0058 (7)
C1	0.0149 (11)	0.0130 (12)	0.0141 (10)	-0.0002 (8)	0.0032 (8)	-0.0014 (8)
C2	0.0147 (11)	0.0174 (11)	0.0131 (11)	-0.0005 (8)	0.0020 (8)	-0.0005 (9)
C7	0.0130 (10)	0.0171 (11)	0.0109 (10)	-0.0006 (8)	0.0010 (8)	-0.0011 (8)
C6	0.0176 (11)	0.0180 (12)	0.0140 (11)	0.0003 (9)	0.0030 (9)	0.0010 (9)
C4	0.0147 (11)	0.0188 (12)	0.0249 (13)	-0.0037 (9)	0.0070 (9)	-0.0054 (10)
C3	0.0149 (11)	0.0231 (13)	0.0181 (12)	0.0001 (9)	0.0017 (9)	-0.0031 (9)
C5	0.0204 (11)	0.0137 (11)	0.0194 (12)	-0.0014 (8)	0.0106 (9)	0.0005 (9)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

I1—C5	2.089 (2)	C1—C6	1.404 (3)
O2—C8	1.253 (3)	C1—C2	1.408 (3)
N2—C8	1.369 (3)	C1—C7	1.460 (3)
N2—N1	1.369 (3)	C2—C3	1.391 (3)
N2—H2	0.80 (4)	C7—H7	0.9500
C8—N3	1.333 (3)	C6—C5	1.385 (3)
N3—H3A	0.82 (4)	C6—H6	0.9500
N3—H3B	0.81 (4)	C4—C5	1.387 (4)
N1—C7	1.282 (3)	C4—C3	1.387 (4)
O1—C2	1.355 (3)	C4—H4	0.9500

O1—H1	0.841 (18)	C3—H3	0.9500
C8—N2—N1	120.5 (2)	C3—C2—C1	120.0 (2)
C8—N2—H2	118 (2)	N1—C7—C1	120.9 (2)
N1—N2—H2	120 (2)	N1—C7—H7	119.6
O2—C8—N3	122.8 (2)	C1—C7—H7	119.6
O2—C8—N2	118.5 (2)	C5—C6—C1	120.4 (2)
N3—C8—N2	118.7 (2)	C5—C6—H6	119.8
C8—N3—H3A	121 (3)	C1—C6—H6	119.8
C8—N3—H3B	118 (3)	C5—C4—C3	119.9 (2)
H3A—N3—H3B	120 (4)	C5—C4—H4	120.0
C7—N1—N2	116.5 (2)	C3—C4—H4	120.0
C2—O1—H1	108 (3)	C4—C3—C2	120.4 (2)
C6—C1—C2	118.8 (2)	C4—C3—H3	119.8
C6—C1—C7	118.9 (2)	C2—C3—H3	119.8
C2—C1—C7	122.3 (2)	C6—C5—C4	120.4 (2)
O1—C2—C3	118.2 (2)	C6—C5—I1	120.0 (2)
O1—C2—C1	121.8 (2)	C4—C5—I1	119.59 (17)
N1—N2—C8—O2	169.1 (2)	C2—C1—C6—C5	0.2 (4)
N1—N2—C8—N3	-11.9 (3)	C7—C1—C6—C5	178.5 (2)
C8—N2—N1—C7	-175.9 (2)	C5—C4—C3—C2	-0.3 (4)
C6—C1—C2—O1	178.8 (2)	O1—C2—C3—C4	-178.8 (2)
C7—C1—C2—O1	0.6 (4)	C1—C2—C3—C4	0.1 (4)
C6—C1—C2—C3	-0.1 (4)	C1—C6—C5—C4	-0.4 (4)
C7—C1—C2—C3	-178.3 (2)	C1—C6—C5—I1	-179.51 (18)
N2—N1—C7—C1	177.8 (2)	C3—C4—C5—C6	0.5 (4)
C6—C1—C7—N1	179.1 (2)	C3—C4—C5—I1	179.59 (19)
C2—C1—C7—N1	-2.6 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—H3B...O2 <sup>i</sup>	0.81 (4)	2.13 (4)	2.920 (3)	163 (3)
N2—H2...O2 <sup>ii</sup>	0.80 (4)	2.00 (4)	2.800 (3)	176 (3)
O1—H1...N1	0.84 (2)	1.88 (3)	2.628 (3)	147 (4)

Symmetry codes: (i)  $-x+3, y, -z+3/2$ ; (ii)  $-x+3, -y+2, -z+2$ .