

4-[5-(Furan-2-yl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide

Abdullah M. Asiri,^{a,b,‡} Hassan M. Faidallah,^a Khalid A. Alamry,^a Seik Weng Ng^{c,a} and Edward R. T. Tiekink^{c*}

^aChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203, Jeddah, Saudi Arabia, ^bThe Center of Excellence for Advanced Materials Research, King Abdulaziz University, Jeddah, PO Box 80203, Saudi Arabia, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: edward.tiekink@gmail.com


Received 17 March 2012; accepted 19 March 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; disorder in main residue; R factor = 0.069; wR factor = 0.189; data-to-parameter ratio = 16.3.

In the title compound, $\text{C}_{14}\text{H}_{10}\text{F}_3\text{N}_3\text{O}_3\text{S}$, there are significant twists in the molecule, as seen in the values of the dihedral angles between the pyrazole ring and each of the furan [31.1 (2)°] and benzene rings [55.58 (10)°]. The amino N atom occupies a position almost normal to the benzene ring [$\text{N}-\text{S}-\text{C}_{\text{ar}}-\text{C}_{\text{ar}}$ (ar = aromatic) torsion angle = 83.70 (19)°]. One amino H atom forms a hydrogen bond to the tricoordinate pyrazole N atom and the other interacts with a sulfonamide O atom, forming a supramolecular chain along [010]. The chains are consolidated into a supramolecular layers *via* $\text{C}-\text{H}\cdots\text{O}$ interactions involving the second sulfonamide O atom; layers stack along [10 $\bar{1}$]. The furan ring was found to be disordered over two diagonally opposite orientations of equal occupancy.

Related literature

For background to the biological applications of sulfonamides, see: Croitoru *et al.* (2004); Dogruer *et al.* (2010). For the biological efficacy of F and CF_3 in medicinal chemistry, see: Fokin & Kolomiyets (1988); Bonaccorso *et al.* (2006). For related structures, see: Asiri *et al.* (2011, 2012).



[‡] Additional correspondence author, e-mail: aasiri2@kau.edu.sa.

Experimental

Crystal data

$\text{C}_{14}\text{H}_{10}\text{F}_3\text{N}_3\text{O}_3\text{S}$
 $M_r = 357.31$
Monoclinic, $P2_1/n$
 $a = 16.0536$ (13) Å
 $b = 4.8173$ (4) Å
 $c = 20.6202$ (15) Å
 $\beta = 110.728$ (8)°

$V = 1491.4$ (2) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.27$ mm⁻¹
 $T = 100$ K
0.25 × 0.10 × 0.05 mm

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2011)
 $T_{\text{min}} = 0.935$, $T_{\text{max}} = 0.987$

6503 measured reflections
3431 independent reflections
2400 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.069$
 $wR(F^2) = 0.189$
 $S = 1.06$
3431 reflections
210 parameters
33 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.57$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.63$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N3}-\text{H1}\cdots\text{O3}^{\text{i}}$	0.88 (1)	2.00 (2)	2.830 (4)	158 (3)
$\text{N3}-\text{H2}\cdots\text{N1}^{\text{ii}}$	0.88 (1)	2.17 (1)	3.032 (4)	170 (4)
$\text{C8}-\text{H8}\cdots\text{O2}^{\text{iii}}$	0.95	2.36	3.185 (7)	145
$\text{C10}-\text{H10}\cdots\text{O2}^{\text{iv}}$	0.95	2.44	3.092 (4)	126

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (iv) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *PUBLICIF* (Westrip, 2010).

The authors are thankful to the Center of Excellence for Advanced Materials Research and the Chemistry Department at King Abdulaziz University for providing the research facilities. We also thank the Ministry of Higher Education (Malaysia) for funding structural studies through the High-Impact Research scheme (UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6693).

References

- Agilent (2011). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
Asiri, A. M., Al-Youbi, A. O., Faidallah, H. M., Ng, S. W. & Tiekink, E. R. T. (2011). *Acta Cryst.* **E67**, o2424.
Asiri, A. M., Faidallah, H. M., Ng, S. W. & Tiekink, E. R. T. (2012). *Acta Cryst.* **E68**, o762–o763.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.

- Bonacorso, H. G., Wentz, A. P., Lourega, R. V., Cechinel, C. A., Moraes, T. S., Coelho, H. S., Zanatta, N., Martins, M. A. P., Hoerner, M. & Alves, S. H. (2006). *J. Fluorine Chem.* **127**, 1066–1072.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Croitoru, M., Pintilie, L., Tanase, C., Caproiu, M. T. & Draghici, C. (2004). *Rev. Chem. (Bucharest)*, **55**, 993–997.
- Dogruer, D. S., Urtu, S., Onkol, T., Ozcelik, B. & Sahin, M. F. (2010). *Turk. J. Chem.* **34**, 57–65.
- Fokin, A. V. & Kolomiyets, A. F. (1988). *J. Fluorine Chem.* **40**, 247–259.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o1168–o1169 [doi:10.1107/S1600536812011920]

4-[5-(Furan-2-yl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide

Abdullah M. Asiri, Hassan M. Faidallah, Khalid A. Alamry, Seik Weng Ng and Edward R. T. Tiekink

Comment

The title CF₃-derivatized sulfonamide (I), was investigated. in connection with on-going studies of sulfonamides, both biological (Croitoru *et al.*, 2004; Dogruer *et al.*, 2010) and crystallographic (Asiri *et al.*, 2011; Asiri *et al.*, 2012). In particular, fluoride in the form of a trifluoromethyl group, which has long been recognized in medicinal chemistry for its ability to alter the physico-chemical and biological characteristics of molecules (Fokin & Kolomiyets, 1988; Bonacorso *et al.*, 2006), is featured in the new molecule to promote enhanced biological properties.

In (I), Fig. 1, the dihedral angle of 31.1 (2)° between the furanyl and pyrazole rings indicates a significant twist between these rings. Similarly, the dihedral angle of 55.58 (10)° between the pyrazole ring and the benzene ring to which it is connected indicates a significant twist. The amino-N3 atom occupies a position almost normal to the benzene ring, forming a N3—S1—C12—C13 torsion angle of 83.70 (19)°. This allows the participation of both N—H atoms in hydrogen bonding interactions.

One amino-H atom forms a hydrogen bond to the pyrazole-N2 atom of a centrosymmetrically related molecule to form an 18-membered {···HN₂SC₄NN₂}₂ synthon, Table 1. These are connected into a supramolecular chain *via* a N—H···O(sulfonamide) hydrogen bonding, Fig. 2 and Table 1. The second sulfonamide-O2 atom forms two C—H···O interactions, Table 1, to consolidate the chains into a supramolecular layers. No specific intermolecular interactions occur between the layers that stack along [1 0 $\bar{1}$], Fig. 3.

Experimental

A solution of 4,4,4-trifluoro-1-phenyl-1,3-butanedione (2.16 g, 0.01 mmol) in ethanol (50 ml) was refluxed with 4-hydrazinobenzenesulfonamide hydrochloride (2.2 g, 0.01 mmol) for 4 h, concentrated and cooled. The precipitated crude product was filtered and recrystallized from ethanol as colourless prisms. Yield: 74%. *M.pt.*: 467–468 K.

Refinement


Carbon-bound H-atoms were placed in calculated positions [C—H = 0.95 Å; $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation. The N—H atoms were located in a difference Fourier map, and were refined with a distance restraint of N—H = 0.88±0.01 Å; their U_{iso} values were refined.

The furyl ring is disordered over two positions; the disorder could not be refined, and was assumed as a 1:1 type of disorder. The ring was refined as a rigid pentagon of 1.35 Å sides. The U_{ij} values of C6' was equated to those of O6 (as well as the O1'/C6, C5'/C5, C6'/O1, C7'/C8 and C8'/C7 pairs). The benzene ring was refined as a rigid hexagon of 1.39 Å sides.

Owing to poor agreement, the ($\bar{8}$ 0 10) reflection was omitted from the final cycles of refinement.

Computing details

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO* (Agilent, 2011); data reduction: *CrysAlis PRO* (Agilent, 2011); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

**Figure 1**

The molecular structure of (I) showing displacement ellipsoids at the 70% probability level. Only one orientation of the disordered furanyl ring is shown.





Figure 2

Supramolecular chain along [010] in (I) sustained by N—H \cdots N and N—H \cdots O hydrogen bonds shown as blue and orange dashed lines, respectively. Hydrogen atoms not participating in hydrogen-bonding interactions have been omitted for reasons of clarity.

**Figure 3**

A view in projection down the b axis of the unit-cell contents of (I). The N—H...N, N—H...O and C—H...O interactions are shown as orange, blue and brown dashed lines, respectively.

4-[5-(Furan-2-yl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide

Crystal data

$C_{14}H_{10}F_3N_3O_3S$
 $M_r = 357.31$
 Monoclinic, $P2_1/n$
 Hall symbol: -P 2yn
 $a = 16.0536$ (13) Å
 $b = 4.8173$ (4) Å
 $c = 20.6202$ (15) Å
 $\beta = 110.728$ (8)°
 $V = 1491.4$ (2) Å³
 $Z = 4$

$F(000) = 728$
 $D_x = 1.591$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 1899 reflections
 $\theta = 2.5$ – 27.5 °
 $\mu = 0.27$ mm⁻¹
 $T = 100$ K
 Prism, colourless
 $0.25 \times 0.10 \times 0.05$ mm

Data collection

Agilent SuperNova Dual
 diffractometer with an Atlas detector
 Radiation source: SuperNova (Mo) X-ray
 Source
 Mirror monochromator
 Detector resolution: 10.4041 pixels mm⁻¹
 ω scan
 Absorption correction: multi-scan
 (CrysAlis PRO; Agilent, 2011)

$T_{\min} = 0.935$, $T_{\max} = 0.987$
 6503 measured reflections
 3431 independent reflections
 2400 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$
 $\theta_{\max} = 27.6$ °, $\theta_{\min} = 2.7$ °
 $h = -20 \rightarrow 15$
 $k = -6 \rightarrow 6$
 $l = -26 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.069$

$wR(F^2) = 0.189$

$S = 1.06$

3431 reflections

210 parameters

33 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0768P)^2 + 1.5964P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.57 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.63 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.55947 (6)	0.78623 (17)	0.68278 (5)	0.0271 (3)	
F1	0.67465 (17)	-0.1684 (5)	0.34173 (13)	0.0506 (7)	
F2	0.7248 (2)	0.1593 (5)	0.29748 (11)	0.0559 (7)	
F3	0.81166 (17)	-0.1668 (6)	0.35146 (14)	0.0605 (8)	
O1	0.8796 (3)	0.7601 (14)	0.6001 (3)	0.0229 (13)	0.50
C5	0.9023 (5)	0.5273 (19)	0.5738 (4)	0.0223 (8)	0.50
C6	0.9876 (5)	0.4658 (18)	0.6122 (5)	0.0330 (13)	0.50
H6	1.0207	0.3124	0.6052	0.040*	0.50
C7	1.0177 (3)	0.6606 (15)	0.6621 (3)	0.0298 (17)	0.50
H7	1.0757	0.6683	0.6965	0.036*	0.50
C8	0.9510 (4)	0.8425 (11)	0.6547 (3)	0.035 (2)	0.50
H8	0.9537	1.0006	0.6829	0.042*	0.50
O1'	0.9946 (4)	0.4216 (15)	0.5961 (3)	0.0330 (13)	0.50
C5'	0.9085 (4)	0.501 (2)	0.5745 (4)	0.0223 (8)	0.50
C6'	0.9012 (4)	0.7080 (18)	0.6164 (4)	0.0229 (13)	0.50
H6'	0.8480	0.8029	0.6130	0.028*	0.50
C7'	0.9829 (4)	0.7561 (13)	0.6638 (3)	0.035 (2)	0.50
H7'	0.9972	0.8908	0.6998	0.042*	0.50
C8'	1.0406 (3)	0.5791 (14)	0.6513 (3)	0.0298 (17)	0.50
H8'	1.1027	0.5675	0.6768	0.036*	0.50
O2	0.61854 (18)	0.7401 (7)	0.75194 (14)	0.0456 (8)	
O3	0.5296 (2)	1.0621 (5)	0.65989 (17)	0.0512 (9)	
N1	0.70506 (19)	0.2448 (6)	0.44062 (14)	0.0242 (6)	
N2	0.75031 (18)	0.3840 (6)	0.50020 (14)	0.0233 (6)	
N3	0.4716 (2)	0.6093 (6)	0.67061 (15)	0.0246 (6)	
C1	0.7446 (2)	-0.0041 (8)	0.35257 (17)	0.0289 (8)	
C2	0.7692 (2)	0.1556 (7)	0.41875 (17)	0.0263 (7)	
C3	0.8543 (2)	0.2312 (7)	0.46230 (17)	0.0254 (7)	
H3	0.9094	0.1894	0.4571	0.030*	
C4	0.8404 (2)	0.3810 (7)	0.51480 (17)	0.0240 (7)	
C9	0.70348 (15)	0.4776 (5)	0.54273 (10)	0.0215 (7)	
C10	0.72944 (14)	0.3863 (5)	0.61097 (11)	0.0441 (11)	
H10	0.7778	0.2604	0.6285	0.053*	
C11	0.68462 (16)	0.4790 (6)	0.65348 (8)	0.0384 (10)	

H11	0.7024	0.4165	0.7001	0.046*
C12	0.61384 (15)	0.6630 (5)	0.62776 (10)	0.0238 (7)
C13	0.58788 (17)	0.7544 (5)	0.55953 (12)	0.0555 (14)
H13	0.5395	0.8802	0.5419	0.067*
C14	0.63270 (18)	0.6617 (5)	0.51701 (9)	0.0457 (11)
H14	0.6150	0.7242	0.4704	0.055*
H1	0.481 (2)	0.430 (3)	0.6746 (19)	0.032 (10)*
H2	0.4245 (17)	0.659 (8)	0.6352 (14)	0.041 (12)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0345 (5)	0.0213 (4)	0.0295 (5)	-0.0030 (4)	0.0161 (4)	-0.0064 (3)
F1	0.0543 (16)	0.0558 (15)	0.0475 (15)	-0.0290 (13)	0.0251 (13)	-0.0239 (12)
F2	0.094 (2)	0.0472 (14)	0.0228 (12)	-0.0098 (14)	0.0162 (13)	-0.0029 (11)
F3	0.0408 (15)	0.0808 (19)	0.0521 (16)	0.0169 (13)	0.0069 (12)	-0.0314 (14)
O1	0.012 (3)	0.026 (3)	0.025 (3)	0.009 (2)	-0.001 (2)	-0.002 (3)
C5	0.0209 (18)	0.024 (2)	0.0233 (16)	0.0009 (15)	0.0097 (14)	0.0001 (15)
C6	0.0217 (18)	0.035 (3)	0.042 (3)	0.0019 (17)	0.012 (2)	-0.011 (2)
C7	0.011 (3)	0.042 (5)	0.032 (3)	-0.003 (3)	0.002 (2)	0.001 (3)
C8	0.048 (5)	0.032 (4)	0.030 (3)	-0.008 (4)	0.020 (3)	-0.008 (3)
O1'	0.0217 (18)	0.035 (3)	0.042 (3)	0.0019 (17)	0.012 (2)	-0.011 (2)
C5'	0.0209 (18)	0.024 (2)	0.0233 (16)	0.0009 (15)	0.0097 (14)	0.0001 (15)
C6'	0.012 (3)	0.026 (3)	0.025 (3)	0.009 (2)	-0.001 (2)	-0.002 (3)
C7'	0.048 (5)	0.032 (4)	0.030 (3)	-0.008 (4)	0.020 (3)	-0.008 (3)
C8'	0.011 (3)	0.042 (5)	0.032 (3)	-0.003 (3)	0.002 (2)	0.001 (3)
O2	0.0350 (15)	0.077 (2)	0.0248 (14)	-0.0097 (14)	0.0100 (12)	-0.0201 (14)
O3	0.082 (2)	0.0163 (13)	0.079 (2)	0.0030 (13)	0.058 (2)	-0.0024 (13)
N1	0.0237 (14)	0.0303 (15)	0.0172 (14)	-0.0032 (12)	0.0054 (11)	-0.0016 (11)
N2	0.0215 (14)	0.0286 (14)	0.0202 (14)	-0.0026 (12)	0.0079 (12)	-0.0006 (11)
N3	0.0285 (16)	0.0206 (14)	0.0247 (16)	0.0029 (12)	0.0094 (13)	0.0014 (12)
C1	0.0273 (18)	0.0350 (19)	0.0242 (17)	-0.0031 (15)	0.0089 (15)	-0.0037 (15)
C2	0.0258 (18)	0.0289 (17)	0.0242 (18)	-0.0009 (14)	0.0090 (15)	-0.0004 (14)
C3	0.0226 (17)	0.0306 (18)	0.0243 (17)	-0.0025 (14)	0.0100 (14)	-0.0011 (14)
C4	0.0199 (17)	0.0304 (17)	0.0224 (17)	-0.0006 (13)	0.0084 (14)	0.0028 (14)
C9	0.0215 (16)	0.0230 (16)	0.0200 (16)	-0.0025 (13)	0.0073 (13)	0.0004 (13)
C10	0.033 (2)	0.078 (3)	0.0251 (19)	0.031 (2)	0.0155 (17)	0.020 (2)
C11	0.027 (2)	0.066 (3)	0.0211 (18)	0.0152 (19)	0.0077 (15)	0.0105 (18)
C12	0.0291 (18)	0.0192 (15)	0.0249 (18)	-0.0011 (13)	0.0118 (15)	-0.0011 (13)
C13	0.082 (4)	0.057 (3)	0.036 (2)	0.054 (3)	0.032 (2)	0.024 (2)
C14	0.071 (3)	0.046 (2)	0.028 (2)	0.033 (2)	0.026 (2)	0.0176 (18)

Geometric parameters (\AA , $^\circ$)

S1—O2	1.423 (3)	C7'—H7'	0.9500
S1—O3	1.435 (3)	C8'—H8'	0.9500
S1—N3	1.590 (3)	N1—C2	1.334 (4)
S1—C12	1.7614 (17)	N1—N2	1.362 (4)
F1—C1	1.327 (4)	N2—C4	1.369 (4)
F2—C1	1.325 (4)	N2—C9	1.416 (3)

F3—C1	1.338 (4)	N3—H1	0.878 (10)
O1—C5	1.3500	N3—H2	0.878 (10)
O1—C8	1.3500	C1—C2	1.493 (5)
C5—C6	1.3500	C2—C3	1.391 (5)
C5—C4	1.452 (5)	C3—C4	1.383 (5)
C6—C7	1.3500	C3—H3	0.9500
C6—H6	0.9500	C9—C10	1.3900
C7—C8	1.3500	C9—C14	1.3900
C7—H7	0.9500	C10—C11	1.3900
C8—H8	0.9500	C10—H10	0.9500
O1'—C5'	1.3500	C11—C12	1.3900
O1'—C8'	1.3500	C11—H11	0.9500
C5'—C6'	1.3500	C12—C13	1.3900
C5'—C4	1.446 (5)	C13—C14	1.3900
C6'—C7'	1.3500	C13—H13	0.9500
C6'—H6'	0.9500	C14—H14	0.9500
C7'—C8'	1.3500		
O2—S1—O3	120.06 (19)	S1—N3—H2	116 (3)
O2—S1—N3	108.29 (17)	H1—N3—H2	115 (4)
O3—S1—N3	105.67 (18)	F2—C1—F1	106.1 (3)
O2—S1—C12	106.65 (14)	F2—C1—F3	106.5 (3)
O3—S1—C12	106.52 (14)	F1—C1—F3	106.6 (3)
N3—S1—C12	109.37 (14)	F2—C1—C2	112.5 (3)
C5—O1—C8	108.0	F1—C1—C2	113.3 (3)
C6—C5—O1	108.0	F3—C1—C2	111.3 (3)
C6—C5—C4	129.5 (5)	N1—C2—C3	113.4 (3)
O1—C5—C4	122.5 (5)	N1—C2—C1	119.3 (3)
C5—C6—C7	108.0	C3—C2—C1	127.4 (3)
C5—C6—H6	126.0	C4—C3—C2	104.4 (3)
C7—C6—H6	126.0	C4—C3—H3	127.8
C8—C7—C6	108.0	C2—C3—H3	127.8
C8—C7—H7	126.0	N2—C4—C3	106.5 (3)
C6—C7—H7	126.0	N2—C4—C5'	127.1 (4)
C7—C8—O1	108.0	C3—C4—C5'	126.3 (4)
C7—C8—H8	126.0	N2—C4—C5	122.4 (4)
O1—C8—H8	126.0	C3—C4—C5	131.1 (5)
C5'—O1'—C8'	108.0	C10—C9—C14	120.0
O1'—C5'—C6'	108.0	C10—C9—N2	119.44 (17)
O1'—C5'—C4	122.9 (5)	C14—C9—N2	120.56 (17)
C6'—C5'—C4	129.0 (5)	C9—C10—C11	120.0
C7'—C6'—C5'	108.0	C9—C10—H10	120.0
C7'—C6'—H6'	126.0	C11—C10—H10	120.0
C5'—C6'—H6'	126.0	C12—C11—C10	120.0
C8'—C7'—C6'	108.0	C12—C11—H11	120.0
C8'—C7'—H7'	126.0	C10—C11—H11	120.0
C6'—C7'—H7'	126.0	C13—C12—C11	120.0
C7'—C8'—O1'	108.0	C13—C12—S1	120.39 (12)
C7'—C8'—H8'	126.0	C11—C12—S1	119.58 (12)

O1'—C8'—H8'	126.0	C12—C13—C14	120.0
C2—N1—N2	103.6 (3)	C12—C13—H13	120.0
N1—N2—C4	112.2 (3)	C14—C13—H13	120.0
N1—N2—C9	119.0 (3)	C13—C14—C9	120.0
C4—N2—C9	128.2 (3)	C13—C14—H14	120.0
S1—N3—H1	113 (3)	C9—C14—H14	120.0
C8—O1—C5—C6	0.0	C2—C3—C4—C5	-176.4 (7)
C8—O1—C5—C4	179.3 (12)	O1'—C5'—C4—N2	159.5 (5)
O1—C5—C6—C7	0.0	C6'—C5'—C4—N2	-22.8 (12)
C4—C5—C6—C7	-179.3 (13)	O1'—C5'—C4—C3	-18.1 (12)
C5—C6—C7—C8	0.0	C6'—C5'—C4—C3	159.7 (6)
C6—C7—C8—O1	0.0	O1'—C5'—C4—C5	-159 (10)
C5—O1—C8—C7	0.0	C6'—C5'—C4—C5	18 (9)
C8'—O1'—C5'—C6'	0.0	C6—C5—C4—N2	150.0 (6)
C8'—O1'—C5'—C4	178.2 (12)	O1—C5—C4—N2	-29.1 (12)
O1'—C5'—C6'—C7'	0.0	C6—C5—C4—C3	-33.5 (12)
C4—C5'—C6'—C7'	-178.0 (13)	O1—C5—C4—C3	147.3 (6)
C5'—C6'—C7'—C8'	0.0	C6—C5—C4—C5'	8 (9)
C6'—C7'—C8'—O1'	0.0	O1—C5—C4—C5'	-171 (10)
C5'—O1'—C8'—C7'	0.0	N1—N2—C9—C10	121.0 (3)
C2—N1—N2—C4	0.0 (3)	C4—N2—C9—C10	-49.5 (4)
C2—N1—N2—C9	-171.9 (3)	N1—N2—C9—C14	-59.7 (3)
N2—N1—C2—C3	0.3 (4)	C4—N2—C9—C14	129.8 (3)
N2—N1—C2—C1	-178.7 (3)	C14—C9—C10—C11	0.0
F2—C1—C2—N1	85.1 (4)	N2—C9—C10—C11	179.3 (2)
F1—C1—C2—N1	-35.3 (5)	C9—C10—C11—C12	0.0
F3—C1—C2—N1	-155.4 (3)	C10—C11—C12—C13	0.0
F2—C1—C2—C3	-93.8 (4)	C10—C11—C12—S1	-178.32 (19)
F1—C1—C2—C3	145.8 (4)	O2—S1—C12—C13	-159.41 (19)
F3—C1—C2—C3	25.7 (5)	O3—S1—C12—C13	-30.1 (2)
N1—C2—C3—C4	-0.5 (4)	N3—S1—C12—C13	83.70 (19)
C1—C2—C3—C4	178.4 (3)	O2—S1—C12—C11	18.9 (2)
N1—N2—C4—C3	-0.3 (4)	O3—S1—C12—C11	148.3 (2)
C9—N2—C4—C3	170.6 (3)	N3—S1—C12—C11	-98.0 (2)
N1—N2—C4—C5'	-178.3 (7)	C11—C12—C13—C14	0.0
C9—N2—C4—C5'	-7.3 (8)	S1—C12—C13—C14	178.31 (19)
N1—N2—C4—C5	176.9 (6)	C12—C13—C14—C9	0.0
C9—N2—C4—C5	-12.1 (8)	C10—C9—C14—C13	0.0
C2—C3—C4—N2	0.5 (4)	N2—C9—C14—C13	-179.3 (2)
C2—C3—C4—C5'	178.4 (7)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—H1...O3 ⁱ	0.88 (1)	2.00 (2)	2.830 (4)	158 (3)
N3—H2...N1 ⁱⁱ	0.88 (1)	2.17 (1)	3.032 (4)	170 (4)

C8—H8···O2 ⁱⁱⁱ	0.95	2.36	3.185 (7)	145
C10—H10···O2 ^{iv}	0.95	2.44	3.092 (4)	126

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, -y+1, -z+1$; (iii) $-x+3/2, y+1/2, -z+3/2$; (iv) $-x+3/2, y-1/2, -z+3/2$.