

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(5-Bromo-2-hydroxyphenyl)(4-propyl-cyclohexyl)methanone

Kang Meng, Miao Cao, Zhuo-Ling An, Jie Zhang and Li-Hong Liu*

Pharmacy Department of The Second Artillery General Hospital, Beijing 100088, People's Republic of China

Correspondence e-mail: lihongliu2011@yahoo.cn


Received 11 March 2012; accepted 17 March 2012

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.028; wR factor = 0.062; data-to-parameter ratio = 19.4.

In the title compound, $\text{C}_{16}\text{H}_{21}\text{BrO}_2$, the cyclohexane ring adopts a chair conformation. The hydroxy and carbonyl groups are involved in an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond. In the crystal, weak $\text{C}-\text{H}\cdots\text{O}$ interactions link the molecules into zigzag chains along [010].

Related literature

For details of the biological activity of SGLT2 inhibitors, see: Meng *et al.* (2008); Gao *et al.* (2010); Shao *et al.* (2011). For related structures, see: Robinson *et al.* (2002); Wang *et al.* (2011).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{21}\text{BrO}_2$
 $M_r = 325.24$
 Monoclinic, $P2_1/c$
 $a = 14.5826$ (12) Å
 $b = 8.5467$ (8) Å
 $c = 12.6369$ (10) Å
 $\beta = 113.037$ (5)°

$V = 1449.4$ (2) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 2.83$ mm⁻¹
 $T = 113$ K
 $0.20 \times 0.18 \times 0.12$ mm

Data collection

Rigaku Saturn 724 CCD area-detector diffractometer
 Absorption correction: multi-scan (*CrystalClear-SM Expert*; Rigaku/MS, 2009)
 $T_{\min} = 0.601$, $T_{\max} = 0.727$
 17988 measured reflections
 3441 independent reflections
 3069 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.062$
 $S = 1.04$
 3441 reflections
 177 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.81$ e Å⁻³
 $\Delta\rho_{\min} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{O2}$	0.81 (2)	1.82 (2)	2.5527 (16)	148 (3)
$\text{C3}-\text{H3}\cdots\text{O1}^i$	0.95	2.59	3.483 (2)	157

Symmetry code: (i) $-x + 2, y - \frac{1}{2}, -z - \frac{1}{2}$.

Data collection: *CrystalClear-SM Expert* (Rigaku/MS, 2009); cell refinement: *CrystalClear-SM Expert*; data reduction: *CrystalClear-SM Expert*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Dr Haibin Song, Nankai University, for the X-ray crystallographic determination.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5261).

References

- Gao, Y. L., Zhao, G. L., Liu, W., Wang, Y. L., Xu, W. R. & Wang, J. W. (2010). *Chin. J. Chem.* **28**, 605–612.
- Meng, M., Ellsworth, B. A., Nirschl, A. A., McCann, P. J., Patel, M., Girotra, R. N., Wu, G., Sher, P. M., Morrison, E. P., Biller, S. A., Zahler, R., Deshpande, P. P., Pullockaran, A., Hagan, D. L., Morgan, N., Taylor, J. R., Obermeier, M. T., Humphreys, W. G., Khanna, A., Discenza, L., Robertson, J. M., Wang, A., Han, S., Wetterau, J. R., Janovitz, E. B., Flint, O. P., Whaley, J. M. & Washburn, W. N. (2008). *J. Med. Chem.* **51**, 1145–1149.
- Rigaku/MS (2009). *CrystalClear-SM Expert*. Rigaku/MS, The Woodlands, Texas, USA.
- Robinson, P. D., Meyers, C. Y. & Kolb, V. M. (2002). *Acta Cryst.* **E58**, o1288–o1289.
- Shao, H., Gao, Y. L., Lou, Y. Y., Wang, Y. L., Liu, W., Xu, W. R., Wang, J. W., Zhao, G. L. & Tang, L. D. (2011). *Chin. J. Org. Chem.* **31**, 836–842.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, L., Chang, Z., Ding, C., Shao, H. & Sun, J. (2011). *Acta Cryst.* **E67**, o1173.

supplementary materials

Acta Cryst. (2012). E68, o1181 [doi:10.1107/S1600536812011634]

(5-Bromo-2-hydroxyphenyl)(4-propylcyclohexyl)methanone**Kang Meng, Miao Cao, Zhuo-Ling An, Jie Zhang and Li-Hong Liu****Comment**

SGLT2 inhibitors represent a new class of potential hypoglycemic agents (Meng *et al.*, 2008). During the development of our own cyclohexane-bearing SGLT2 inhibitors (Gao *et al.*, 2010; Shao *et al.*, 2011), the title compound (I) was prepared as a key intermediate.

In (I) (Fig. 1), all bond lengths and angles are normal and in a good agreement with those reported previously for related compounds (Robinson *et al.*, 2002; Wang *et al.*, 2011). The cyclohexane ring (C8—C13) adopts a chair conformation. Weak intermolecular C—H \cdots O interactions (Table 1) link the molecules into zigzag chains in [010].

Experimental


A dried 100-ml round-bottomed flask was charged with 1.89 g (10 mmol) of *trans*-4-propylcyclohexanecarboxylic acid chloride, 1.87 g (10 mmol) of 4-bromoanisole and 20 ml of dried dichloromethane. The mixture was stirred on an ice-water bath, followed by addition of 1.60 g (12 mmol) of anhydrous aluminium chloride in a portionwise manner. After addition, the reaction mixture was stirred at room temperature for 1 h and at reflux overnight, and poured into 300 ml of ice-water. The mixture thus formed was exacted with three 50-ml portions of dichloromethane, and the combined exacts were washed with saturated brine, dried over sodium sulfate and evaporated on a rotary evaporator to afford the crude title compound. Pure title compound was obtained by column chromatography. Crystals suitable for X-ray diffraction were obtained through slow evaporation of a solution of the pure title compound in dichloromethane/petroleum ether (1/30 by volume).

Refinement

Hydroxy atom H1 was located on a difference map and isotropically refined. C-bound H atoms were geometrically positioned [C—H = 0.95–1.00 Å], and included in the final cycles of refinement using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{C})$ for the methyl H atoms.

Computing details

Data collection: *CrystalClear-SM Expert* (Rigaku/MSK, 2009); cell refinement: *CrystalClear-SM Expert* (Rigaku/MSK, 2009); data reduction: *CrystalClear-SM Expert* (Rigaku/MSK, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).


Figure 1

View of the title compound, with displacement ellipsoids drawn at the 40% probability level.

(5-Bromo-2-hydroxyphenyl)(4-propylcyclohexyl)methanone
Crystal data
 $C_{16}H_{21}BrO_2$
 $M_r = 325.24$

 Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 14.5826$ (12) Å

 $b = 8.5467$ (8) Å

 $c = 12.6369$ (10) Å

 $\beta = 113.037$ (5)°

 $V = 1449.4$ (2) Å³
 $Z = 4$
 $F(000) = 672$
 $D_x = 1.490$ Mg m⁻³

 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5072 reflections

 $\theta = 1.8$ – 27.9 °

 $\mu = 2.83$ mm⁻¹
 $T = 113$ K

Prism, colourless

 $0.20 \times 0.18 \times 0.12$ mm

Data collection

 Rigaku Saturn 724 CCD area-detector
diffractometer

Radiation source: rotating anode

Multilayer monochromator

 Detector resolution: 14.22 pixels mm⁻¹
 ω and φ scans

Absorption correction: multi-scan

 (*CrystalClear-SM Expert*; Rigaku/MS, 2009)

 $T_{\min} = 0.601$, $T_{\max} = 0.727$

17988 measured reflections

3441 independent reflections

 3069 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$
 $\theta_{\max} = 27.9$ °, $\theta_{\min} = 2.8$ °

 $h = -19$ → 19
 $k = -11$ → 11
 $l = -16$ → 16
Refinement

 Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.062$
 $S = 1.04$

3441 reflections

177 parameters

0 restraints

 Primary atom site location: structure-invariant
direct methods

 Secondary atom site location: difference Fourier
map

 Hydrogen site location: inferred from
neighbouring sites

 H atoms treated by a mixture of independent
and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0334P)^2 + 0.0947P]$

 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.81$ e Å⁻³
 $\Delta\rho_{\min} = -0.34$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.101943 (12)	0.65152 (2)	0.057741 (15)	0.02149 (7)
O1	0.80421 (9)	1.03963 (14)	-0.31892 (10)	0.0200 (3)
H1	0.759 (2)	1.066 (3)	-0.301 (2)	0.058 (8)*
O2	0.69906 (9)	1.05226 (14)	-0.19829 (10)	0.0202 (3)
C1	0.92537 (13)	0.83440 (18)	-0.04336 (14)	0.0169 (3)
H1A	0.9181	0.8160	0.0271	0.020*
C2	1.00541 (12)	0.77214 (19)	-0.06037 (14)	0.0174 (3)
C3	1.01921 (13)	0.8002 (2)	-0.16215 (15)	0.0197 (4)
H3	1.0747	0.7565	-0.1733	0.024*
C4	0.95127 (13)	0.8920 (2)	-0.24620 (15)	0.0193 (4)
H4	0.9610	0.9135	-0.3148	0.023*
C5	0.86871 (13)	0.95348 (19)	-0.23174 (14)	0.0170 (3)
C6	0.85425 (12)	0.92496 (19)	-0.12913 (14)	0.0153 (3)
C7	0.76554 (12)	0.99105 (19)	-0.11519 (14)	0.0157 (3)
C8	0.75600 (12)	0.98403 (19)	-0.00005 (13)	0.0153 (3)
H8	0.8240	0.9946	0.0623	0.018*
C9	0.69103 (13)	1.11868 (19)	0.01104 (14)	0.0174 (4)
H9A	0.7221	1.2196	0.0053	0.021*
H9B	0.6247	1.1133	-0.0530	0.021*
C10	0.67844 (13)	1.11146 (19)	0.12561 (14)	0.0169 (4)
H10A	0.7440	1.1282	0.1893	0.020*
H10B	0.6335	1.1968	0.1283	0.020*
C11	0.63570 (12)	0.95477 (18)	0.14288 (13)	0.0154 (3)
H11	0.5675	0.9435	0.0812	0.018*
C12	0.70000 (13)	0.82152 (19)	0.12971 (14)	0.0174 (4)
H12A	0.6697	0.7205	0.1368	0.021*
H12B	0.7668	0.8276	0.1928	0.021*
C13	0.71144 (13)	0.82583 (19)	0.01480 (14)	0.0164 (3)
H13A	0.6456	0.8109	-0.0487	0.020*
H13B	0.7556	0.7396	0.0116	0.020*
C14	0.62659 (13)	0.9434 (2)	0.25955 (14)	0.0183 (4)
H14A	0.5998	0.8389	0.2656	0.022*
H14B	0.6942	0.9509	0.3211	0.022*
C15	0.56089 (13)	1.0669 (2)	0.28140 (15)	0.0202 (4)
H15A	0.4955	1.0692	0.2156	0.024*
H15B	0.5923	1.1709	0.2869	0.024*
C16	0.54478 (14)	1.0347 (2)	0.39195 (15)	0.0247 (4)

H16A	0.5087	0.9359	0.3843	0.037*
H16B	0.5059	1.1201	0.4056	0.037*
H16C	0.6095	1.0276	0.4568	0.037*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.01672 (10)	0.02144 (10)	0.02637 (11)	0.00224 (7)	0.00849 (8)	0.00249 (7)
O1	0.0228 (7)	0.0202 (6)	0.0201 (6)	0.0024 (5)	0.0115 (6)	0.0036 (5)
O2	0.0184 (6)	0.0240 (6)	0.0187 (6)	0.0029 (5)	0.0077 (5)	0.0018 (5)
C1	0.0178 (8)	0.0165 (8)	0.0184 (8)	-0.0028 (7)	0.0094 (7)	-0.0016 (7)
C2	0.0158 (8)	0.0140 (8)	0.0215 (9)	-0.0014 (7)	0.0063 (7)	-0.0009 (7)
C3	0.0195 (9)	0.0169 (8)	0.0267 (9)	-0.0027 (7)	0.0135 (8)	-0.0054 (7)
C4	0.0225 (9)	0.0189 (8)	0.0213 (9)	-0.0041 (7)	0.0139 (8)	-0.0025 (7)
C5	0.0202 (9)	0.0138 (8)	0.0176 (8)	-0.0040 (7)	0.0079 (7)	-0.0033 (7)
C6	0.0168 (8)	0.0129 (8)	0.0173 (8)	-0.0028 (7)	0.0080 (7)	-0.0023 (6)
C7	0.0163 (8)	0.0130 (8)	0.0185 (8)	-0.0029 (6)	0.0075 (7)	-0.0016 (7)
C8	0.0142 (8)	0.0163 (8)	0.0162 (8)	-0.0005 (7)	0.0069 (7)	-0.0006 (6)
C9	0.0200 (9)	0.0151 (8)	0.0201 (9)	0.0005 (7)	0.0112 (7)	0.0011 (7)
C10	0.0195 (9)	0.0148 (8)	0.0199 (9)	0.0009 (7)	0.0115 (7)	-0.0007 (7)
C11	0.0155 (8)	0.0158 (8)	0.0150 (8)	0.0009 (6)	0.0061 (7)	0.0003 (7)
C12	0.0192 (9)	0.0152 (8)	0.0198 (9)	0.0020 (7)	0.0098 (7)	0.0039 (7)
C13	0.0179 (8)	0.0146 (8)	0.0185 (8)	0.0006 (6)	0.0089 (7)	-0.0017 (7)
C14	0.0194 (9)	0.0187 (8)	0.0181 (8)	-0.0002 (7)	0.0089 (7)	0.0014 (7)
C15	0.0205 (9)	0.0218 (9)	0.0202 (9)	0.0020 (7)	0.0100 (7)	0.0006 (7)
C16	0.0285 (10)	0.0260 (10)	0.0245 (9)	-0.0005 (8)	0.0157 (8)	-0.0008 (8)

Geometric parameters (\AA , $^\circ$)

Br1—C2	1.9046 (17)	C10—C11	1.528 (2)
O1—C5	1.353 (2)	C10—H10A	0.9900
O1—H1	0.81 (2)	C10—H10B	0.9900
O2—C7	1.233 (2)	C11—C12	1.525 (2)
C1—C2	1.374 (2)	C11—C14	1.534 (2)
C1—C6	1.403 (2)	C11—H11	1.0000
C1—H1A	0.9500	C12—C13	1.525 (2)
C2—C3	1.398 (2)	C12—H12A	0.9900
C3—C4	1.379 (2)	C12—H12B	0.9900
C3—H3	0.9500	C13—H13A	0.9900
C4—C5	1.390 (2)	C13—H13B	0.9900
C4—H4	0.9500	C14—C15	1.522 (2)
C5—C6	1.414 (2)	C14—H14A	0.9900
C6—C7	1.484 (2)	C14—H14B	0.9900
C7—C8	1.516 (2)	C15—C16	1.530 (2)
C8—C9	1.531 (2)	C15—H15A	0.9900
C8—C13	1.543 (2)	C15—H15B	0.9900
C8—H8	1.0000	C16—H16A	0.9800
C9—C10	1.530 (2)	C16—H16B	0.9800
C9—H9A	0.9900	C16—H16C	0.9800
C9—H9B	0.9900		

C5—O1—H1	107.5 (19)	C9—C10—H10B	109.2
C2—C1—C6	120.60 (15)	H10A—C10—H10B	107.9
C2—C1—H1A	119.7	C12—C11—C10	109.65 (13)
C6—C1—H1A	119.7	C12—C11—C14	110.22 (13)
C1—C2—C3	120.96 (16)	C10—C11—C14	112.72 (13)
C1—C2—Br1	119.98 (13)	C12—C11—H11	108.0
C3—C2—Br1	119.03 (13)	C10—C11—H11	108.0
C4—C3—C2	119.17 (16)	C14—C11—H11	108.0
C4—C3—H3	120.4	C11—C12—C13	112.75 (13)
C2—C3—H3	120.4	C11—C12—H12A	109.0
C3—C4—C5	120.81 (16)	C13—C12—H12A	109.0
C3—C4—H4	119.6	C11—C12—H12B	109.0
C5—C4—H4	119.6	C13—C12—H12B	109.0
O1—C5—C4	117.44 (15)	H12A—C12—H12B	107.8
O1—C5—C6	122.35 (15)	C12—C13—C8	110.19 (13)
C4—C5—C6	120.21 (16)	C12—C13—H13A	109.6
C1—C6—C5	118.23 (15)	C8—C13—H13A	109.6
C1—C6—C7	122.23 (15)	C12—C13—H13B	109.6
C5—C6—C7	119.54 (15)	C8—C13—H13B	109.6
O2—C7—C6	119.47 (14)	H13A—C13—H13B	108.1
O2—C7—C8	119.83 (14)	C15—C14—C11	115.33 (14)
C6—C7—C8	120.70 (14)	C15—C14—H14A	108.4
C7—C8—C9	110.51 (13)	C11—C14—H14A	108.4
C7—C8—C13	110.65 (13)	C15—C14—H14B	108.4
C9—C8—C13	110.01 (13)	C11—C14—H14B	108.4
C7—C8—H8	108.5	H14A—C14—H14B	107.5
C9—C8—H8	108.5	C14—C15—C16	111.95 (14)
C13—C8—H8	108.5	C14—C15—H15A	109.2
C10—C9—C8	111.34 (13)	C16—C15—H15A	109.2
C10—C9—H9A	109.4	C14—C15—H15B	109.2
C8—C9—H9A	109.4	C16—C15—H15B	109.2
C10—C9—H9B	109.4	H15A—C15—H15B	107.9
C8—C9—H9B	109.4	C15—C16—H16A	109.5
H9A—C9—H9B	108.0	C15—C16—H16B	109.5
C11—C10—C9	112.15 (13)	H16A—C16—H16B	109.5
C11—C10—H10A	109.2	C15—C16—H16C	109.5
C9—C10—H10A	109.2	H16A—C16—H16C	109.5
C11—C10—H10B	109.2	H16B—C16—H16C	109.5
C6—C1—C2—C3	1.4 (3)	O2—C7—C8—C9	-27.0 (2)
C6—C1—C2—Br1	179.29 (12)	C6—C7—C8—C9	152.95 (14)
C1—C2—C3—C4	0.1 (3)	O2—C7—C8—C13	95.07 (18)
Br1—C2—C3—C4	-177.77 (13)	C6—C7—C8—C13	-84.95 (18)
C2—C3—C4—C5	-1.4 (3)	C7—C8—C9—C10	178.66 (13)
C3—C4—C5—O1	-178.60 (15)	C13—C8—C9—C10	56.18 (18)
C3—C4—C5—C6	1.2 (3)	C8—C9—C10—C11	-56.10 (18)
C2—C1—C6—C5	-1.6 (2)	C9—C10—C11—C12	54.50 (18)
C2—C1—C6—C7	178.51 (15)	C9—C10—C11—C14	177.69 (14)

O1—C5—C6—C1	-179.88 (15)	C10—C11—C12—C13	-55.68 (18)
C4—C5—C6—C1	0.3 (2)	C14—C11—C12—C13	179.67 (14)
O1—C5—C6—C7	0.0 (2)	C11—C12—C13—C8	57.25 (18)
C4—C5—C6—C7	-179.81 (15)	C7—C8—C13—C12	-178.73 (13)
C1—C6—C7—O2	-170.67 (15)	C9—C8—C13—C12	-56.33 (17)
C5—C6—C7—O2	9.5 (2)	C12—C11—C14—C15	-177.88 (14)
C1—C6—C7—C8	9.3 (2)	C10—C11—C14—C15	59.25 (19)
C5—C6—C7—C8	-170.52 (15)	C11—C14—C15—C16	172.51 (15)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...O2	0.81 (2)	1.82 (2)	2.5527 (16)	148 (3)
C3—H3...O1 ⁱ	0.95	2.59	3.483 (2)	157

Symmetry code: (i) $-x+2, y-1/2, -z-1/2$.