

Bis(3-acetylpyridine- κN)diaquabis(selenocyanato- κN)cobalt(II)

Julia Werner,* Jan Boeckmann, Inke Jess and Christian Näther

Institut für Anorganische Chemie, Christian-Albrechts-Universität Kiel, Max-Eyth-Strasse 2, 24118 Kiel, Germany

Correspondence e-mail: jwerner@ac.uni-kiel.de


Received 23 April 2012; accepted 24 April 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.029; wR factor = 0.066; data-to-parameter ratio = 19.3.

In the crystal structure of the title compound, $[\text{Co}(\text{NCSe})_2(\text{C}_7\text{H}_7\text{NO})_2(\text{H}_2\text{O})_2]$, the Co^{2+} cation is coordinated by two selenocyanate anions, two 3-acetylpyridine ligands and two water molecules within a slightly distorted CoN_4O_2 octahedron. The asymmetric unit consists of one Co^{2+} cation, which is located on a center of inversion, as well as one selenocyanate anion, one 3-acetylpyridine ligand and one water molecule in general positions. Whereas one of the water H atoms makes a classical $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond, the other shows a $\text{O}-\text{H}\cdots\text{Se}$ interaction.

Related literature

For general background to this work, see: Näther & Greve (2003). For the synthesis, structures and properties of the corresponding compounds with pyridine, see: Boeckmann & Näther (2010, 2011, 2012).



Experimental

Crystal data

$[\text{Co}(\text{NCSe})_2(\text{C}_7\text{H}_7\text{NO})_2(\text{H}_2\text{O})_2]$	$V = 2025.13$ (13) Å ³
$M_r = 547.19$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 19.1098$ (6) Å	$\mu = 4.47$ mm ⁻¹
$b = 9.0064$ (4) Å	$T = 293$ K
$c = 14.9734$ (5) Å	$0.35 \times 0.27 \times 0.19$ mm
$\beta = 128.203$ (2)°	

Data collection

Stoe IPDS-2 diffractometer	16689 measured reflections
Absorption correction: numerical (<i>X-SHAPE</i> and <i>X-RED32</i> ; Stoe & Cie, 2008)	2409 independent reflections
$T_{\min} = 0.240$, $T_{\max} = 0.423$	2260 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$	125 parameters
$wR(F^2) = 0.066$	H-atom parameters constrained
$S = 1.12$	$\Delta\rho_{\max} = 0.50$ e Å ⁻³
2409 reflections	$\Delta\rho_{\min} = -0.45$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1O}\cdots\text{O11}^{\text{i}}$	0.82	1.97	2.780 (2)	169
$\text{O1}-\text{H2O}\cdots\text{Se1}^{\text{ii}}$	0.82	2.57	3.338 (2)	157

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2008); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008) and *DIAMOND* (Brandenburg, 2011); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

We gratefully acknowledge financial support by the DFG (project No. NA 720/3-1) and the State of Schleswig-Holstein. We thank Professor Dr Wolfgang Bensch for access to his experimental facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5898).

References

- Boeckmann, J. & Näther, C. (2010). *Dalton Trans.* **39**, 11019–11026.
 Boeckmann, J. & Näther, C. (2011). *Chem. Commun.* **47**, 7104–7106.
 Boeckmann, J. & Näther, C. (2012). *Polyhedron*, **31**, 587–595.
 Brandenburg, K. (2011). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Näther, C. & Greve, J. (2003). *J. Solid State Chem.* **176**, 259–265.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Stoe & Cie (2008). *X-AREA*, *X-RED32* and *X-SHAPE*. Stoe & Cie, Darmstadt, Germany.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, m703 [doi:10.1107/S1600536812018387]

Bis(3-acetylpyridine- κ N)diaquabis(selenocyanato- κ N)cobalt(II)

Julia Werner, Jan Boeckmann, Inke Jess and Christian Näther

Comment

Recently we have reported on the magnetic properties of 1D coordination polymers based on paramagnetic transition metal cations that are coordinated by pyridine as co-ligand and that are μ -1,3-bridged into chains by thio- or selenocyanato anions (Näther & Greve, 2003). Dependent on the nature of the metal cation, anti- or ferromagnetic ordering is observed and for the compounds with Co(II) as cation we have found a slow relaxation of the magnetization (Boeckmann & Näther 2010, 2011 and 2012). To investigate the influence of the co-ligand on the magnetic properties we tried to prepare similar compounds based on 3-acetylpyridine, which resulted in the formation of the title compound in which the anionic ligands are only terminal N-coordinated. Further investigations also show that this compound cannot be transformed into the corresponding 1D coordination polymers and therefore, it was characterized only by single crystal X-ray diffraction.

In the crystal structure the cobalt(II) cations are coordinated by four nitrogen atoms of two terminal N-bonded selenocyanato anions and two terminal bonded 3-acetylpyridine co-ligands as well as two water molecules into discrete complexes (Fig. 1). The coordination polyhedron of the Co cations can be described as a slightly distorted octahedron with the Co cation located on a centre of inversion. The discrete cobalt complexes are bridged by two pairs of intermolecular O—H \cdots O hydrogen bonding into 16-membered rings that are located on centres of inversion (Fig. 2). These rings are further linked into hydrogen bonded chains that are parallel to the *b* axis (Fig. 2 and Table 1).

Experimental


Potassium selenocyanate and 3-acetylpyridine were purchased from Alfa Aesar and the Co(NO₃)₂·6 H₂O obtained from Merck. The title compound was prepared by the reaction of 72.8 mg Co(NO₃)₂·6 H₂O (0.25 mmol), 64.8 mg KSeCN (0.45 mmol) and 54.6 μ L 3-acetylpyridine (0.50 mmol) in 1.5 mL H₂O at RT in a closed 3 ml snap cap vial. After three days pink blocks of the title compound were obtained.

Refinement

The C-H H atoms were positioned with idealized geometry (methyl H atoms allowed to rotate but not to tip) and were refined isotropic with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ for aromatic H atoms (1.5 for methyl H atoms) using a riding model with C—H = 0.93 Å (aromatic) and with C—H = 0.96 Å (methyl). The O-H H atom were located in difference map, their bond lengths set to ideal values of 0.84 Å and afterwards they were refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$.


Computing details

Data collection: *X-AREA* (Stoe & Cie, 2008); cell refinement: *X-AREA* (Stoe & Cie, 2008); data reduction: *X-AREA* (Stoe & Cie, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008) and *DIAMOND* (Brandenburg, 2011); software used to prepare material for publication: *publCIF* (Westrip, 2010).

**Figure 1**

Crystal structure of the title compound with labelling and displacement ellipsoids drawn at the 50% probability level.

Symmetry code: $i = -x+1, -y+1, -z$.


Figure 2

Crystal structure showing the discrete complexes that are linked by hydrogen bonding into chains that elongate in the direction of the crystallographic *b* axis. Hydrogen bonding is shown as dashed lines and for clarification only the O-H H atoms are shown.

Bis(3-acetylpyridine- κ N)diaquabis(selenocyanato- κ N)cobalt(II)
Crystal data
 $[\text{Co}(\text{NCSe})_2(\text{C}_7\text{H}_7\text{NO})_2(\text{H}_2\text{O})_2]$
 $M_r = 547.19$

 Monoclinic, $C2/c$

 Hall symbol: $-C\ 2yc$
 $a = 19.1098\ (6)\ \text{\AA}$
 $b = 9.0064\ (4)\ \text{\AA}$
 $c = 14.9734\ (5)\ \text{\AA}$
 $\beta = 128.203\ (2)^\circ$
 $V = 2025.13\ (13)\ \text{\AA}^3$
 $Z = 4$
 $F(000) = 1076$
 $D_x = 1.795\ \text{Mg m}^{-3}$

 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 16689 reflections

 $\theta = 2.6\text{--}27.9^\circ$
 $\mu = 4.47\ \text{mm}^{-1}$
 $T = 293\ \text{K}$

Block, pink

 $0.35 \times 0.27 \times 0.19\ \text{mm}$
Data collection

Stoe IPDS-2

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scans

Absorption correction: numerical

 (*X-SHAPE* and *X-RED32*; Stoe & Cie, 2008)

 $T_{\min} = 0.240$, $T_{\max} = 0.423$

16689 measured reflections

2409 independent reflections

 2260 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\max} = 27.9^\circ$, $\theta_{\min} = 2.6^\circ$
 $h = -25 \rightarrow 24$
 $k = -11 \rightarrow 11$
 $l = -19 \rightarrow 19$
Refinement

 Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.066$
 $S = 1.12$

2409 reflections

125 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0275P)^2 + 2.5784P]$

 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.50\ \text{e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.45\ \text{e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.5000	0.5000	0.0000	0.03364 (10)
Se1	0.725951 (16)	0.09800 (3)	0.20771 (2)	0.05020 (9)
C1	0.65776 (13)	0.2600 (2)	0.13699 (18)	0.0372 (4)
N1	0.61139 (12)	0.3598 (2)	0.09124 (17)	0.0431 (4)
O1	0.55451 (11)	0.64233 (17)	0.14168 (13)	0.0450 (4)
H1O	0.5265	0.7200	0.1249	0.068*
H2O	0.6080	0.6587	0.1792	0.068*
N11	0.43968 (12)	0.36819 (19)	0.05628 (16)	0.0393 (4)
C11	0.44407 (14)	0.2201 (2)	0.06235 (18)	0.0373 (4)
H11	0.4741	0.1709	0.0406	0.045*
C12	0.40582 (14)	0.1366 (2)	0.09965 (17)	0.0370 (4)
C13	0.36145 (18)	0.2091 (3)	0.1322 (2)	0.0499 (6)
H13	0.3339	0.1558	0.1560	0.060*
C14	0.3585 (2)	0.3627 (3)	0.1289 (3)	0.0574 (7)
H14	0.3303	0.4146	0.1524	0.069*
C15	0.39775 (18)	0.4364 (3)	0.0905 (2)	0.0497 (6)
H15	0.3952	0.5395	0.0880	0.060*
C16	0.41692 (16)	-0.0284 (2)	0.10701 (19)	0.0423 (5)
C17	0.3525 (2)	-0.1217 (3)	0.1065 (3)	0.0614 (7)
H17A	0.2945	-0.1129	0.0342	0.092*
H17B	0.3500	-0.0890	0.1655	0.092*
H17C	0.3713	-0.2236	0.1194	0.092*
O11	0.47857 (13)	-0.08133 (19)	0.11398 (17)	0.0558 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.03083 (19)	0.02384 (18)	0.0412 (2)	0.00276 (13)	0.01973 (16)	0.00073 (14)
Se1	0.04556 (14)	0.04665 (15)	0.06007 (16)	0.01964 (10)	0.03351 (12)	0.01574 (11)
C1	0.0309 (9)	0.0380 (10)	0.0403 (11)	0.0027 (8)	0.0208 (9)	-0.0010 (8)
N1	0.0335 (9)	0.0369 (9)	0.0504 (10)	0.0059 (7)	0.0217 (8)	0.0035 (8)
O1	0.0406 (8)	0.0337 (8)	0.0491 (9)	0.0013 (6)	0.0218 (7)	-0.0061 (6)
N11	0.0400 (9)	0.0303 (9)	0.0499 (10)	0.0021 (7)	0.0289 (8)	0.0011 (7)
C11	0.0387 (10)	0.0302 (10)	0.0435 (11)	0.0021 (8)	0.0258 (9)	-0.0007 (8)
C12	0.0389 (10)	0.0306 (10)	0.0382 (10)	0.0001 (8)	0.0222 (9)	0.0004 (8)
C13	0.0598 (14)	0.0450 (13)	0.0603 (14)	-0.0013 (11)	0.0449 (13)	-0.0002 (11)
C14	0.0723 (18)	0.0462 (14)	0.0800 (18)	0.0040 (13)	0.0603 (16)	-0.0056 (13)

C15	0.0586 (15)	0.0316 (11)	0.0690 (16)	0.0030 (10)	0.0445 (14)	-0.0046 (10)
C16	0.0500 (12)	0.0330 (10)	0.0390 (10)	0.0001 (9)	0.0250 (10)	0.0013 (8)
C17	0.0768 (19)	0.0403 (13)	0.0705 (18)	-0.0095 (13)	0.0473 (16)	0.0032 (12)
O11	0.0598 (11)	0.0342 (8)	0.0701 (12)	0.0088 (8)	0.0386 (10)	0.0034 (8)

Geometric parameters (Å, °)

Co1—N1 ⁱ	2.0962 (18)	C11—H11	0.9300
Co1—N1	2.0962 (18)	C12—C13	1.376 (3)
Co1—O1 ⁱ	2.1202 (15)	C12—C16	1.495 (3)
Co1—O1	2.1202 (15)	C13—C14	1.384 (4)
Co1—N11	2.1562 (19)	C13—H13	0.9300
Co1—N11 ⁱ	2.1562 (19)	C14—C15	1.367 (4)
Se1—C1	1.800 (2)	C14—H14	0.9300
C1—N1	1.146 (3)	C15—H15	0.9300
O1—H1O	0.8201	C16—O11	1.214 (3)
O1—H2O	0.8200	C16—C17	1.487 (4)
N11—C11	1.336 (3)	C17—H17A	0.9600
N11—C15	1.338 (3)	C17—H17B	0.9600
C11—C12	1.384 (3)	C17—H17C	0.9600
N1 ⁱ —Co1—N1	180.00 (8)	N11—C11—H11	118.6
N1 ⁱ —Co1—O1 ⁱ	92.26 (7)	C12—C11—H11	118.6
N1—Co1—O1 ⁱ	87.74 (7)	C13—C12—C11	118.7 (2)
N1 ⁱ —Co1—O1	87.74 (7)	C13—C12—C16	122.4 (2)
N1—Co1—O1	92.26 (7)	C11—C12—C16	118.9 (2)
O1 ⁱ —Co1—O1	180.00 (9)	C12—C13—C14	118.9 (2)
N1 ⁱ —Co1—N11	90.77 (7)	C12—C13—H13	120.6
N1—Co1—N11	89.23 (7)	C14—C13—H13	120.6
O1 ⁱ —Co1—N11	90.43 (7)	C15—C14—C13	118.6 (2)
O1—Co1—N11	89.57 (7)	C15—C14—H14	120.7
N1 ⁱ —Co1—N11 ⁱ	89.23 (7)	C13—C14—H14	120.7
N1—Co1—N11 ⁱ	90.77 (7)	N11—C15—C14	123.6 (2)
O1 ⁱ —Co1—N11 ⁱ	89.57 (7)	N11—C15—H15	118.2
O1—Co1—N11 ⁱ	90.43 (7)	C14—C15—H15	118.2
N11—Co1—N11 ⁱ	180.00 (9)	O11—C16—C17	122.3 (2)
N1—C1—Se1	177.1 (2)	O11—C16—C12	118.8 (2)
C1—N1—Co1	163.86 (18)	C17—C16—C12	118.8 (2)
Co1—O1—H1O	112.7	C16—C17—H17A	109.5
Co1—O1—H2O	115.2	C16—C17—H17B	109.5
H1O—O1—H2O	110.9	H17A—C17—H17B	109.5
C11—N11—C15	117.4 (2)	C16—C17—H17C	109.5
C11—N11—Co1	123.37 (15)	H17A—C17—H17C	109.5
C15—N11—Co1	119.25 (15)	H17B—C17—H17C	109.5
N11—C11—C12	122.9 (2)		

Symmetry code: (i) $-x+1, -y+1, -z$.

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1O \cdots O11 ⁱⁱ	0.82	1.97	2.780 (2)	169
O1—H2O \cdots Se1 ⁱⁱⁱ	0.82	2.57	3.338 (2)	157

Symmetry codes: (ii) $x, y+1, z$; (iii) $-x+3/2, y+1/2, -z+1/2$.