

2-{4-[Acetyl(ethyl)amino]benzene-sulfonamido}benzoic acid

Ghulam Mustafa,^a Islam Ullah Khan,^a Farhan Mehmood Khan^a and Mehmet Akkurt^{b*}

^aDepartment of Chemistry, GC University, Lahore 54000, Pakistan, and ^bDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey
Correspondence e-mail: akkurt@erciyes.edu.tr, gmustafa884@yahoo.com


Received 30 March 2012; accepted 30 March 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.035; wR factor = 0.093; data-to-parameter ratio = 16.7.

In the title compound, $\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}_5\text{S}$, the dihedral angle between the aromatic rings is $68.59(10)^\circ$ and the $\text{C}-\text{S}-\text{N}-\text{C}$ torsion angle is $-81.84(18)^\circ$. The molecular conformation is stabilized by an intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond, generating an $S(6)$ ring. In the crystal, molecules are linked by $\text{C}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds into a three-dimensional network.

Related literature

For related structures and background to the biological properties of sulfonamides, see: Mustafa *et al.* (2010, 2011); Khan *et al.* (2011).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}_5\text{S}$

$M_r = 362.40$

Orthorhombic, $Pna2_1$

$a = 18.0371(7)\text{ \AA}$

$b = 12.0249(4)\text{ \AA}$

$c = 7.8430(2)\text{ \AA}$

$V = 1701.10(10)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.22\text{ mm}^{-1}$

$T = 296\text{ K}$
 $0.35 \times 0.29 \times 0.27\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer
9278 measured reflections

3880 independent reflections
3327 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.093$
 $S = 0.99$
3880 reflections
233 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.35\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.18\text{ e \AA}^{-3}$
Absolute structure: Flack (1983), 1612 Freidel pairs
Flack parameter: 0.05 (6)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots O5 ⁱ	0.82	1.82	2.599 (2)	158
N1—H1N \cdots O2	0.85 (2)	1.93 (2)	2.627 (2)	138 (2)
C12—H12 \cdots O3 ⁱⁱ	0.93	2.45	3.356 (2)	164
C15—H15C \cdots O3 ⁱⁱ	0.96	2.53	3.400 (3)	150
C17—H17A \cdots O2 ⁱⁱ	0.96	2.58	3.486 (4)	158

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, z + \frac{3}{2}$; (ii) $x, y, z - 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors are grateful to Mr Muhammad Nadeem Arshad for his assistance and the Higher Education Commission (HEC), Pakistan, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6720).

References

- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Khan, I. U., Mustafa, G. & Akkurt, M. (2011). *Acta Cryst. E* **67**, o1857.
- Mustafa, G., Akkurt, M., Khan, I. U., Naseem, R. & Sajjad, B. (2010). *Acta Cryst. E* **66**, o1768.
- Mustafa, G., Khan, I. U., Zia-ur-Rehman, M., Sharif, S. & Arshad, M. N. (2011). *Acta Cryst. E* **67**, o1018.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2012). E68, o1305 [doi:10.1107/S1600536812013864]

2-{4-[Acetyl(ethyl)amino]benzenesulfonamido}benzoic acid

Ghulam Mustafa, Islam Ullah Khan, Farhan Mehmood Khan and Mehmet Akkurt

Comment

As part of our ongoing studies of sulfonamides with potential biological properties (Mustafa *et al.*, 2010, 2011; Khan *et al.*, 2011), we now describe the title compound, (I).

In the title compound (I), (Fig. 1), the C2—C7 benzene ring makes a dihedral angle of 68.59 (10) $^{\circ}$ with the C8—C13 benzene ring. All the bond lengths are similar to those of the related molecules (Mustafa *et al.*, 2010; 2011; Khan *et al.*, 2011). The torsion angles C3—N1—S1—C8, C11—N2—C16—C17 and C11—N2—C14—C15 are -81.84 (18), 94.2 (3), and 7.0 (3) $^{\circ}$, respectively.

The molecular conformation of (I) is stabilized by intramolecular N—H \cdots O hydrogen-bond interaction (Table 1), which generate one S(6) ring. In the crystal, molecules are linked by C—H \cdots O and O—H \cdots O hydrogen bonds to form a three-dimensional network structure (Table 1, Fig. 2).

Experimental

To an aqueous solution of *p*-amino benzoic acid (1.0 g, 7.3 mmol), sodium carbonate (1 N) was added to adjust the pH 8. Then added 4-(acetylamino) benzenesulfonyl chloride (2.21 g, 9.48 mmol) and the mixture was stirred at room temperature keeping the pH of the mixture up to 8.0 with occasional addition of sodium carbonate solution. Progress and completion of the reaction was confirmed by TLC and conversion of suspension into clear solution. After 2 h, whole mixture was poured into a beaker and the pH was adjusted to 2.0 by 1 N HCl. Precipitates were produced which were filtered and washed with distilled water. The prepared sulfonamide (2-((4-(acetylamino)phenyl)sulfonyl)amino) benzoic acid (1.0 g, 3 mmol), DMF (10 ml) and n-hexane washed sodium hydride (0.22 g, 9.0 mmol) were stirred at room temperature for 40 min followed by the addition of ethyl iodide (0.61 g, 3.9 mmol). The whole reaction mixture was stirred till the completion of the reaction and poured into crushed ice in a beaker. The pH of the mixture was adjusted to 4.0 with 1 N HCl. Precipitates were produced, filtered and washed twice with distilled water and crystallized from chloroform solution as brown blocks.

Refinement

The NH H atom was located in a difference map and refined with the distance restraint N—H = 0.86 (2) Å freely. The remaining H atoms were placed in calculated positions [O—H = 0.82 Å, C—H = 0.93, 0.96 and 0.97 Å], with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}, \text{O})$ for the methyl and hydroxyl H atoms and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for the other H atoms. Four reflections giving bad agreements with Fc, *viz.* (200), (110), (011) and (122), were omitted during the final cycles of refinement.

Computing details

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used

to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).




Figure 1

The molecular structure of the title compound, showing displacement ellipsoids drawn at the 30% probability level.




Figure 2

View of the packing diagram and the hydrogen bonding of (I) along the c axis. Hydrogen atoms that not involved in the hydrogen-bonding (dashed lines) have been omitted for clarity.

2-{4-[Acetyl(ethyl)amino]benzenesulfonamido}benzoic acid*Crystal data*

$C_{17}H_{18}N_2O_5S$
 $M_r = 362.40$
Orthorhombic, $Pna2_1$
Hall symbol: P 2c -2n
 $a = 18.0371 (7)$ Å
 $b = 12.0249 (4)$ Å
 $c = 7.8430 (2)$ Å
 $V = 1701.10 (10)$ Å³
 $Z = 4$

$F(000) = 760$
 $D_x = 1.415 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 4557 reflections
 $\theta = 2.3\text{--}28.2^\circ$
 $\mu = 0.22 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
Block, brown
 $0.35 \times 0.29 \times 0.27$ mm

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: sealed tube
Graphite monochromator
 φ and ω scans
9278 measured reflections
3880 independent reflections

3327 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$
 $\theta_{\text{max}} = 28.3^\circ, \theta_{\text{min}} = 2.8^\circ$
 $h = -24 \rightarrow 19$
 $k = -13 \rightarrow 16$
 $l = -10 \rightarrow 10$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.093$
 $S = 0.99$
3880 reflections
233 parameters
2 restraints
Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0604P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.35 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.18 \text{ e } \text{\AA}^{-3}$
Absolute structure: Flack (1983), 1612 Friedel pairs
Flack parameter: 0.05 (6)

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating - R -factor-obs etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.09768 (2)	1.03872 (4)	1.00764 (6)	0.0403 (1)
O1	0.06408 (9)	0.65010 (13)	1.42241 (19)	0.0555 (5)
O2	0.12860 (8)	0.80244 (12)	1.3690 (2)	0.0570 (5)
O3	0.13309 (9)	1.10513 (13)	1.13429 (18)	0.0551 (5)

O4	0.02644 (8)	1.06853 (14)	0.9452 (2)	0.0568 (5)
O5	0.36221 (10)	1.08713 (16)	0.1903 (2)	0.0660 (6)
N1	0.09488 (10)	0.91519 (15)	1.0928 (2)	0.0473 (6)
N2	0.30964 (11)	1.02317 (15)	0.4272 (3)	0.0544 (6)
C1	0.08374 (10)	0.73379 (15)	1.3240 (2)	0.0402 (5)
C2	0.04690 (9)	0.73281 (15)	1.1545 (2)	0.0376 (5)
C3	0.05278 (10)	0.82314 (14)	1.0418 (2)	0.0371 (5)
C4	0.01818 (12)	0.81716 (18)	0.8837 (3)	0.0494 (6)
C5	-0.02159 (13)	0.72412 (18)	0.8387 (3)	0.0569 (8)
C6	-0.02717 (13)	0.63543 (19)	0.9473 (3)	0.0584 (7)
C7	0.00674 (11)	0.63937 (17)	1.1041 (3)	0.0486 (6)
C8	0.15845 (10)	1.03090 (13)	0.8330 (2)	0.0347 (5)
C9	0.23390 (11)	1.02380 (17)	0.8644 (3)	0.0461 (6)
C10	0.28288 (12)	1.02284 (18)	0.7301 (3)	0.0497 (7)
C11	0.25724 (11)	1.02912 (16)	0.5651 (2)	0.0428 (6)
C12	0.18157 (11)	1.03649 (16)	0.5338 (2)	0.0456 (6)
C13	0.13254 (11)	1.03771 (17)	0.6682 (2)	0.0428 (6)
C14	0.32075 (12)	1.10446 (17)	0.3124 (3)	0.0479 (6)
C15	0.28352 (13)	1.21419 (17)	0.3350 (4)	0.0609 (8)
C16	0.35281 (15)	0.9186 (2)	0.4124 (4)	0.0765 (10)
C17	0.3174 (2)	0.8394 (3)	0.2929 (7)	0.1147 (17)
H1	0.09620	0.63950	1.49470	0.0830*
H1N	0.1178 (11)	0.9097 (19)	1.187 (2)	0.047 (6)*
H4	0.02200	0.87640	0.80800	0.0590*
H5	-0.04500	0.72140	0.73310	0.0680*
H6	-0.05380	0.57270	0.91500	0.0700*
H7	0.00280	0.57890	1.17760	0.0580*
H9	0.25130	1.01970	0.97580	0.0550*
H10	0.33350	1.01790	0.75080	0.0600*
H12	0.16410	1.04060	0.42240	0.0550*
H13	0.08190	1.04310	0.64770	0.0510*
H15A	0.31060	1.27030	0.27400	0.0910*
H15B	0.28220	1.23280	0.45390	0.0910*
H15C	0.23380	1.21010	0.29150	0.0910*
H16A	0.40250	0.93560	0.37260	0.0920*
H16B	0.35700	0.88430	0.52390	0.0920*
H17A	0.27050	0.81600	0.33830	0.1720*
H17B	0.34890	0.77570	0.27820	0.1720*
H17C	0.30990	0.87500	0.18470	0.1720*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0487 (2)	0.0385 (2)	0.0338 (2)	-0.0018 (2)	0.0058 (2)	0.0024 (2)
O1	0.0657 (9)	0.0549 (8)	0.0460 (8)	-0.0130 (7)	-0.0115 (7)	0.0178 (7)
O2	0.0614 (9)	0.0576 (8)	0.0520 (8)	-0.0156 (7)	-0.0192 (7)	0.0128 (7)
O3	0.0783 (10)	0.0481 (8)	0.0388 (7)	-0.0115 (7)	0.0075 (7)	-0.0068 (6)
O4	0.0479 (8)	0.0636 (9)	0.0590 (9)	0.0107 (7)	0.0093 (7)	0.0114 (8)
O5	0.0780 (12)	0.0699 (11)	0.0501 (9)	-0.0151 (8)	0.0208 (8)	0.0087 (8)
N1	0.0584 (11)	0.0472 (9)	0.0363 (9)	-0.0136 (8)	-0.0088 (8)	0.0139 (8)

N2	0.0615 (11)	0.0523 (10)	0.0495 (10)	0.0049 (8)	0.0198 (9)	0.0117 (9)
C1	0.0401 (9)	0.0410 (9)	0.0394 (9)	0.0035 (7)	-0.0001 (7)	0.0065 (8)
C2	0.0317 (8)	0.0421 (9)	0.0390 (9)	0.0000 (7)	0.0032 (7)	0.0026 (8)
C3	0.0350 (9)	0.0390 (8)	0.0374 (10)	-0.0010 (7)	0.0016 (7)	0.0016 (7)
C4	0.0558 (12)	0.0521 (11)	0.0402 (10)	-0.0040 (9)	-0.0063 (9)	0.0083 (9)
C5	0.0664 (14)	0.0582 (13)	0.0462 (12)	-0.0054 (10)	-0.0154 (10)	-0.0039 (10)
C6	0.0641 (13)	0.0492 (12)	0.0618 (13)	-0.0110 (10)	-0.0135 (11)	-0.0042 (10)
C7	0.0503 (11)	0.0422 (10)	0.0534 (12)	-0.0061 (8)	-0.0039 (9)	0.0050 (9)
C8	0.0404 (10)	0.0322 (8)	0.0315 (8)	-0.0021 (6)	0.0018 (7)	0.0012 (7)
C9	0.0462 (11)	0.0583 (12)	0.0339 (9)	-0.0017 (9)	-0.0074 (8)	0.0039 (9)
C10	0.0383 (11)	0.0615 (12)	0.0494 (12)	-0.0003 (9)	-0.0015 (9)	0.0063 (10)
C11	0.0455 (11)	0.0425 (10)	0.0405 (10)	-0.0007 (7)	0.0081 (8)	0.0031 (8)
C12	0.0526 (11)	0.0530 (11)	0.0312 (10)	-0.0037 (9)	-0.0065 (8)	0.0027 (8)
C13	0.0378 (10)	0.0531 (11)	0.0376 (10)	-0.0022 (8)	-0.0065 (8)	0.0040 (8)
C14	0.0513 (11)	0.0468 (10)	0.0457 (11)	-0.0119 (8)	0.0002 (9)	0.0054 (9)
C15	0.0686 (15)	0.0468 (11)	0.0673 (15)	-0.0098 (10)	-0.0047 (12)	0.0065 (11)
C16	0.0816 (19)	0.0763 (17)	0.0716 (17)	0.0207 (14)	0.0323 (14)	0.0162 (15)
C17	0.117 (3)	0.0541 (16)	0.173 (4)	-0.0027 (17)	0.033 (3)	-0.008 (2)

Geometric parameters (\AA , $^\circ$)

S1—O3	1.4256 (16)	C9—C10	1.375 (3)
S1—O4	1.4211 (15)	C10—C11	1.376 (3)
S1—N1	1.6295 (18)	C11—C12	1.390 (3)
S1—C8	1.7568 (17)	C12—C13	1.376 (2)
O1—C1	1.317 (2)	C14—C15	1.491 (3)
O2—C1	1.209 (2)	C16—C17	1.481 (5)
O5—C14	1.233 (3)	C4—H4	0.9300
O1—H1	0.8200	C5—H5	0.9300
N1—C3	1.401 (2)	C6—H6	0.9300
N2—C14	1.344 (3)	C7—H7	0.9300
N2—C16	1.484 (3)	C9—H9	0.9300
N2—C11	1.438 (3)	C10—H10	0.9300
N1—H1N	0.849 (17)	C12—H12	0.9300
C1—C2	1.486 (2)	C13—H13	0.9300
C2—C3	1.404 (2)	C15—H15A	0.9600
C2—C7	1.394 (3)	C15—H15B	0.9600
C3—C4	1.390 (3)	C15—H15C	0.9600
C4—C5	1.375 (3)	C16—H16A	0.9700
C5—C6	1.369 (3)	C16—H16B	0.9700
C6—C7	1.374 (3)	C17—H17A	0.9600
C8—C9	1.386 (3)	C17—H17B	0.9600
C8—C13	1.377 (2)	C17—H17C	0.9600
O3—S1—O4	120.26 (10)	O5—C14—C15	121.0 (2)
O3—S1—N1	103.82 (9)	N2—C14—C15	119.8 (2)
O3—S1—C8	107.08 (9)	N2—C16—C17	111.6 (2)
O4—S1—N1	110.07 (9)	C3—C4—H4	120.00
O4—S1—C8	108.00 (9)	C5—C4—H4	120.00
N1—S1—C8	106.87 (8)	C4—C5—H5	120.00

C1—O1—H1	109.00	C6—C5—H5	120.00
S1—N1—C3	128.33 (13)	C5—C6—H6	120.00
C11—N2—C16	116.48 (19)	C7—C6—H6	120.00
C14—N2—C16	119.1 (2)	C2—C7—H7	120.00
C11—N2—C14	124.45 (18)	C6—C7—H7	120.00
C3—N1—H1N	116.8 (15)	C8—C9—H9	120.00
S1—N1—H1N	114.4 (15)	C10—C9—H9	120.00
O1—C1—C2	113.44 (16)	C9—C10—H10	120.00
O1—C1—O2	122.08 (16)	C11—C10—H10	120.00
O2—C1—C2	124.47 (16)	C11—C12—H12	120.00
C1—C2—C3	121.54 (15)	C13—C12—H12	120.00
C1—C2—C7	119.49 (17)	C8—C13—H13	120.00
C3—C2—C7	118.96 (16)	C12—C13—H13	120.00
C2—C3—C4	119.18 (16)	C14—C15—H15A	109.00
N1—C3—C2	118.19 (15)	C14—C15—H15B	109.00
N1—C3—C4	122.62 (16)	C14—C15—H15C	109.00
C3—C4—C5	120.4 (2)	H15A—C15—H15B	109.00
C4—C5—C6	120.8 (2)	H15A—C15—H15C	109.00
C5—C6—C7	119.8 (2)	H15B—C15—H15C	110.00
C2—C7—C6	120.85 (19)	N2—C16—H16A	109.00
C9—C8—C13	120.26 (17)	N2—C16—H16B	109.00
S1—C8—C13	121.13 (14)	C17—C16—H16A	109.00
S1—C8—C9	118.52 (14)	C17—C16—H16B	109.00
C8—C9—C10	119.7 (2)	H16A—C16—H16B	108.00
C9—C10—C11	120.3 (2)	C16—C17—H17A	109.00
N2—C11—C10	118.92 (19)	C16—C17—H17B	109.00
N2—C11—C12	121.05 (16)	C16—C17—H17C	109.00
C10—C11—C12	119.98 (17)	H17A—C17—H17B	109.00
C11—C12—C13	119.78 (15)	H17A—C17—H17C	110.00
C8—C13—C12	120.02 (18)	H17B—C17—H17C	109.00
O5—C14—N2	119.2 (2)		
O3—S1—N1—C3	165.17 (17)	O2—C1—C2—C3	-10.4 (3)
O4—S1—N1—C3	35.20 (19)	O1—C1—C2—C7	-10.4 (2)
C8—S1—N1—C3	-81.84 (18)	C1—C2—C7—C6	-179.19 (19)
N1—S1—C8—C13	109.88 (16)	C7—C2—C3—C4	0.3 (3)
N1—S1—C8—C9	-73.62 (16)	C7—C2—C3—N1	-178.43 (17)
O3—S1—C8—C9	37.13 (17)	C1—C2—C3—C4	178.99 (17)
O4—S1—C8—C9	167.99 (15)	C3—C2—C7—C6	-0.5 (3)
O3—S1—C8—C13	-139.37 (15)	C1—C2—C3—N1	0.3 (2)
O4—S1—C8—C13	-8.52 (18)	C2—C3—C4—C5	0.3 (3)
S1—N1—C3—C2	-166.53 (14)	N1—C3—C4—C5	178.98 (19)
S1—N1—C3—C4	14.8 (3)	C3—C4—C5—C6	-0.8 (3)
C14—N2—C11—C12	64.0 (3)	C4—C5—C6—C7	0.6 (4)
C16—N2—C14—C15	-174.0 (2)	C5—C6—C7—C2	0.0 (3)
C16—N2—C11—C10	62.2 (3)	S1—C8—C9—C10	-176.96 (16)
C11—N2—C14—C15	7.0 (3)	C9—C8—C13—C12	0.6 (3)
C14—N2—C16—C17	-85.0 (3)	C13—C8—C9—C10	-0.4 (3)
C11—N2—C16—C17	94.2 (3)	S1—C8—C13—C12	177.02 (15)

C11—N2—C14—O5	−173.6 (2)	C8—C9—C10—C11	0.1 (3)
C14—N2—C11—C10	−118.7 (2)	C9—C10—C11—C12	0.0 (3)
C16—N2—C14—O5	5.5 (3)	C9—C10—C11—N2	−177.39 (19)
C16—N2—C11—C12	−115.1 (2)	C10—C11—C12—C13	0.1 (3)
O2—C1—C2—C7	168.24 (18)	N2—C11—C12—C13	177.48 (18)
O1—C1—C2—C3	170.93 (16)	C11—C12—C13—C8	−0.4 (3)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···O5 ⁱ	0.82	1.82	2.599 (2)	158
N1—H1N···O2	0.85 (2)	1.93 (2)	2.627 (2)	138 (2)
C12—H12···O3 ⁱⁱ	0.93	2.45	3.356 (2)	164
C15—H15C···O3 ⁱⁱ	0.96	2.53	3.400 (3)	150
C17—H17A···O2 ⁱⁱ	0.96	2.58	3.486 (4)	158

Symmetry codes: (i) $-x+1/2, y-1/2, z+3/2$; (ii) $x, y, z-1$.