

## 2,2'-(Ethane-1,2-diyl)bis(1*H*-benzimidazole)

Xu Chen,<sup>a</sup> Xiao-Yong Wu<sup>b</sup> and Guo-Liang Zhao<sup>a,b\*</sup>

<sup>a</sup>Zhejiang Normal University Xingzhi College, Jinhua, Zhejiang 321004, People's Republic of China, and <sup>b</sup>College of Chemistry and Life Sciences, Zhejiang Normal University, Jinhua 321004, Zhejiang, People's Republic of China  
Correspondence e-mail: sky53@zjnu.cn


Received 16 March 2012; accepted 30 March 2012

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.045;  $wR$  factor = 0.122; data-to-parameter ratio = 16.2.

The complete molecule of the title compound,  $\text{C}_{16}\text{H}_{14}\text{N}_4$ , is generated by crystallographic inversion symmetry. In the crystal, molecules are linked by  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bonds, generating (001) sheets. Weak aromatic  $\pi-\pi$  stacking interactions [centroid–centroid distances = 3.7383 (13) and 3.7935 (14)  $\text{\AA}$ ] are also observed.

### Related literature

For background to metal-organic frameworks, see: van Albada *et al.* (2007). For the synthesis, see: Wang & Joulli (1957).



### Experimental

#### Crystal data

$\text{C}_{16}\text{H}_{14}\text{N}_4$

$M_r = 262.31$

Orthorhombic,  $Pbca$   
 $a = 8.4295 (18)\text{ \AA}$   
 $b = 9.924 (2)\text{ \AA}$   
 $c = 15.351 (4)\text{ \AA}$   
 $V = 1284.2 (5)\text{ \AA}^3$

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.08\text{ mm}^{-1}$   
 $T = 296\text{ K}$   
 $0.32 \times 0.25 \times 0.19\text{ mm}$

#### Data collection

Bruker APEXII CCD  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.975$ ,  $T_{\max} = 0.984$

10702 measured reflections  
1475 independent reflections  
966 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.052$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.122$   
 $S = 1.02$   
1475 reflections

91 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.15\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.21\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2B $\cdots$ N1 <sup>i</sup>	0.86	2.04	2.8568 (18)	159
Symmetry code: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, z$ .				

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6688).

### References

- Albada, G. A. van, Mutikainen, I., Turpeinen, U. & Reedijk, J. (2007). *J. Chem. Cryst.* **37**, 489–496.
- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wang, L. L.-Y. & Joulli, M. M. (1957). *J. Am. Chem. Soc.* **79**, 5706–5708.

# supplementary materials

*Acta Cryst.* (2012). E68, o1321 [doi:10.1107/S1600536812013839]

## 2,2'-(Ethane-1,2-diyl)bis(1*H*-benzimidazole)

**Xu Chen, Xiao-Yong Wu and Guo-Liang Zhao**

### Comment

We present a polyaromatic compound (1) that contains multiple functional groups that can develop a series of metal-organic frameworks with potential applications (e.g. van Albada *et al.* (2007)). The molecular structure of (1), shown in Fig. 1, consists of two symmetrical benzimidazole rings. The two benzimidazolyl rings are nearly parallel, with a dihedral angle of 2.645 (6) $^{\circ}$  between them. Molecules are linked *via* a network of hydrogen bonds (N2—H2B $\cdots$ N1; Table 2).  $\pi$ — $\pi$  stacking interactions are observed between nearly parallel benzimidazolyl benzene rings. The centroid-to-centroid distance between C1—C6 benzene rings is 3.7379 Å, while between C1A—C6A benzene rings it is 3.7944 Å (the symmetry operation:  $-x + 1, -y, -z$ ). The hydrogen bonds and  $\pi$ — $\pi$  weak non-covalent interactions lend stability to the structure. The stacking plot of this compound was shown in Fig. 2.

### Experimental


*o*-Phenylenediamine (1.081 g, 10 mmol) and succinic acid (0.590 g, 5 mmol) were refluxed for 4 h in 30 ml of 10% hydrochloric acid solution. The reaction mixture was cooled to room temperature, the precipitation that formed was filtered and then recrystallized from water. The colourless crystals were obtained from water after a week (Wang *et al.*, 1957).

### Refinement


The H atoms bonded to C and N atoms were positioned geometrically and refined using a riding model [aliphatic C—H = 0.96 Å ( $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ )), aromatic C—H = 0.93 Å ( $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ ) and N—H = 0.86 Å with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ ].

### Computing details

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT* (Bruker, 2006); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).

**Figure 1**

The molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level. Atoms with suffix A are at the symmetry position (1-x, 1-y, -z).

**Figure 2**

The stacking plot of the title compound, showing H-bond interactions (dashed lines) and  $\pi-\pi$  stacking interactions.

### **2,2'-(Ethane-1,2-diyl)bis(1H-benzimidazole)**

#### *Crystal data*

$C_{16}H_{14}N_4$   
 $M_r = 262.31$   
Orthorhombic,  $Pbca$   
Hall symbol: -P 2ac 2ab  
 $a = 8.4295 (18) \text{ \AA}$   
 $b = 9.924 (2) \text{ \AA}$   
 $c = 15.351 (4) \text{ \AA}$   
 $V = 1284.2 (5) \text{ \AA}^3$   
 $Z = 4$

$F(000) = 552$   
 $D_x = 1.357 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
 $\theta = 2.7-27.6^\circ$   
 $\mu = 0.08 \text{ mm}^{-1}$   
 $T = 296 \text{ K}$   
Block, colourless  
 $0.32 \times 0.25 \times 0.19 \text{ mm}$

*Data collection*

Bruker APEXII CCD	10702 measured reflections
diffractometer	1475 independent reflections
Radiation source: fine-focus sealed tube	966 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.052$
phi and $\omega$ scans	$\theta_{\text{max}} = 27.6^\circ, \theta_{\text{min}} = 2.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.975, T_{\text{max}} = 0.984$	$k = -12 \rightarrow 12$
	$l = -19 \rightarrow 16$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.045$	H-atom parameters constrained
$wR(F^2) = 0.122$	$w = 1/[\sigma^2(F_o^2) + (0.0554P)^2 + 0.2274P]$
$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
1475 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
91 parameters	$\Delta\rho_{\text{max}} = 0.15 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
N2	0.18376 (15)	0.39276 (12)	0.06131 (8)	0.0378 (4)
H2B	0.1935	0.3084	0.0498	0.045*
N1	0.23667 (16)	0.61293 (12)	0.06218 (9)	0.0402 (4)
C5	0.09915 (18)	0.58969 (15)	0.11024 (10)	0.0357 (4)
C8	0.28216 (19)	0.49246 (14)	0.03497 (11)	0.0375 (4)
C4	0.06495 (18)	0.45137 (15)	0.11012 (10)	0.0354 (4)
C7	0.42738 (19)	0.46475 (17)	-0.01726 (12)	0.0452 (4)
H7A	0.4092	0.4932	-0.0769	0.054*
H7B	0.4467	0.3684	-0.0178	0.054*
C3	-0.0631 (2)	0.39827 (18)	0.15503 (11)	0.0475 (5)
H3A	-0.0860	0.3066	0.1537	0.057*
C6	0.0040 (2)	0.67805 (16)	0.15747 (11)	0.0461 (5)
H6A	0.0247	0.7701	0.1581	0.055*
C2	-0.1552 (2)	0.48740 (18)	0.20187 (12)	0.0524 (5)
H2A	-0.2416	0.4551	0.2333	0.063*
C1	-0.1215 (2)	0.62512 (18)	0.20317 (12)	0.0520 (5)
H1A	-0.1856	0.6824	0.2357	0.062*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N2	0.0346 (8)	0.0281 (6)	0.0506 (8)	0.0003 (5)	0.0010 (6)	-0.0046 (6)
N1	0.0373 (8)	0.0317 (7)	0.0515 (8)	-0.0013 (5)	0.0053 (6)	-0.0039 (6)
C5	0.0323 (9)	0.0347 (8)	0.0402 (9)	0.0015 (7)	-0.0014 (7)	0.0005 (6)
C8	0.0328 (9)	0.0329 (8)	0.0469 (9)	-0.0014 (6)	-0.0024 (7)	-0.0044 (7)
C4	0.0326 (9)	0.0330 (8)	0.0407 (9)	0.0013 (6)	-0.0027 (7)	-0.0005 (6)
C7	0.0343 (9)	0.0450 (9)	0.0563 (11)	-0.0019 (7)	0.0050 (8)	-0.0102 (8)
C3	0.0458 (11)	0.0432 (9)	0.0537 (10)	-0.0071 (8)	0.0034 (9)	0.0023 (8)
C6	0.0473 (11)	0.0371 (9)	0.0541 (11)	0.0069 (8)	0.0020 (9)	-0.0036 (7)
C2	0.0435 (11)	0.0626 (12)	0.0512 (11)	-0.0013 (9)	0.0105 (9)	0.0027 (9)
C1	0.0458 (11)	0.0571 (11)	0.0529 (11)	0.0130 (9)	0.0083 (9)	-0.0057 (9)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

N2—C8	1.3530 (19)	C7—H7A	0.9700
N2—C4	1.3795 (19)	C7—H7B	0.9700
N2—H2B	0.8600	C3—C2	1.379 (2)
N1—C8	1.3233 (18)	C3—H3A	0.9300
N1—C5	1.393 (2)	C6—C1	1.374 (2)
C5—C6	1.392 (2)	C6—H6A	0.9300
C5—C4	1.403 (2)	C2—C1	1.396 (2)
C8—C7	1.489 (2)	C2—H2A	0.9300
C4—C3	1.385 (2)	C1—H1A	0.9300
C7—C7 <sup>i</sup>	1.506 (3)		
C8—N2—C4	107.40 (12)	C8—C7—H7B	109.0
C8—N2—H2B	126.3	C7 <sup>i</sup> —C7—H7B	109.0
C4—N2—H2B	126.3	H7A—C7—H7B	107.8
C8—N1—C5	104.99 (12)	C2—C3—C4	117.01 (16)
C6—C5—N1	130.61 (14)	C2—C3—H3A	121.5
C6—C5—C4	119.91 (15)	C4—C3—H3A	121.5
N1—C5—C4	109.41 (13)	C1—C6—C5	117.96 (15)
N1—C8—N2	112.88 (14)	C1—C6—H6A	121.0
N1—C8—C7	125.11 (14)	C5—C6—H6A	121.0
N2—C8—C7	122.00 (13)	C3—C2—C1	121.40 (17)
N2—C4—C3	132.51 (15)	C3—C2—H2A	119.3
N2—C4—C5	105.32 (13)	C1—C2—H2A	119.3
C3—C4—C5	122.14 (15)	C6—C1—C2	121.57 (16)
C8—C7—C7 <sup>i</sup>	113.14 (16)	C6—C1—H1A	119.2
C8—C7—H7A	109.0	C2—C1—H1A	119.2
C7 <sup>i</sup> —C7—H7A	109.0		
C8—N1—C5—C6	176.72 (17)	N1—C5—C4—C3	178.13 (14)
C8—N1—C5—C4	-0.20 (18)	N1—C8—C7—C7 <sup>i</sup>	46.1 (3)
C5—N1—C8—N2	0.41 (18)	N2—C8—C7—C7 <sup>i</sup>	-132.3 (2)
C5—N1—C8—C7	-178.13 (15)	N2—C4—C3—C2	176.40 (16)
C4—N2—C8—N1	-0.47 (19)	C5—C4—C3—C2	-1.2 (2)
C4—N2—C8—C7	178.12 (14)	N1—C5—C6—C1	-176.40 (16)

C8—N2—C4—C3	−177.62 (17)	C4—C5—C6—C1	0.2 (2)
C8—N2—C4—C5	0.31 (17)	C4—C3—C2—C1	0.6 (3)
C6—C5—C4—N2	−177.37 (14)	C5—C6—C1—C2	−0.9 (3)
N1—C5—C4—N2	−0.07 (17)	C3—C2—C1—C6	0.4 (3)
C6—C5—C4—C3	0.8 (2)		

Symmetry code: (i)  $-x+1, -y+1, -z$ .

*Hydrogen-bond geometry ( $\text{\AA}$ , °)*

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N2—H2B $\cdots$ N1 <sup>ii</sup>	0.86	2.04	2.8568 (18)	159

Symmetry code: (ii)  $-x+1/2, y-1/2, z$ .