

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-[1,1-Dioxo-2-(2,4,5-trifluorobenzyl)-2H-1,2-benzothiazin-4-yl]acetic acid

Yanchun Yang,^a Youzhu Yu^b and Changjin Zhu^{a*}

^aDepartment of Applied Chemistry, Beijing Institute of Technology, Zhongguancun South Street, 100081 Beijing, People's Republic of China, and ^bDepartment of Chemistry and Environmental Engineering, Anyang Institute of Technology, Henan 455000, People's Republic of China
Correspondence e-mail: zcj@bit.edu.cn


Received 13 March 2012; accepted 3 April 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.039; wR factor = 0.108; data-to-parameter ratio = 17.4.

In the title compound, $\text{C}_{17}\text{H}_{12}\text{F}_3\text{NO}_4\text{S}$, the heterocyclic thiazine ring adopts a half-chair conformation with the S and the N atoms displaced by -0.608 (3) and 0.105 (3) Å, respectively, from the mean plane formed by the remaining ring atoms. The dihedral angle between the two benzene rings is 36.63 (8)° and the acetic acid group is inclined at right angles [89.78 (8)°] to the mean plane formed by the C atoms of the thiazine ring. The crystal structure features $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For pharmaceutical properties of benzothiazines, see: Zia-ur-Rehman *et al.* (2006). For synthetic details of the title compound, see: Chen *et al.* (2011). For related structures, see: Ahmad *et al.* (2008); Zia-ur-Rehman *et al.* (2008).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{12}\text{F}_3\text{NO}_4\text{S}$
 $M_r = 383.34$

Monoclinic, $P2_1/n$
 $a = 8.3290$ (17) Å

$b = 23.141$ (5) Å
 $c = 8.6692$ (17) Å
 $\beta = 90.93$ (3)°
 $V = 1670.7$ (6) Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.25$ mm⁻¹
 $T = 293$ K
 $0.20 \times 0.20 \times 0.20$ mm

Data collection

Bruker APEXII CCD
diffractometer
22476 measured reflections

4158 independent reflections
3646 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.108$
 $S = 1.03$
4158 reflections
239 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.32$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.40$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{O2}^{\text{i}}$	0.84 (3)	1.83 (3)	2.675 (2)	179 (3)
$\text{C8}-\text{H8}\cdots\text{O3}^{\text{ii}}$	0.93	2.55	3.211 (2)	129
$\text{C9}-\text{H9A}\cdots\text{O3}^{\text{ii}}$	0.97	2.30	3.207 (2)	155
$\text{C11}-\text{H11B}\cdots\text{O4}^{\text{iii}}$	0.97	2.40	3.162 (2)	135

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$; (iii) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

This work was supported by the Beijing Natural Science Foundation (No. 7102091) and the Research Fund for the Doctoral Program of Higher Education of China (No. 20111101110042).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2523).

References

- Ahmad, M., Latif Siddiqui, H., Zia-ur-Rehman, M., Tizzard, G. J. & Ahmad, S. (2008). *Acta Cryst.* **E64**, o1392.
Bruker (2001). *SAINTE-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2005). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
Chen, X., Zhang, S., Yang, Y., Hussain, S., He, M., Gui, D., Ma, B., Jing, C., Qiao, Z., Zhu, C. & Yu, Q. (2011). *Bioorg. Med. Chem.* **19**, 7262–7269.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Zia-ur-Rehman, M., Choudary, J. A., Ahmad, S. & Siddiqui, H. L. (2006). *Chem. Pharm. Bull.* **54**, 1175–1178.
Zia-ur-Rehman, M., Choudary, J. A., Elsegood, M. R. J., Akbar, N. & Latif Siddiqui, H. (2008). *Acta Cryst.* **E64**, o1508.

supplementary materials

Acta Cryst. (2012). E68, o1364 [doi:10.1107/S1600536812015942]

2-[1,1-Dioxo-2-(2,4,5-trifluorobenzyl)-2H-1,2-benzothiazin-4-yl]acetic acid**Yanchun Yang, Youzhu Yu and Changjin Zhu****Comment**

Benzothiazine moiety is present as the skeletal structure in several pharmaceuticals such as antibacterial, diuretic, hypoglycemic, antithyroid, and antitumor drugs (Zia-ur-Rehman *et al.*, 2006). In this article, we report the crystal structure of the title compound which has been used as aldose reductase inhibitor (Chen *et al.*, 2011).

In the title compound (Fig. 1), the heterocyclic thiazine ring adopts a half chair conformation with the S1 and the N1 atoms displaced by -0.608 (3) and 0.105 (3) Å, respectively, from the mean plane formed by the remaining ring atoms (C1/C6/C7/C8). The dihedral angle between the two benze rings (C1–C6) and (C12–C17) is 36.63 (8)° and the acetate group (O1/O2/C9/C10) is inclined at right angles (89.78 (8) °) to the mean plane formed by the C-atoms of the thiazine ring. The crystal structure is stabilized by intermolecular O—H···O and C—H···O hydrogen bonds forming a three-dimensional network (Fig. 2).

Experimental

A mixture of methyl 2-(1,1-dioxido-2-(2,4,5-trifluorobenzyl)-2H-benzo[e][1,2] thiazin-4-yl)acetate (0.5 mmol), 1,4-dioxane (5 ml) and saturated aqueous sodium hydroxide (8 ml) was stirred at room temperature for 12 h. The alkaline suspension was adjusted to be acidic with 0.1 M HCl and extracted with ethyl acetate (3 x 30 ml). The combined organic layers were dried over MgSO₄ and filtered. Crystals suitable for X-ray diffraction were obtained by slow evaporation of a solution of the title compound in methanol (yield = 69%).

Refinement

H atom bonded to O1 was located from a different Fourier map and refined freely. The remaining H atoms were positioned geometrically, with C—H = 0.93 and 0.97 Å for aromatic and methylene H, respectively, and constrained to ride on their parent atoms with $U_{iso}(H) = 1.2U_{eq}(C)$.

Computing details

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINTE-Plus* (Bruker, 2001); data reduction: *SAINTE-Plus* (Bruker, 2001); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).




Figure 1

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as small spheres of arbitrary radius.




Figure 2

A view of the O—H...O and C—H...O hydrogen bonds (dotted lines) in the crystal structure of the title compound.

2-[1,1-Dioxo-2-(2,4,5-trifluorobenzyl)-2H-1,2-benzothiazin-4-yl]acetic acid

Crystal data

$C_{17}H_{12}F_3NO_4S$	$F(000) = 784$
$M_r = 383.34$	$D_x = 1.524 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-P 2_1n$	Cell parameters from 9975 reflections
$a = 8.3290 (17) \text{ \AA}$	$\theta = 2.5\text{--}28.4^\circ$
$b = 23.141 (5) \text{ \AA}$	$\mu = 0.25 \text{ mm}^{-1}$
$c = 8.6692 (17) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 90.93 (3)^\circ$	Block, colorless
$V = 1670.7 (6) \text{ \AA}^3$	$0.20 \times 0.20 \times 0.20 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD diffractometer	3646 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.023$
Graphite monochromator	$\theta_{\text{max}} = 28.4^\circ$, $\theta_{\text{min}} = 2.5^\circ$
φ and ω scans	$h = -11 \rightarrow 11$
22476 measured reflections	$k = -27 \rightarrow 30$
4158 independent reflections	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.039$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.108$	$w = 1/[\sigma^2(F_o^2) + (0.0536P)^2 + 0.5773P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
4158 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
239 parameters	$\Delta\rho_{\text{max}} = 0.32 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.40 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.11950 (4)	0.232698 (14)	0.97865 (4)	0.03556 (11)
F1	0.69758 (14)	0.21289 (5)	0.89004 (16)	0.0722 (3)
F2	0.6328 (2)	0.02123 (6)	0.73067 (19)	0.0989 (5)
F3	0.3601 (2)	0.01403 (5)	0.8862 (2)	0.0997 (5)
O1	0.4665 (2)	0.48147 (7)	0.79699 (18)	0.0807 (5)

H1	0.504 (4)	0.5074 (14)	0.855 (3)	0.107 (10)*
O2	0.41807 (19)	0.43626 (6)	1.01497 (15)	0.0686 (4)
O3	0.08188 (15)	0.19719 (5)	1.10801 (14)	0.0526 (3)
O4	0.08941 (14)	0.20919 (5)	0.82864 (12)	0.0463 (3)
N1	0.30918 (15)	0.25127 (5)	0.99625 (14)	0.0389 (3)
C1	0.02950 (17)	0.30051 (6)	0.99039 (15)	0.0348 (3)
C2	-0.11807 (18)	0.30537 (7)	1.06150 (18)	0.0419 (3)
H2	-0.1616	0.2741	1.1136	0.050*
C3	-0.19919 (19)	0.35725 (8)	1.0537 (2)	0.0504 (4)
H3	-0.2977	0.3614	1.1014	0.060*
C4	-0.1331 (2)	0.40320 (7)	0.9744 (2)	0.0525 (4)
H4	-0.1901	0.4376	0.9658	0.063*
C5	0.0162 (2)	0.39876 (7)	0.90794 (19)	0.0453 (3)
H5	0.0589	0.4304	0.8568	0.054*
C6	0.10414 (17)	0.34714 (6)	0.91652 (15)	0.0354 (3)
C7	0.26885 (18)	0.34231 (6)	0.86404 (16)	0.0379 (3)
C8	0.36224 (18)	0.29773 (6)	0.90869 (17)	0.0387 (3)
H8	0.4691	0.2980	0.8792	0.046*
C9	0.3439 (2)	0.39103 (6)	0.77500 (19)	0.0448 (3)
H9A	0.4296	0.3755	0.7128	0.054*
H9B	0.2636	0.4074	0.7054	0.054*
C10	0.41079 (18)	0.43834 (6)	0.87590 (19)	0.0433 (3)
C11	0.4227 (2)	0.21175 (7)	1.07534 (18)	0.0448 (3)
H11A	0.3728	0.1972	1.1679	0.054*
H11B	0.5170	0.2336	1.1071	0.054*
C12	0.47589 (19)	0.16095 (7)	0.97972 (17)	0.0419 (3)
C13	0.6136 (2)	0.16311 (7)	0.8936 (2)	0.0480 (4)
C14	0.6717 (2)	0.11724 (9)	0.8094 (2)	0.0594 (4)
H14	0.7667	0.1200	0.7549	0.071*
C15	0.5833 (3)	0.06758 (8)	0.8099 (2)	0.0636 (5)
C16	0.4443 (3)	0.06386 (8)	0.8907 (3)	0.0635 (5)
C17	0.3901 (2)	0.10940 (7)	0.9768 (2)	0.0532 (4)
H17	0.2964	0.1058	1.0329	0.064*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.04325 (19)	0.02765 (17)	0.03590 (18)	-0.00299 (12)	0.00411 (13)	0.00177 (12)
F1	0.0606 (7)	0.0529 (6)	0.1035 (9)	-0.0118 (5)	0.0109 (6)	-0.0096 (6)
F2	0.1302 (13)	0.0558 (8)	0.1110 (11)	0.0240 (8)	0.0069 (9)	-0.0322 (7)
F3	0.1114 (11)	0.0384 (7)	0.1489 (14)	-0.0155 (7)	-0.0099 (10)	-0.0094 (7)
O1	0.1217 (14)	0.0547 (9)	0.0655 (9)	-0.0464 (9)	-0.0027 (9)	0.0112 (7)
O2	0.0991 (11)	0.0490 (8)	0.0580 (8)	-0.0326 (7)	0.0064 (7)	-0.0001 (6)
O3	0.0639 (7)	0.0421 (6)	0.0523 (6)	-0.0011 (5)	0.0144 (5)	0.0156 (5)
O4	0.0566 (6)	0.0365 (6)	0.0457 (6)	-0.0028 (5)	-0.0032 (5)	-0.0089 (4)
N1	0.0412 (6)	0.0332 (6)	0.0424 (6)	-0.0001 (5)	-0.0004 (5)	0.0018 (5)
C1	0.0411 (7)	0.0302 (6)	0.0331 (6)	-0.0014 (5)	0.0012 (5)	-0.0017 (5)
C2	0.0420 (7)	0.0416 (8)	0.0423 (7)	-0.0052 (6)	0.0048 (6)	0.0002 (6)
C3	0.0416 (8)	0.0525 (9)	0.0573 (9)	0.0042 (7)	0.0078 (7)	-0.0013 (7)
C4	0.0495 (9)	0.0419 (9)	0.0662 (10)	0.0100 (7)	0.0044 (7)	0.0025 (7)

C5	0.0517 (8)	0.0329 (7)	0.0514 (8)	0.0015 (6)	0.0044 (7)	0.0037 (6)
C6	0.0423 (7)	0.0295 (7)	0.0344 (6)	-0.0023 (5)	0.0020 (5)	-0.0015 (5)
C7	0.0459 (7)	0.0279 (6)	0.0402 (7)	-0.0049 (5)	0.0085 (6)	-0.0032 (5)
C8	0.0403 (7)	0.0327 (7)	0.0434 (7)	-0.0047 (5)	0.0063 (6)	-0.0041 (5)
C9	0.0538 (8)	0.0320 (7)	0.0491 (8)	-0.0048 (6)	0.0167 (7)	-0.0006 (6)
C10	0.0424 (7)	0.0308 (7)	0.0572 (9)	-0.0036 (6)	0.0103 (6)	0.0024 (6)
C11	0.0513 (8)	0.0433 (8)	0.0394 (7)	0.0062 (7)	-0.0103 (6)	-0.0030 (6)
C12	0.0484 (8)	0.0356 (7)	0.0413 (7)	0.0047 (6)	-0.0097 (6)	0.0019 (6)
C13	0.0512 (8)	0.0382 (8)	0.0543 (9)	0.0021 (6)	-0.0063 (7)	-0.0013 (7)
C14	0.0631 (11)	0.0552 (11)	0.0601 (11)	0.0133 (9)	0.0033 (8)	-0.0029 (8)
C15	0.0838 (13)	0.0409 (9)	0.0659 (11)	0.0163 (9)	-0.0075 (10)	-0.0110 (8)
C16	0.0782 (13)	0.0320 (8)	0.0796 (13)	-0.0022 (8)	-0.0153 (10)	-0.0013 (8)
C17	0.0579 (10)	0.0395 (8)	0.0621 (10)	0.0000 (7)	-0.0032 (8)	0.0049 (7)

Geometric parameters (Å, °)

S1—O4	1.4281 (11)	C5—C6	1.403 (2)
S1—O3	1.4292 (11)	C5—H5	0.9300
S1—N1	1.6420 (13)	C6—C7	1.457 (2)
S1—C1	1.7427 (14)	C7—C8	1.345 (2)
F1—C13	1.348 (2)	C7—C9	1.5076 (19)
F2—C15	1.342 (2)	C8—H8	0.9300
F3—C16	1.350 (2)	C9—C10	1.503 (2)
O1—C10	1.2998 (19)	C9—H9A	0.9700
O1—H1	0.84 (3)	C9—H9B	0.9700
O2—C10	1.207 (2)	C11—C12	1.509 (2)
N1—C8	1.3923 (18)	C11—H11A	0.9700
N1—C11	1.4765 (19)	C11—H11B	0.9700
C1—C2	1.388 (2)	C12—C13	1.379 (2)
C1—C6	1.4049 (19)	C12—C17	1.391 (2)
C2—C3	1.379 (2)	C13—C14	1.380 (2)
C2—H2	0.9300	C14—C15	1.365 (3)
C3—C4	1.385 (2)	C14—H14	0.9300
C3—H3	0.9300	C15—C16	1.366 (3)
C4—C5	1.383 (2)	C16—C17	1.372 (3)
C4—H4	0.9300	C17—H17	0.9300
O4—S1—O3	117.26 (8)	C10—C9—C7	113.55 (13)
O4—S1—N1	109.80 (7)	C10—C9—H9A	108.9
O3—S1—N1	107.48 (8)	C7—C9—H9A	108.9
O4—S1—C1	109.08 (7)	C10—C9—H9B	108.9
O3—S1—C1	111.83 (7)	C7—C9—H9B	108.9
N1—S1—C1	99.97 (7)	H9A—C9—H9B	107.7
C10—O1—H1	111 (2)	O2—C10—O1	122.93 (16)
C8—N1—C11	121.67 (13)	O2—C10—C9	124.36 (14)
C8—N1—S1	117.70 (10)	O1—C10—C9	112.65 (15)
C11—N1—S1	119.28 (11)	N1—C11—C12	114.76 (12)
C2—C1—C6	122.78 (13)	N1—C11—H11A	108.6
C2—C1—S1	118.92 (11)	C12—C11—H11A	108.6
C6—C1—S1	118.10 (10)	N1—C11—H11B	108.6

C3—C2—C1	119.04 (14)	C12—C11—H11B	108.6
C3—C2—H2	120.5	H11A—C11—H11B	107.6
C1—C2—H2	120.5	C13—C12—C17	116.91 (15)
C2—C3—C4	119.63 (15)	C13—C12—C11	121.54 (14)
C2—C3—H3	120.2	C17—C12—C11	121.54 (15)
C4—C3—H3	120.2	F1—C13—C12	118.65 (15)
C5—C4—C3	121.14 (15)	F1—C13—C14	117.28 (16)
C5—C4—H4	119.4	C12—C13—C14	124.07 (16)
C3—C4—H4	119.4	C15—C14—C13	116.87 (18)
C4—C5—C6	120.89 (14)	C15—C14—H14	121.6
C4—C5—H5	119.6	C13—C14—H14	121.6
C6—C5—H5	119.6	F2—C15—C14	120.1 (2)
C5—C6—C1	116.36 (13)	F2—C15—C16	118.81 (19)
C5—C6—C7	122.84 (13)	C14—C15—C16	121.11 (17)
C1—C6—C7	120.62 (13)	F3—C16—C15	118.92 (19)
C8—C7—C6	120.81 (13)	F3—C16—C17	119.8 (2)
C8—C7—C9	118.58 (13)	C15—C16—C17	121.31 (17)
C6—C7—C9	120.18 (13)	C16—C17—C12	119.69 (18)
C7—C8—N1	124.19 (13)	C16—C17—H17	120.2
C7—C8—H8	117.9	C12—C17—H17	120.2
N1—C8—H8	117.9		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots O2 ⁱ	0.84 (3)	1.83 (3)	2.675 (2)	179 (3)
C8—H8 \cdots O3 ⁱⁱ	0.93	2.55	3.211 (2)	129
C9—H9A \cdots O3 ⁱⁱ	0.97	2.30	3.207 (2)	155
C11—H11A \cdots O3	0.97	2.47	2.877 (2)	105
C11—H11B \cdots F1	0.97	2.47	2.818 (2)	101
C11—H11B \cdots O4 ⁱⁱⁱ	0.97	2.40	3.162 (2)	135

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $x+1/2, -y+1/2, z-1/2$; (iii) $x+1/2, -y+1/2, z+1/2$.