

Dimethyl 3-(cyclopropylcarbonyl)-pyrrolo[2,1-a]isoquinoline-1,2-dicarboxylate

Honglong Xing,^a Fan Tang^a and Wei Wang^{b*}

^aSchool of Chemical Engineering, Anhui University of Science and Technology, Huainan 232001, People's Republic of China, and ^bKey Laboratory for Advanced Technology in Environmental Protection of Jiangsu Province, School of Chemical and Biological Engineering, Yancheng Institute of Technology, Yancheng 224051, People's Republic of China
Correspondence e-mail: wangw@ycit.edu.cn


Received 30 March 2012; accepted 10 April 2012

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.056; wR factor = 0.179; data-to-parameter ratio = 13.1.

In the molecular structure of the title compound, $\text{C}_{20}\text{H}_{17}\text{NO}_5$, two intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond generate six- and seven-membered ring motifs. The dihedral angles between the almost planar 13-atom triple-fused-ring system (r.m.s. deviation = 0.003 \AA) and the planes of the two methoxy-carbonyl substituents are $61.7(2)$ and $33.01(10)^\circ$.

Related literature

For chemical background, see: Michael (2004); Sriram *et al.* (2005); Alonso *et al.* (1985). For the biological activity of indolizine derivatives, see: Shen *et al.* (2010).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{17}\text{NO}_5$
 $M_r = 351.35$

Monoclinic, $P2_1/c$
 $a = 7.5910(15)\text{ \AA}$

$b = 18.436(4)\text{ \AA}$
 $c = 12.162(2)\text{ \AA}$
 $\beta = 94.43(3)^\circ$
 $V = 1697.0(6)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.10\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.30 \times 0.20 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer
Absorption correction: ψ scan (*XCAD4*; Harms & Wocadlo, 1995)
 $T_{\min} = 0.971$, $T_{\max} = 0.990$
3321 measured reflections

3078 independent reflections
2060 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$
3 standard reflections every 200 reflections
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.179$
 $S = 1.01$
3078 reflections

235 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.30\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.24\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C13—H13A \cdots O1	0.93	2.27	2.872 (4)	122
C20—H20A \cdots O4	0.93	2.18	3.019 (4)	150

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

We greatly appreciate the financial support of the Foundation of the Key Laboratory for Advanced Technology in Environmental Protection of Jiangsu Province (AE201017).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FF2061).

References

- Alonso, R., Castedo, L. & Dominguez, D. (1985). *Tetrahedron Lett.* **26**, 2925–2928.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Michael, J. P. (2004). *Nat. Prod. Rep.* **21**, 625–649.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shen, Y. M., Lv, P. C., Chen, W., Liu, P. G., Zhang, M. Z. & Zhu, H. L. (2010). *Eur. J. Med. Chem.* **45**, 3184–3190.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Sriram, D., Yogeeshwari, P., Thirumurugan, R. & Bal, T. R. (2005). *Nat. Prod. Res.* **19**, 393–412.

supplementary materials

Acta Cryst. (2012). E68, o1430 [doi:10.1107/S1600536812015656]

Dimethyl 3-(cyclopropylcarbonyl)pyrrolo[2,1-a]isoquinoline-1,2-dicarboxylate

Honglong Xing, Fan Tang and Wei Wang

Comment

The indolizine and hydrogenated indolizine structures are found in many alkaloids such as amorine, crythraline, swainsonine, cryptaustoline, cryptowoline (Sriram *et al.* 2005), camptothesin (Michael, 2004), nuevamine (Alonso *et al.*, 1985), *etc.* These natural and many synthetic indolizine derivatives have been found to have a variety of biological activity (Shen *et al.*, 2010).

We report here the synthesis and crystal structure of the title compound (I). The molecular structure of (I) is shown in Fig. 1. In the title compound, intramolecular C13—H13A···O1 and C20—H20A···O4 hydrogen bond generates extra two ring motifs. The dihedral angle between the indolizine ring system and the C20—H20A···O4 plane is 20.40 (4)°.

Experimental


A mixture of the 2-(2-cyclopropyl-2-oxoethyl)isoquinolinium bromide (10 mmol), acrylonitrile (40 mmol), triethylamine (2 ml) and TPCD (4 g) in DMF (40 ml) was heated at 90° C for 5 h. After cooling, the reaction mixture was poured into an aqueous hydrochloric acid solution (5%, 100 ml), the precipitated crude product was collected by filtration and further purified by silica gel column chromatography with petroleum ether (bp 60–90 °C)-ethyl acetate as eluents. Yellow crystal. m.p. 388–389 K, Yield 76%. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in petroleum ether -ethyl acetate(4:1), at room temperature.

Refinement

The H atoms were fixed geometrically and were treated as riding on their parent C atoms, with C—H distances in the range of 0.93–0.97 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent atom})$, or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_\text{methyl})$.

Computing details

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software* (Enraf–Nonius, 1989); data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

**Figure 1**

The molecular structure of the title compound, with 30% probability displacement ellipsoids. Dashed lines indicate hydrogen bonds.

Dimethyl 3-(cyclopropylcarbonyl)pyrrolo[2,1-a]isoquinoline-1,2-dicarboxylate

Crystal data

$C_{20}H_{17}NO_5$
 $M_r = 351.35$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 7.5910 (15)$ Å
 $b = 18.436 (4)$ Å
 $c = 12.162 (2)$ Å
 $\beta = 94.43 (3)$ °
 $V = 1697.0 (6)$ Å³
 $Z = 4$

$F(000) = 736$
 $D_x = 1.375$ Mg m⁻³
Melting point: 388 K
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 25 reflections
 $\theta = 9\text{--}12$ °
 $\mu = 0.10$ mm⁻¹
 $T = 293$ K
Block, yellow
 $0.30 \times 0.20 \times 0.10$ mm

Data collection

Enraf–Nonius CAD-4
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 $\omega/2\theta$ scans
Absorption correction: ψ scan
(XCAD4; Harms & Wocadlo, 1995)
 $T_{\min} = 0.971$, $T_{\max} = 0.990$
3321 measured reflections

3078 independent reflections
2060 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$
 $\theta_{\max} = 25.3$ °, $\theta_{\min} = 2.0$ °
 $h = 0 \rightarrow 9$
 $k = 0 \rightarrow 22$
 $l = -14 \rightarrow 14$
3 standard reflections every 200 reflections
intensity decay: 1%

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.056$$

$$wR(F^2) = 0.179$$

$$S = 1.01$$

3078 reflections

235 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 0.4P]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.30 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.24 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N	0.2043 (3)	0.45752 (11)	0.56131 (17)	0.0407 (5)
O1	0.0490 (3)	0.31335 (12)	0.57398 (18)	0.0729 (7)
C1	0.0128 (5)	0.2518 (2)	0.7977 (3)	0.0763 (10)
H1A	0.0035	0.2474	0.8765	0.092*
H1B	-0.0987	0.2485	0.7533	0.092*
O2	0.3127 (2)	0.44915 (12)	0.92593 (15)	0.0583 (6)
C2	0.1687 (5)	0.22111 (19)	0.7524 (4)	0.0815 (11)
H2A	0.1537	0.1991	0.6799	0.098*
H2B	0.2560	0.1979	0.8032	0.098*
O3	0.0196 (3)	0.44054 (13)	0.88912 (17)	0.0657 (6)
C3	0.1495 (4)	0.30195 (16)	0.7597 (3)	0.0591 (8)
H3A	0.2247	0.3260	0.8179	0.071*
O4	0.4069 (3)	0.64344 (13)	0.76734 (18)	0.0715 (7)
C4	0.1160 (4)	0.34276 (15)	0.6567 (2)	0.0463 (7)
O5	0.1973 (3)	0.59404 (11)	0.86128 (17)	0.0613 (6)
C5	0.2880 (5)	0.4387 (3)	1.0410 (3)	0.0879 (13)
H5A	0.4009	0.4381	1.0824	0.132*
H5B	0.2287	0.3934	1.0507	0.132*
H5C	0.2179	0.4776	1.0667	0.132*
C6	0.1646 (4)	0.45114 (15)	0.8593 (2)	0.0462 (7)
C7	0.2330 (5)	0.64687 (18)	0.9471 (3)	0.0694 (10)
H7A	0.1510	0.6407	1.0026	0.104*
H7B	0.2206	0.6947	0.9164	0.104*
H7C	0.3513	0.6405	0.9795	0.104*
C8	0.3006 (3)	0.59626 (15)	0.7761 (2)	0.0452 (7)
C9	0.2639 (3)	0.53321 (14)	0.7048 (2)	0.0403 (6)

C10	0.2004 (3)	0.46707 (14)	0.7437 (2)	0.0403 (6)
C11	0.1670 (3)	0.41976 (14)	0.6564 (2)	0.0405 (6)
C12	0.2688 (3)	0.52648 (14)	0.5891 (2)	0.0406 (6)
C13	0.1793 (3)	0.43333 (16)	0.4534 (2)	0.0472 (7)
H13A	0.1353	0.3870	0.4387	0.057*
C14	0.2183 (4)	0.47657 (17)	0.3706 (2)	0.0555 (8)
H14A	0.1957	0.4607	0.2983	0.067*
C15	0.2945 (4)	0.54702 (17)	0.3911 (2)	0.0512 (7)
C16	0.3216 (3)	0.57249 (15)	0.5008 (2)	0.0449 (7)
C17	0.3413 (4)	0.5916 (2)	0.3040 (3)	0.0644 (9)
H17A	0.3216	0.5754	0.2317	0.077*
C18	0.4146 (4)	0.6579 (2)	0.3238 (3)	0.0693 (10)
H18A	0.4429	0.6871	0.2654	0.083*
C19	0.4473 (4)	0.68183 (18)	0.4315 (3)	0.0653 (9)
H19A	0.5020	0.7264	0.4449	0.078*
C20	0.4002 (4)	0.64068 (16)	0.5189 (3)	0.0553 (8)
H20A	0.4206	0.6582	0.5905	0.066*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N	0.0414 (12)	0.0421 (12)	0.0390 (12)	0.0036 (10)	0.0048 (9)	-0.0006 (10)
O1	0.0976 (17)	0.0614 (14)	0.0594 (14)	-0.0228 (13)	0.0031 (12)	-0.0076 (11)
C1	0.076 (2)	0.080 (3)	0.075 (2)	-0.0009 (19)	0.0169 (19)	0.0190 (19)
O2	0.0539 (12)	0.0747 (15)	0.0457 (12)	0.0098 (11)	0.0005 (9)	0.0041 (10)
C2	0.101 (3)	0.053 (2)	0.092 (3)	0.0147 (19)	0.022 (2)	0.0156 (19)
O3	0.0523 (12)	0.0861 (16)	0.0604 (13)	-0.0108 (12)	0.0159 (10)	-0.0020 (12)
C3	0.074 (2)	0.0412 (16)	0.061 (2)	-0.0008 (15)	-0.0017 (16)	0.0013 (14)
O4	0.0759 (15)	0.0702 (15)	0.0703 (15)	-0.0316 (13)	0.0186 (12)	-0.0128 (12)
C4	0.0449 (15)	0.0450 (16)	0.0495 (16)	0.0019 (13)	0.0059 (12)	-0.0051 (13)
O5	0.0749 (14)	0.0487 (12)	0.0634 (14)	-0.0122 (10)	0.0248 (11)	-0.0177 (10)
C5	0.092 (3)	0.130 (4)	0.0406 (19)	0.021 (3)	-0.0004 (17)	0.008 (2)
C6	0.0499 (16)	0.0419 (15)	0.0471 (16)	0.0023 (13)	0.0060 (13)	-0.0026 (12)
C7	0.086 (2)	0.062 (2)	0.062 (2)	-0.0102 (18)	0.0194 (18)	-0.0194 (17)
C8	0.0352 (13)	0.0472 (16)	0.0531 (17)	-0.0003 (13)	0.0036 (12)	0.0001 (13)
C9	0.0349 (13)	0.0427 (15)	0.0437 (15)	0.0056 (11)	0.0049 (11)	-0.0003 (12)
C10	0.0379 (13)	0.0417 (15)	0.0413 (15)	0.0045 (11)	0.0035 (11)	0.0006 (11)
C11	0.0373 (13)	0.0423 (15)	0.0421 (15)	0.0050 (11)	0.0038 (11)	0.0011 (12)
C12	0.0339 (13)	0.0402 (14)	0.0479 (16)	0.0071 (11)	0.0039 (11)	0.0020 (12)
C13	0.0448 (15)	0.0531 (17)	0.0428 (16)	0.0060 (13)	-0.0014 (12)	-0.0032 (13)
C14	0.0547 (17)	0.069 (2)	0.0426 (17)	0.0116 (15)	0.0020 (13)	-0.0036 (15)
C15	0.0462 (15)	0.0619 (19)	0.0461 (17)	0.0136 (14)	0.0069 (12)	0.0123 (14)
C16	0.0330 (13)	0.0472 (16)	0.0551 (17)	0.0110 (12)	0.0066 (11)	0.0095 (13)
C17	0.0523 (18)	0.083 (2)	0.059 (2)	0.0119 (17)	0.0109 (14)	0.0175 (18)
C18	0.057 (2)	0.078 (2)	0.075 (2)	0.0110 (18)	0.0161 (17)	0.034 (2)
C19	0.0588 (19)	0.0517 (18)	0.087 (3)	0.0033 (15)	0.0161 (17)	0.0196 (18)
C20	0.0491 (16)	0.0516 (17)	0.066 (2)	0.0040 (14)	0.0112 (14)	0.0080 (15)

Geometric parameters (\AA , $^\circ$)

N—C13	1.385 (3)	C6—C10	1.483 (4)
N—C12	1.395 (3)	C7—H7A	0.9600
N—C11	1.398 (3)	C7—H7B	0.9600
O1—C4	1.218 (3)	C7—H7C	0.9600
C1—C2	1.458 (5)	C8—C9	1.464 (4)
C1—C3	1.490 (5)	C9—C10	1.406 (4)
C1—H1A	0.9700	C9—C12	1.416 (4)
C1—H1B	0.9700	C10—C11	1.382 (4)
O2—C6	1.334 (3)	C12—C16	1.449 (4)
O2—C5	1.440 (4)	C13—C14	1.335 (4)
C2—C3	1.501 (4)	C13—H13A	0.9300
C2—H2A	0.9700	C14—C15	1.436 (4)
C2—H2B	0.9700	C14—H14A	0.9300
O3—C6	1.201 (3)	C15—C17	1.408 (4)
C3—C4	1.467 (4)	C15—C16	1.414 (4)
C3—H3A	0.9800	C16—C20	1.402 (4)
O4—C8	1.197 (3)	C17—C18	1.356 (5)
C4—C11	1.472 (4)	C17—H17A	0.9300
O5—C8	1.347 (3)	C18—C19	1.387 (5)
O5—C7	1.438 (3)	C18—H18A	0.9300
C5—H5A	0.9600	C19—C20	1.376 (4)
C5—H5B	0.9600	C19—H19A	0.9300
C5—H5C	0.9600	C20—H20A	0.9300
C13—N—C12	122.9 (2)	H7B—C7—H7C	109.5
C13—N—C11	127.0 (2)	O4—C8—O5	121.8 (3)
C12—N—C11	110.1 (2)	O4—C8—C9	128.5 (3)
C2—C1—C3	61.2 (2)	O5—C8—C9	109.6 (2)
C2—C1—H1A	117.6	C10—C9—C12	107.1 (2)
C3—C1—H1A	117.6	C10—C9—C8	122.9 (2)
C2—C1—H1B	117.6	C12—C9—C8	129.8 (2)
C3—C1—H1B	117.6	C11—C10—C9	109.7 (2)
H1A—C1—H1B	114.8	C11—C10—C6	124.6 (2)
C6—O2—C5	115.3 (2)	C9—C10—C6	125.7 (2)
C1—C2—C3	60.5 (2)	C10—C11—N	106.5 (2)
C1—C2—H2A	117.7	C10—C11—C4	129.8 (2)
C3—C2—H2A	117.7	N—C11—C4	123.5 (2)
C1—C2—H2B	117.7	N—C12—C9	106.6 (2)
C3—C2—H2B	117.7	N—C12—C16	117.8 (2)
H2A—C2—H2B	114.8	C9—C12—C16	135.7 (3)
C4—C3—C1	120.3 (3)	C14—C13—N	120.0 (3)
C4—C3—C2	118.0 (3)	C14—C13—H13A	120.0
C1—C3—C2	58.3 (2)	N—C13—H13A	120.0
C4—C3—H3A	116.0	C13—C14—C15	121.3 (3)
C1—C3—H3A	116.0	C13—C14—H14A	119.4
C2—C3—H3A	116.0	C15—C14—H14A	119.4
O1—C4—C3	120.7 (3)	C17—C15—C16	119.3 (3)
O1—C4—C11	121.4 (3)	C17—C15—C14	121.3 (3)

C3—C4—C11	117.9 (2)	C16—C15—C14	119.4 (3)
C8—O5—C7	116.7 (2)	C20—C16—C15	118.4 (3)
O2—C5—H5A	109.5	C20—C16—C12	123.1 (3)
O2—C5—H5B	109.5	C15—C16—C12	118.5 (3)
H5A—C5—H5B	109.5	C18—C17—C15	121.1 (3)
O2—C5—H5C	109.5	C18—C17—H17A	119.4
H5A—C5—H5C	109.5	C15—C17—H17A	119.4
H5B—C5—H5C	109.5	C17—C18—C19	119.7 (3)
O3—C6—O2	124.1 (3)	C17—C18—H18A	120.1
O3—C6—C10	123.9 (3)	C19—C18—H18A	120.1
O2—C6—C10	111.9 (2)	C20—C19—C18	121.1 (3)
O5—C7—H7A	109.5	C20—C19—H19A	119.5
O5—C7—H7B	109.5	C18—C19—H19A	119.5
H7A—C7—H7B	109.5	C19—C20—C16	120.4 (3)
O5—C7—H7C	109.5	C19—C20—H20A	119.8
H7A—C7—H7C	109.5	C16—C20—H20A	119.8
C2—C1—C3—C4	106.2 (3)	C3—C4—C11—C10	-19.1 (4)
C1—C2—C3—C4	-110.1 (3)	O1—C4—C11—N	-23.9 (4)
C1—C3—C4—O1	-44.0 (4)	C3—C4—C11—N	155.6 (2)
C2—C3—C4—O1	23.8 (5)	C13—N—C12—C9	175.8 (2)
C1—C3—C4—C11	136.5 (3)	C11—N—C12—C9	-2.7 (3)
C2—C3—C4—C11	-155.7 (3)	C13—N—C12—C16	-3.9 (3)
C5—O2—C6—O3	-4.7 (4)	C11—N—C12—C16	177.5 (2)
C5—O2—C6—C10	176.1 (3)	C10—C9—C12—N	1.4 (3)
C7—O5—C8—O4	-6.1 (4)	C8—C9—C12—N	-173.7 (2)
C7—O5—C8—C9	172.0 (3)	C10—C9—C12—C16	-178.9 (3)
O4—C8—C9—C10	150.7 (3)	C8—C9—C12—C16	6.0 (5)
O5—C8—C9—C10	-27.3 (3)	C12—N—C13—C14	0.4 (4)
O4—C8—C9—C12	-34.9 (5)	C11—N—C13—C14	178.7 (2)
O5—C8—C9—C12	147.1 (3)	N—C13—C14—C15	3.1 (4)
C12—C9—C10—C11	0.3 (3)	C13—C14—C15—C17	177.8 (3)
C8—C9—C10—C11	175.9 (2)	C13—C14—C15—C16	-2.9 (4)
C12—C9—C10—C6	-176.5 (2)	C17—C15—C16—C20	-2.2 (4)
C8—C9—C10—C6	-1.0 (4)	C14—C15—C16—C20	178.5 (2)
O3—C6—C10—C11	-63.4 (4)	C17—C15—C16—C12	178.6 (2)
O2—C6—C10—C11	115.8 (3)	C14—C15—C16—C12	-0.8 (4)
O3—C6—C10—C9	113.0 (3)	N—C12—C16—C20	-175.2 (2)
O2—C6—C10—C9	-67.8 (3)	C9—C12—C16—C20	5.1 (4)
C9—C10—C11—N	-2.0 (3)	N—C12—C16—C15	4.0 (3)
C6—C10—C11—N	174.9 (2)	C9—C12—C16—C15	-175.7 (3)
C9—C10—C11—C4	173.5 (2)	C16—C15—C17—C18	1.2 (4)
C6—C10—C11—C4	-9.6 (4)	C14—C15—C17—C18	-179.4 (3)
C13—N—C11—C10	-175.5 (2)	C15—C17—C18—C19	1.2 (5)
C12—N—C11—C10	2.9 (3)	C17—C18—C19—C20	-2.6 (5)
C13—N—C11—C4	8.6 (4)	C18—C19—C20—C16	1.6 (5)
C12—N—C11—C4	-172.9 (2)	C15—C16—C20—C19	0.8 (4)
O1—C4—C11—C10	161.3 (3)	C12—C16—C20—C19	-180.0 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C13—H13A···O1	0.93	2.27	2.872 (4)	122
C20—H20A···O4	0.93	2.18	3.019 (4)	150