

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[[diaqua(1,10-phenanthroline- κ^2N,N')zinc]- μ -4-hydroxy-3-sulfonatobenzoato- $\kappa^2O^3:O^1$] sesquihydrate]

Xiang-Qian Fang,^a Shan Gao^a and Seik Weng Ng^{b,c*}

^aKey Laboratory of Functional Inorganic Material Chemistry, Ministry of Education, Heilongjiang University, Harbin 150080, People's Republic of China, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: seikweng@um.edu.my


Received 23 April 2012; accepted 26 April 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.035; wR factor = 0.104; data-to-parameter ratio = 14.5.

The 1,10-phenanthroline-chelated Zn atom in the polymeric title compound, $\{[Zn(C_7H_4O_6S)(C_{12}H_8N_2)(H_2O)_2] \cdot 1.5H_2O\}_n$, is connected to the sulfonate O atom of one 4-hydroxy-3-sulfonatobenzoate dianion and to the carboxylate O atom of another dianion. It is also coordinated by two water molecules in an overall octahedral environment. The dianion links adjacent metal atoms into a chain running along [110]. The chains are linked by O—H...O hydrogen bonds into a three-dimensional network.

Related literature

For the isostructural Mn^{II} derivative, see: Fang *et al.* (2011) and for the isostructural Co^{II} derivative, see: Fang *et al.* (2012).



Experimental

Crystal data

$[Zn(C_7H_4O_6S)(C_{12}H_8N_2)(H_2O)_2] \cdot 1.5H_2O$
 $M_r = 524.79$
 Monoclinic, $C2/c$

$a = 8.3682$ (3) Å
 $b = 17.3251$ (6) Å
 $c = 28.6686$ (10) Å
 $\beta = 92.848$ (1)°

$V = 4151.2$ (3) Å³
 $Z = 8$
 Mo $K\alpha$ radiation

$\mu = 1.34$ mm⁻¹
 $T = 293$ K
 $0.19 \times 0.15 \times 0.11$ mm

Data collection

Rigaku R-Axis RAPID IP diffractometer
 Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
 $T_{min} = 0.784$, $T_{max} = 0.866$

20011 measured reflections
 4741 independent reflections
 3098 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.104$
 $S = 1.16$
 4741 reflections
 326 parameters
 8 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{max} = 0.62$ e Å⁻³
 $\Delta\rho_{min} = -0.88$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O4—H4...O4w	0.84 (1)	1.78 (1)	2.615 (3)	172 (4)
O1w—H11...O6 ⁱ	0.84 (1)	1.68 (1)	2.520 (3)	172 (5)
O1w—H12...O3w	0.84 (1)	1.98 (2)	2.793 (2)	164 (4)
O2w—H21...O2 ⁱⁱ	0.84 (1)	1.94 (1)	2.760 (3)	166 (4)
O2w—H22...O1w ⁱⁱⁱ	0.84 (1)	1.93 (1)	2.767 (3)	174 (4)
O3w—H31...O2	0.84 (1)	1.95 (2)	2.752 (3)	160 (4)
O4w—H41...O4 ^{iv}	0.84 (1)	2.23 (3)	2.939 (3)	142 (4)
O4w—H42...O5 ^v	0.84 (1)	1.95 (1)	2.789 (3)	177 (4)

Symmetry codes: (i) $x - \frac{1}{2}, y - \frac{1}{2}, z$; (ii) $x - 1, y, z$; (iii) $-x + 1, y, -z + \frac{1}{2}$; (iv) $-x + \frac{3}{2}, -y + \frac{1}{2}, -z + 1$; (v) $-x + 2, -y + 1, -z + 1$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MS, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

This work was supported by the Key Project of the Natural Science Foundation of Heilongjiang Province (No. ZD200903), the Key Project of the Education Bureau of Heilongjiang Province (Nos. 12511z023 and 2011CJHB006), the Innovation Team of the Education Bureau of Heilongjiang Province (No. 2010 t d03), Heilongjiang University (Hdtd2010-04) and the Ministry of Higher Education of Malaysia (grant No. UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5893).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Fang, X.-Q., Chen, P.-G., Zhu, Z.-B., Deng, Z.-P. & Gao, S. (2011). *Chin. J. Inorg. Chem.* **27**, 1733–1737.
 Fang, X.-Q., Gao, S. & Ng, S. W. (2012). *Acta Cryst.* **E68**, m721.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku/MS (2002). *CrystalClear*. Rigaku/MS Inc., The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, m722 [doi:10.1107/S1600536812018740]

catena-Poly[[[diaqua(1,10-phenanthroline- κ^2N,N')zinc]- μ -4-hydroxy-3-sulfonatobenzoato- $\kappa^2O^3:O^1$] sesquihydrate]

Xiang-Qian Fang, Shan Gao and Seik Weng Ng

Comment

The title zinc compound (Scheme I, Fig. 1) is isostructural with the manganese(II) derivative (Fang *et al.*, 2011) and with the cobalt(II) derivative (Fang *et al.*, 2012). The 1,10-phenanthroline chelated Zn atom is connected to the sulfonate O atom of one (C₇H₄O₆S) dianion and to the carboxylate O atom of another dianion. It is also coordinated by two water molecules in an octahedral environment. The dianion links adjacent metal atoms into a chain along [1 1 0]. The chains are linked by O–H \cdots O hydrogen bonds into a three-dimensional network (Table 1).

Experimental

A methanol solution (5 ml) of 1,10-phenanthroline (1 mmol) was added to an aqueous solution (10 ml) of zinc(II) dichloride (1 mmol), 2-hydroxy-5-carboxybenzenesulfonic acid (2 mmol) and lithium hydroxide (4 mmol). Colorless crystals were isolated from the solution after several days.


Refinement

All H atoms were located in a difference map. Carbon-bound H-atoms were placed in calculated positions (C–H 0.93 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U(C)$. H-atoms bonded to O were isotropically refined with a distance restraint of O–H 0.84 \pm 0.01 Å.

Omitted owing to bad disagreement were (4 0 4), (5 17 4), (-7 11 20), (4 10 27) and (5 11 26).

Computing details

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO* (Rigaku, 1998); data reduction: *CrystalClear* (Rigaku/MSK, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).


Figure 1

Anisotropic displacement ellipsoid plot (Barbour, 2001) of a portion of the structure of polymeric $[\text{Zn}(\text{H}_2\text{O})_2(\text{C}_{12}\text{H}_8\text{N}_2)(\text{C}_7\text{H}_4\text{O}_6\text{S})]_n \cdot 1.5n\text{H}_2\text{O}$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

catena-Poly[[[diaqua(1,10-phenanthroline- κ^2N,N')zinc]- μ -4-hydroxy-3-sulfonatobenzoato- $\kappa^2O^3:O^1$] sesquihydrate]

Crystal data

$[\text{Zn}(\text{C}_7\text{H}_4\text{O}_6\text{S})(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_2] \cdot 1.5\text{H}_2\text{O}$

$M_r = 524.79$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 8.3682\ (3)\ \text{\AA}$

$b = 17.3251\ (6)\ \text{\AA}$

$c = 28.6686\ (10)\ \text{\AA}$

$\beta = 92.848\ (1)^\circ$

$V = 4151.2\ (3)\ \text{\AA}^3$

$Z = 8$

$F(000) = 2152$

$D_x = 1.679\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 12500 reflections

$\theta = 3.0\text{--}27.5^\circ$

$\mu = 1.34\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Prism, colorless

$0.19 \times 0.15 \times 0.11\ \text{mm}$

Data collection

Rigaku R-AXIS RAPID IP

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scan

Absorption correction: multi-scan

(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.784$, $T_{\max} = 0.866$

20011 measured reflections

4741 independent reflections

3098 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.051$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -10 \rightarrow 10$

$k = -18 \rightarrow 22$

$l = -37 \rightarrow 37$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

$wR(F^2) = 0.104$

$S = 1.16$

4741 reflections

326 parameters

8 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0376P)^2 + 5.6207P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$$\Delta\rho_{\max} = 0.62 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.87 \text{ e } \text{\AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.61591 (4)	0.232574 (19)	0.342240 (11)	0.02942 (12)
S1	1.01800 (9)	0.26433 (4)	0.37668 (3)	0.02952 (18)
O1	0.8591 (2)	0.23027 (12)	0.37134 (7)	0.0340 (5)
O2	1.0907 (3)	0.27321 (13)	0.33170 (8)	0.0489 (7)
O3	1.1182 (3)	0.22346 (14)	0.41081 (9)	0.0523 (7)
O4	0.8751 (3)	0.31120 (12)	0.46469 (7)	0.0415 (6)
O5	1.0863 (3)	0.62949 (11)	0.37689 (7)	0.0331 (5)
O6	1.1071 (3)	0.54964 (13)	0.31619 (8)	0.0518 (7)
O1W	0.6923 (3)	0.17550 (13)	0.28105 (7)	0.0314 (5)
O2W	0.3893 (3)	0.22125 (19)	0.30948 (9)	0.0579 (8)
O3W	1.0000	0.1982 (2)	0.2500	0.0491 (9)
O4W	0.7309 (3)	0.35044 (14)	0.53999 (8)	0.0421 (6)
N1	0.5209 (3)	0.30126 (15)	0.39576 (8)	0.0332 (6)
N2	0.6531 (3)	0.34633 (15)	0.31609 (9)	0.0374 (6)
C1	0.9919 (4)	0.35998 (16)	0.39722 (10)	0.0280 (6)
C2	0.9203 (4)	0.37230 (17)	0.43963 (10)	0.0313 (7)
C3	0.8988 (4)	0.44787 (18)	0.45453 (11)	0.0385 (8)
H3	0.8495	0.4571	0.4824	0.046*
C4	0.9496 (4)	0.50923 (17)	0.42853 (11)	0.0369 (8)
H4A	0.9348	0.5593	0.4392	0.044*
C5	1.0230 (4)	0.49733 (17)	0.38647 (10)	0.0318 (7)
C6	1.0417 (4)	0.42181 (17)	0.37114 (10)	0.0302 (7)
H6	1.0885	0.4128	0.3429	0.036*
C7	1.0760 (4)	0.56319 (17)	0.35757 (10)	0.0332 (7)
C8	0.5305 (4)	0.37811 (19)	0.38722 (11)	0.0377 (8)
C9	0.4569 (4)	0.2779 (2)	0.43478 (12)	0.0429 (8)
H9	0.4499	0.2252	0.4406	0.051*
C10	0.3996 (5)	0.3299 (3)	0.46746 (13)	0.0599 (12)
H10	0.3567	0.3118	0.4947	0.072*
C11	0.4075 (5)	0.4066 (3)	0.45891 (15)	0.0636 (12)
H11A	0.3691	0.4414	0.4803	0.076*
C12	0.4733 (5)	0.4339 (2)	0.41789 (14)	0.0517 (10)
C13	0.4894 (6)	0.5136 (2)	0.40649 (19)	0.0738 (14)
H13	0.4525	0.5509	0.4267	0.089*
C14	0.5563 (6)	0.5357 (2)	0.3673 (2)	0.0795 (15)
H14	0.5647	0.5881	0.3609	0.095*
C15	0.6156 (5)	0.4811 (2)	0.33509 (15)	0.0564 (11)
C16	0.6890 (6)	0.5010 (2)	0.29391 (17)	0.0740 (14)
H16	0.7015	0.5528	0.2862	0.089*
C17	0.7418 (6)	0.4454 (3)	0.26522 (15)	0.0701 (13)
H17	0.7913	0.4584	0.2380	0.084*
C18	0.7207 (5)	0.3679 (2)	0.27727 (12)	0.0506 (10)
H18	0.7555	0.3300	0.2572	0.061*

C19	0.6008 (4)	0.40212 (18)	0.34510 (11)	0.0376 (8)
H4	0.832 (5)	0.328 (2)	0.4885 (9)	0.066 (13)*
H11	0.665 (6)	0.1314 (13)	0.2903 (16)	0.092 (17)*
H12	0.7903 (16)	0.176 (2)	0.2762 (13)	0.054 (12)*
H21	0.301 (3)	0.232 (2)	0.3201 (14)	0.060 (12)*
H22	0.368 (5)	0.204 (2)	0.2826 (7)	0.066 (13)*
H31	1.023 (5)	0.2307 (17)	0.2710 (10)	0.055 (12)*
H41	0.669 (4)	0.3148 (17)	0.5466 (14)	0.064 (13)*
H42	0.784 (4)	0.358 (2)	0.5651 (8)	0.060 (13)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0375 (2)	0.02202 (19)	0.02921 (19)	-0.00039 (15)	0.00628 (15)	-0.00082 (14)
S1	0.0306 (4)	0.0221 (4)	0.0363 (4)	-0.0016 (3)	0.0063 (3)	-0.0058 (3)
O1	0.0303 (11)	0.0265 (11)	0.0448 (12)	-0.0081 (9)	-0.0009 (10)	-0.0059 (9)
O2	0.0613 (16)	0.0375 (14)	0.0506 (14)	-0.0070 (12)	0.0310 (13)	-0.0130 (11)
O3	0.0504 (16)	0.0372 (14)	0.0673 (17)	0.0087 (12)	-0.0172 (13)	-0.0003 (12)
O4	0.0697 (17)	0.0242 (11)	0.0324 (12)	-0.0106 (11)	0.0196 (12)	-0.0017 (9)
O5	0.0482 (14)	0.0223 (11)	0.0292 (10)	-0.0077 (10)	0.0066 (10)	-0.0020 (9)
O6	0.094 (2)	0.0289 (13)	0.0343 (12)	-0.0179 (13)	0.0245 (13)	-0.0051 (10)
O1W	0.0369 (13)	0.0288 (12)	0.0295 (11)	-0.0043 (10)	0.0108 (10)	-0.0007 (9)
O2W	0.0366 (15)	0.100 (2)	0.0372 (14)	0.0095 (15)	0.0010 (13)	-0.0240 (15)
O3W	0.047 (2)	0.055 (2)	0.045 (2)	0.000	0.0070 (19)	0.000
O4W	0.0600 (17)	0.0377 (14)	0.0292 (12)	-0.0082 (12)	0.0086 (12)	-0.0005 (10)
N1	0.0329 (14)	0.0366 (15)	0.0302 (13)	0.0011 (12)	0.0038 (12)	-0.0056 (11)
N2	0.0476 (17)	0.0278 (14)	0.0372 (14)	0.0006 (12)	0.0052 (13)	0.0011 (11)
C1	0.0295 (16)	0.0228 (15)	0.0319 (15)	-0.0045 (12)	0.0043 (13)	-0.0046 (12)
C2	0.0408 (18)	0.0253 (15)	0.0283 (15)	-0.0090 (13)	0.0061 (14)	-0.0004 (12)
C3	0.058 (2)	0.0294 (17)	0.0296 (16)	-0.0062 (15)	0.0174 (16)	-0.0054 (13)
C4	0.055 (2)	0.0228 (16)	0.0336 (16)	-0.0083 (15)	0.0118 (16)	-0.0082 (13)
C5	0.0423 (19)	0.0231 (15)	0.0305 (15)	-0.0064 (13)	0.0067 (14)	-0.0005 (12)
C6	0.0371 (17)	0.0295 (16)	0.0249 (14)	-0.0059 (13)	0.0094 (13)	-0.0027 (12)
C7	0.0427 (19)	0.0251 (16)	0.0323 (16)	-0.0073 (14)	0.0069 (15)	-0.0047 (13)
C8	0.0375 (18)	0.0310 (17)	0.0438 (18)	0.0069 (14)	-0.0054 (15)	-0.0119 (15)
C9	0.0344 (18)	0.056 (2)	0.0386 (18)	0.0004 (16)	0.0027 (15)	-0.0060 (16)
C10	0.046 (2)	0.091 (4)	0.044 (2)	0.000 (2)	0.0122 (19)	-0.019 (2)
C11	0.052 (3)	0.079 (3)	0.060 (3)	0.013 (2)	0.004 (2)	-0.038 (2)
C12	0.046 (2)	0.050 (2)	0.059 (2)	0.0124 (18)	-0.0039 (19)	-0.0237 (19)
C13	0.081 (3)	0.044 (3)	0.095 (4)	0.023 (2)	-0.004 (3)	-0.034 (3)
C14	0.106 (4)	0.025 (2)	0.107 (4)	0.009 (2)	-0.004 (3)	-0.014 (2)
C15	0.068 (3)	0.0264 (18)	0.074 (3)	0.0044 (18)	-0.007 (2)	0.0001 (18)
C16	0.105 (4)	0.032 (2)	0.085 (3)	-0.009 (2)	0.002 (3)	0.019 (2)
C17	0.099 (4)	0.050 (3)	0.061 (3)	-0.013 (3)	0.011 (3)	0.020 (2)
C18	0.070 (3)	0.037 (2)	0.045 (2)	-0.0027 (18)	0.013 (2)	0.0070 (16)
C19	0.042 (2)	0.0239 (16)	0.0462 (19)	0.0049 (14)	-0.0069 (16)	-0.0018 (14)

Geometric parameters (Å, °)

Zn1—O5 ⁱ	2.065 (2)	C2—C3	1.392 (4)
Zn1—O2W	2.083 (3)	C3—C4	1.378 (4)
Zn1—N1	2.127 (2)	C3—H3	0.9300
Zn1—N2	2.137 (3)	C4—C5	1.395 (4)
Zn1—O1W	2.139 (2)	C4—H4A	0.9300
Zn1—O1	2.161 (2)	C5—C6	1.392 (4)
S1—O3	1.442 (3)	C5—C7	1.491 (4)
S1—O1	1.456 (2)	C6—H6	0.9300
S1—O2	1.461 (2)	C8—C12	1.407 (4)
S1—C1	1.776 (3)	C8—C19	1.431 (5)
O4—C2	1.344 (3)	C9—C10	1.401 (5)
O4—H4	0.839 (10)	C9—H9	0.9300
O5—C7	1.276 (3)	C10—C11	1.354 (6)
O5—Zn1 ⁱⁱ	2.065 (2)	C10—H10	0.9300
O6—C7	1.249 (4)	C11—C12	1.405 (6)
O1W—H11	0.844 (10)	C11—H11A	0.9300
O1W—H12	0.839 (10)	C12—C13	1.427 (6)
O2W—H21	0.835 (10)	C13—C14	1.335 (6)
O2W—H22	0.836 (10)	C13—H13	0.9300
O3W—H31	0.839 (10)	C14—C15	1.429 (6)
O4W—H41	0.836 (10)	C14—H14	0.9300
O4W—H42	0.838 (10)	C15—C16	1.401 (6)
N1—C9	1.327 (4)	C15—C19	1.404 (5)
N1—C8	1.357 (4)	C16—C17	1.356 (6)
N2—C18	1.327 (4)	C16—H16	0.9300
N2—C19	1.362 (4)	C17—C18	1.398 (5)
C1—C6	1.383 (4)	C17—H17	0.9300
C1—C2	1.398 (4)	C18—H18	0.9300
O5 ⁱ —Zn1—O2W	90.42 (11)	C3—C4—H4A	119.5
O5 ⁱ —Zn1—N1	94.52 (9)	C5—C4—H4A	119.5
O2W—Zn1—N1	90.76 (10)	C6—C5—C4	118.2 (3)
O5 ⁱ —Zn1—N2	171.77 (9)	C6—C5—C7	120.2 (3)
O2W—Zn1—N2	94.24 (12)	C4—C5—C7	121.6 (3)
N1—Zn1—N2	78.66 (10)	C1—C6—C5	121.0 (3)
O5 ⁱ —Zn1—O1W	92.44 (8)	C1—C6—H6	119.5
O2W—Zn1—O1W	83.65 (9)	C5—C6—H6	119.5
N1—Zn1—O1W	171.10 (10)	O6—C7—O5	124.7 (3)
N2—Zn1—O1W	94.81 (9)	O6—C7—C5	117.8 (3)
O5 ⁱ —Zn1—O1	86.06 (8)	O5—C7—C5	117.6 (3)
O2W—Zn1—O1	172.29 (10)	N1—C8—C12	122.4 (3)
N1—Zn1—O1	96.35 (9)	N1—C8—C19	117.9 (3)
N2—Zn1—O1	90.09 (10)	C12—C8—C19	119.6 (3)
O1W—Zn1—O1	89.65 (8)	N1—C9—C10	122.3 (4)
O3—S1—O1	111.69 (15)	N1—C9—H9	118.9
O3—S1—O2	113.39 (16)	C10—C9—H9	118.9
O1—S1—O2	111.59 (14)	C11—C10—C9	119.2 (4)
O3—S1—C1	108.07 (14)	C11—C10—H10	120.4

O1—S1—C1	106.63 (13)	C9—C10—H10	120.4
O2—S1—C1	104.94 (13)	C10—C11—C12	120.5 (3)
S1—O1—Zn1	150.55 (14)	C10—C11—H11A	119.8
C2—O4—H4	108 (3)	C12—C11—H11A	119.8
C7—O5—Zn1 ⁱⁱ	125.22 (19)	C11—C12—C8	116.8 (4)
Zn1—O1W—H11	94 (3)	C11—C12—C13	124.3 (4)
Zn1—O1W—H12	118 (3)	C8—C12—C13	118.9 (4)
H11—O1W—H12	111 (4)	C14—C13—C12	121.2 (4)
Zn1—O2W—H21	128 (3)	C14—C13—H13	119.4
Zn1—O2W—H22	127 (3)	C12—C13—H13	119.4
H21—O2W—H22	105 (4)	C13—C14—C15	121.8 (4)
H41—O4W—H42	103 (4)	C13—C14—H14	119.1
C9—N1—C8	118.8 (3)	C15—C14—H14	119.1
C9—N1—Zn1	128.2 (2)	C16—C15—C19	117.4 (4)
C8—N1—Zn1	113.01 (19)	C16—C15—C14	124.2 (4)
C18—N2—C19	118.4 (3)	C19—C15—C14	118.5 (4)
C18—N2—Zn1	129.1 (2)	C17—C16—C15	120.3 (4)
C19—N2—Zn1	112.6 (2)	C17—C16—H16	119.8
C6—C1—C2	120.4 (3)	C15—C16—H16	119.8
C6—C1—S1	119.9 (2)	C16—C17—C18	118.9 (4)
C2—C1—S1	119.7 (2)	C16—C17—H17	120.5
O4—C2—C3	122.2 (3)	C18—C17—H17	120.5
O4—C2—C1	119.3 (3)	N2—C18—C17	122.8 (4)
C3—C2—C1	118.6 (3)	N2—C18—H18	118.6
C4—C3—C2	120.8 (3)	C17—C18—H18	118.6
C4—C3—H3	119.6	N2—C19—C15	122.2 (3)
C2—C3—H3	119.6	N2—C19—C8	117.9 (3)
C3—C4—C5	121.0 (3)	C15—C19—C8	120.0 (3)
O3—S1—O1—Zn1	173.2 (2)	Zn1 ⁱⁱ —O5—C7—C5	-168.2 (2)
O2—S1—O1—Zn1	-58.7 (3)	C6—C5—C7—O6	13.8 (5)
C1—S1—O1—Zn1	55.4 (3)	C4—C5—C7—O6	-164.3 (3)
O5 ⁱ —Zn1—O1—S1	179.0 (3)	C6—C5—C7—O5	-166.3 (3)
N1—Zn1—O1—S1	-86.9 (3)	C4—C5—C7—O5	15.7 (5)
N2—Zn1—O1—S1	-8.3 (3)	C9—N1—C8—C12	-0.9 (5)
O1W—Zn1—O1—S1	86.5 (3)	Zn1—N1—C8—C12	179.4 (3)
O5 ⁱ —Zn1—N1—C9	-4.3 (3)	C9—N1—C8—C19	179.6 (3)
O2W—Zn1—N1—C9	86.1 (3)	Zn1—N1—C8—C19	-0.1 (4)
N2—Zn1—N1—C9	-179.7 (3)	C8—N1—C9—C10	-0.2 (5)
O1—Zn1—N1—C9	-90.9 (3)	Zn1—N1—C9—C10	179.5 (3)
O5 ⁱ —Zn1—N1—C8	175.4 (2)	N1—C9—C10—C11	0.8 (6)
O2W—Zn1—N1—C8	-94.2 (2)	C9—C10—C11—C12	-0.4 (6)
N2—Zn1—N1—C8	0.0 (2)	C10—C11—C12—C8	-0.6 (6)
O1—Zn1—N1—C8	88.8 (2)	C10—C11—C12—C13	-178.7 (4)
O2W—Zn1—N2—C18	-91.4 (3)	N1—C8—C12—C11	1.2 (5)
N1—Zn1—N2—C18	178.7 (3)	C19—C8—C12—C11	-179.2 (3)
O1W—Zn1—N2—C18	-7.4 (3)	N1—C8—C12—C13	179.5 (4)
O1—Zn1—N2—C18	82.3 (3)	C19—C8—C12—C13	-1.0 (5)
O2W—Zn1—N2—C19	90.1 (2)	C11—C12—C13—C14	178.5 (5)

N1—Zn1—N2—C19	0.1 (2)	C8—C12—C13—C14	0.4 (7)
O1w—Zn1—N2—C19	174.0 (2)	C12—C13—C14—C15	-0.2 (8)
O1—Zn1—N2—C19	-96.3 (2)	C13—C14—C15—C16	-179.0 (5)
O3—S1—C1—C6	120.0 (3)	C13—C14—C15—C19	0.6 (7)
O1—S1—C1—C6	-119.8 (3)	C19—C15—C16—C17	0.4 (7)
O2—S1—C1—C6	-1.3 (3)	C14—C15—C16—C17	180.0 (5)
O3—S1—C1—C2	-60.3 (3)	C15—C16—C17—C18	0.5 (8)
O1—S1—C1—C2	59.9 (3)	C19—N2—C18—C17	0.7 (6)
O2—S1—C1—C2	178.4 (3)	Zn1—N2—C18—C17	-177.8 (3)
C6—C1—C2—O4	-179.0 (3)	C16—C17—C18—N2	-1.1 (7)
S1—C1—C2—O4	1.3 (4)	C18—N2—C19—C15	0.3 (5)
C6—C1—C2—C3	0.8 (5)	Zn1—N2—C19—C15	179.1 (3)
S1—C1—C2—C3	-178.9 (3)	C18—N2—C19—C8	-179.0 (3)
O4—C2—C3—C4	178.6 (3)	Zn1—N2—C19—C8	-0.2 (4)
C1—C2—C3—C4	-1.3 (5)	C16—C15—C19—N2	-0.8 (6)
C2—C3—C4—C5	0.4 (6)	C14—C15—C19—N2	179.5 (4)
C3—C4—C5—C6	0.8 (5)	C16—C15—C19—C8	178.5 (4)
C3—C4—C5—C7	178.9 (3)	C14—C15—C19—C8	-1.2 (6)
C2—C1—C6—C5	0.4 (5)	N1—C8—C19—N2	0.3 (5)
S1—C1—C6—C5	-179.8 (3)	C12—C8—C19—N2	-179.3 (3)
C4—C5—C6—C1	-1.2 (5)	N1—C8—C19—C15	-179.1 (3)
C7—C5—C6—C1	-179.4 (3)	C12—C8—C19—C15	1.4 (5)
Zn1 ⁱⁱ —O5—C7—O6	11.8 (5)		

Symmetry codes: (i) $x-1/2, y-1/2, z$; (ii) $x+1/2, y+1/2, z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O4—H4 \cdots O4w	0.84 (1)	1.78 (1)	2.615 (3)	172 (4)
O1w—H11 \cdots O6 ⁱ	0.84 (1)	1.68 (1)	2.520 (3)	172 (5)
O1w—H12 \cdots O3w	0.84 (1)	1.98 (2)	2.793 (2)	164 (4)
O2w—H21 \cdots O2 ⁱⁱⁱ	0.84 (1)	1.94 (1)	2.760 (3)	166 (4)
O2w—H22 \cdots O1w ^{iv}	0.84 (1)	1.93 (1)	2.767 (3)	174 (4)
O3w—H31 \cdots O2	0.84 (1)	1.95 (2)	2.752 (3)	160 (4)
O4w—H41 \cdots O4 ^v	0.84 (1)	2.23 (3)	2.939 (3)	142 (4)
O4w—H42 \cdots O5 ^{vi}	0.84 (1)	1.95 (1)	2.789 (3)	177 (4)

Symmetry codes: (i) $x-1/2, y-1/2, z$; (iii) $x-1, y, z$; (iv) $-x+1, y, -z+1/2$; (v) $-x+3/2, -y+1/2, -z+1$; (vi) $-x+2, -y+1, -z+1$.