

Benzyltriethylammonium tetrachlorido-ferrate(III)

Lei Jin

College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China
Correspondence e-mail: jinlei8812@163.com


Received 13 May 2012; accepted 20 May 2012

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.008\text{ \AA}$; disorder in main residue; R factor = 0.068; wR factor = 0.170; data-to-parameter ratio = 18.1.

In the title molecular salt, $(\text{C}_{13}\text{H}_{22}\text{N})[\text{FeCl}_4]$, three of the chloride ions of the tetrahedral Fe^{III} -containing anion are disordered over two orientations in a 0.656 (11):0.344 (11) ratio. In the crystal, there are no identifiable directional interactions between cations and anions except for Coulombic forces.

Related literature

For background to molecular-ionic ferroelectrics, see: Zhang *et al.* (2010).



Experimental

Crystal data

$(\text{C}_{13}\text{H}_{22}\text{N})[\text{FeCl}_4]$	$V = 3764.7 (13)\text{ \AA}^3$
$M_r = 389.97$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 15.514 (3)\text{ \AA}$	$\mu = 1.36\text{ mm}^{-1}$
$b = 15.021 (3)\text{ \AA}$	$T = 293\text{ K}$
$c = 16.155 (3)\text{ \AA}$	$0.26 \times 0.24 \times 0.20\text{ mm}$

Data collection

Rigaku Mercury2 CCD diffractometer	32945 measured reflections
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	3691 independent reflections
$T_{\min} = 0.655$, $T_{\max} = 0.734$	2364 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.082$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.068$	156 restraints
$wR(F^2) = 0.170$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.73\text{ e \AA}^{-3}$
3691 reflections	$\Delta\rho_{\text{min}} = -0.38\text{ e \AA}^{-3}$
204 parameters	

Table 1
Selected bond lengths (\AA).

Fe1—Cl1'	2.178 (4)	Fe1—Cl3'	2.256 (5)
Fe1—Cl2'	2.211 (7)	Fe1—Cl4	2.1821 (17)

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The author thanks the Ordered Matter Science Research Centre, Southeast University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6797).

References

- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhang, W., Ye, H. Y., Cai, H. L., Ge, J. Z., Xiong, R. G. & Huang, S. P. (2010). *J. Am. Chem. Soc.* **132**, 7300–7302.

supplementary materials

Acta Cryst. (2012). E68, m805 [doi:10.1107/S1600536812023008]

Benzyltriethylammonium tetrachloridoferrate(III)

Lei Jin

Comment

In our lab, we have been exploring simple molecular-ionic compounds which have potential phase-transition properties. (Zhang *et al.*, 2010). Herein, the title compound has been synthesized and its crystal structure is reported.

The ions of the title compound, ($C_{13}H_{22}N^+$) $FeCl_4^-$ crystallize in the orthorhombic $Pbca$ space group, an asymmetric unit consists of a tetrachloroferrate anion unit, which contains three disordered Cl1, Cl2, Cl3 atoms with a ratio in 1: 1, and benzyltriethylammonium cations (Scheme 1). (Fig 1). In the anion, the bond distances of Fe—Cl being in the range of 2.083 (7)–2.210 (8) Å and the bond angles of Cl—Fe—Cl being in the range of 93.9 (7)–124.7 (4) °, while the bond distances of disordered Fe—Cl being in the range of 2.178 (4)–2.211 (7) Å and the bond angles of disordered Cl—Fe—Cl range from 90.7 to 127.9 °, which means a largely deviation to the ideal structure. In the structure, the benzyltriethylammonium cations interact with the $FeCl_4^-$ anion through non-covalent interaction-static attracting forces like Coulomb and Van der Waals forces to complete a network structure.

Experimental

In room temperature benzyltriethylammoniumchlorine (10 mmol, 2.28 g) were dissolved in 30 ml water, then a solution with $FeCl_3 \cdot 6H_2O$ (5 mmol, 1.35 g) and excessive hydrochloric acid was dropped slowly into the previous solution with properly stirring. An orange solid appeared immediately and the solid was collected by filtration. Orange blocks were obtained by the slow evaporation of the above filtrate after a week in air.


The dielectric constant of the compound as a function of temperature indicates that the permittivity is basically temperature-independent ($\epsilon = C/(T-T_0)$), suggesting that this compound is not ferroelectric or there may be no distinct phase transition occurring within the measured temperature (below the melting point).

Refinement

H atoms were placed in calculated positions (C—H = 0.93 Å for Csp^2 atoms and C—H = 0.96 Å and 0.97 Å for Csp^3 atoms), assigned fixed U_{iso} values [$U_{iso} = 1.2U_{eq}(Csp^2/N)$ and $1.5U_{eq}(Csp^3)$] and allowed to ride.

Computing details


Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear* (Rigaku, 2005); data reduction: *CrystalClear* (Rigaku, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

**Figure 1**

The molecular structure of the title compound, showing 30% probability displacement ellipsoids.

Benzyltriethylammonium tetrachloroferrate(III)

Crystal data



$M_r = 389.97$

Orthorhombic, $Pbca$

Hall symbol: -P 2ac 2ab

$a = 15.514 (3) \text{ \AA}$

$b = 15.021 (3) \text{ \AA}$

$c = 16.155 (3) \text{ \AA}$

$V = 3764.7 (13) \text{ \AA}^3$

$Z = 8$

$F(000) = 1608$

$D_x = 1.376 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

$\theta = 3.0\text{--}26^\circ$

$\mu = 1.36 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, orange

$0.26 \times 0.24 \times 0.20 \text{ mm}$

Data collection

Rigaku Mercury2 CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 13.6612 pixels mm^{-1}

CCD_Profile_fitting scans

Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005)

$T_{\min} = 0.655$, $T_{\max} = 0.734$

32945 measured reflections

3691 independent reflections

2364 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.082$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -19 \rightarrow 19$

$k = -18 \rightarrow 18$

$l = -19 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.068$

$wR(F^2) = 0.170$

$S = 1.04$

3691 reflections

204 parameters

156 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0625P)^2 + 5.6146P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.009$

$\Delta\rho_{\max} = 0.73 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.38 \text{ e \AA}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0092 (7)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.7598 (3)	0.8752 (4)	0.3510 (3)	0.0664 (14)	
H1A	0.8189	0.8604	0.3369	0.080*	
H1B	0.7231	0.8513	0.3078	0.080*	
C2	0.7509 (4)	0.9749 (4)	0.3512 (4)	0.0816 (17)	
H2A	0.6920	0.9906	0.3619	0.122*	
H2B	0.7679	0.9980	0.2982	0.122*	
H2C	0.7871	0.9996	0.3935	0.122*	
C3	0.6454 (3)	0.8524 (3)	0.4582 (3)	0.0585 (13)	
H3A	0.6316	0.8189	0.5078	0.070*	
H3B	0.6437	0.9150	0.4726	0.070*	
C4	0.5764 (3)	0.8349 (4)	0.3949 (4)	0.0789 (17)	
H4A	0.5873	0.8700	0.3464	0.118*	
H4B	0.5212	0.8506	0.4175	0.118*	
H4C	0.5765	0.7729	0.3804	0.118*	
C5	0.7951 (3)	0.8604 (4)	0.5021 (3)	0.0677 (14)	
H5A	0.7825	0.8254	0.5511	0.081*	
H5B	0.7813	0.9220	0.5146	0.081*	
C6	0.8915 (3)	0.8539 (4)	0.4842 (4)	0.0879 (19)	
H6A	0.9040	0.7971	0.4597	0.132*	
H6B	0.9232	0.8601	0.5349	0.132*	
H6C	0.9079	0.9004	0.4466	0.132*	
C7	0.7465 (3)	0.7295 (3)	0.4183 (3)	0.0590 (13)	
H7A	0.7085	0.7123	0.3733	0.071*	
H7B	0.8051	0.7176	0.4006	0.071*	
C8	0.7268 (3)	0.6716 (3)	0.4920 (3)	0.0574 (12)	
C9	0.7909 (4)	0.6451 (4)	0.5460 (4)	0.0751 (15)	
H9	0.8470	0.6648	0.5377	0.090*	
C10	0.7728 (5)	0.5891 (4)	0.6129 (4)	0.0880 (18)	
H10	0.8164	0.5715	0.6488	0.106*	
C11	0.6901 (5)	0.5607 (4)	0.6247 (4)	0.0864 (18)	
H11	0.6771	0.5245	0.6697	0.104*	
C12	0.6270 (4)	0.5847 (4)	0.5717 (5)	0.0929 (19)	
H12	0.5711	0.5642	0.5800	0.111*	
C13	0.6447 (3)	0.6390 (4)	0.5060 (4)	0.0738 (15)	

H13	0.6005	0.6544	0.4698	0.089*	
Cl1	-0.0663 (7)	0.7414 (8)	0.2546 (8)	0.116 (3)	0.344 (11)
Cl2	0.1300 (9)	0.7070 (9)	0.3047 (8)	0.089 (3)	0.344 (11)
Cl2'	0.1493 (4)	0.6778 (5)	0.2948 (5)	0.0878 (15)	0.656 (11)
Cl3	0.0269 (6)	0.5423 (8)	0.1865 (6)	0.109 (2)	0.344 (11)
Cl4	-0.02775 (10)	0.58556 (13)	0.41086 (9)	0.0936 (6)	
Cl1'	-0.0875 (3)	0.7000 (5)	0.2274 (3)	0.1154 (16)	0.656 (11)
Cl3'	0.0285 (3)	0.4954 (5)	0.2216 (4)	0.1169 (19)	0.656 (11)
Fe1	0.01578 (5)	0.62672 (6)	0.28825 (4)	0.0672 (3)	
N1	0.7371 (2)	0.8292 (3)	0.4325 (2)	0.0527 (10)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.065 (3)	0.081 (4)	0.053 (3)	0.003 (3)	0.013 (2)	0.000 (3)
C2	0.085 (4)	0.081 (4)	0.079 (4)	0.000 (3)	0.004 (3)	0.010 (3)
C3	0.045 (2)	0.064 (3)	0.066 (3)	0.008 (2)	0.007 (2)	-0.019 (3)
C4	0.053 (3)	0.094 (4)	0.090 (4)	0.007 (3)	-0.007 (3)	-0.019 (3)
C5	0.059 (3)	0.082 (4)	0.062 (3)	-0.009 (3)	-0.006 (2)	-0.014 (3)
C6	0.052 (3)	0.109 (5)	0.103 (5)	-0.011 (3)	-0.006 (3)	0.004 (4)
C7	0.053 (3)	0.071 (3)	0.053 (3)	0.005 (2)	0.009 (2)	-0.014 (2)
C8	0.051 (3)	0.066 (3)	0.055 (3)	0.005 (2)	0.005 (2)	-0.013 (2)
C9	0.065 (3)	0.088 (4)	0.072 (4)	0.002 (3)	0.002 (3)	0.000 (3)
C10	0.099 (5)	0.096 (5)	0.070 (4)	0.011 (4)	-0.012 (3)	0.001 (3)
C11	0.105 (5)	0.082 (4)	0.072 (4)	-0.011 (4)	0.015 (4)	0.006 (3)
C12	0.080 (4)	0.091 (5)	0.108 (5)	-0.006 (4)	0.016 (4)	0.010 (4)
C13	0.056 (3)	0.081 (4)	0.085 (4)	-0.002 (3)	0.003 (3)	0.003 (3)
Cl1	0.099 (5)	0.130 (6)	0.120 (5)	0.050 (4)	0.052 (4)	0.027 (4)
Cl2	0.095 (6)	0.112 (6)	0.060 (4)	-0.056 (4)	-0.003 (4)	0.003 (4)
Cl2'	0.079 (2)	0.110 (4)	0.075 (2)	-0.026 (2)	0.0146 (16)	0.000 (2)
Cl3	0.125 (4)	0.123 (5)	0.080 (4)	-0.015 (4)	-0.004 (4)	-0.042 (4)
Cl4	0.0898 (11)	0.1249 (14)	0.0659 (9)	-0.0371 (10)	0.0069 (8)	0.0092 (9)
Cl1'	0.096 (2)	0.153 (4)	0.097 (3)	0.017 (2)	-0.0180 (19)	0.018 (2)
Cl3'	0.103 (2)	0.142 (4)	0.105 (3)	-0.004 (2)	0.010 (2)	-0.063 (3)
Fe1	0.0567 (5)	0.0958 (7)	0.0492 (4)	-0.0014 (4)	0.0059 (3)	-0.0069 (4)
N1	0.046 (2)	0.064 (3)	0.048 (2)	0.0034 (18)	0.0056 (17)	-0.0143 (19)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.503 (8)	C7—N1	1.522 (6)
C1—N1	1.528 (6)	C7—H7A	0.9700
C1—H1A	0.9700	C7—H7B	0.9700
C1—H1B	0.9700	C8—C9	1.381 (7)
C2—H2A	0.9600	C8—C13	1.384 (7)
C2—H2B	0.9600	C9—C10	1.397 (8)
C2—H2C	0.9600	C9—H9	0.9300
C3—C4	1.505 (7)	C10—C11	1.366 (9)
C3—N1	1.521 (5)	C10—H10	0.9300
C3—H3A	0.9700	C11—C12	1.350 (9)
C3—H3B	0.9700	C11—H11	0.9300

C4—H4A	0.9600	C12—C13	1.367 (8)
C4—H4B	0.9600	C12—H12	0.9300
C4—H4C	0.9600	C13—H13	0.9300
C5—N1	1.516 (6)	Cl1—Fe1	2.210 (8)
C5—C6	1.526 (7)	Cl2—Fe1	2.160 (12)
C5—H5A	0.9700	Cl3—Fe1	2.083 (7)
C5—H5B	0.9700	Fe1—Cl1'	2.178 (4)
C6—H6A	0.9600	Fe1—Cl2'	2.211 (7)
C6—H6B	0.9600	Fe1—Cl3'	2.256 (5)
C6—H6C	0.9600	Fe1—Cl4	2.1821 (17)
C7—C8	1.506 (7)		
C2—C1—N1	115.3 (4)	C13—C8—C7	121.3 (5)
C2—C1—H1A	108.4	C8—C9—C10	121.2 (5)
N1—C1—H1A	108.4	C8—C9—H9	119.4
C2—C1—H1B	108.4	C10—C9—H9	119.4
N1—C1—H1B	108.4	C11—C10—C9	118.9 (6)
H1A—C1—H1B	107.5	C11—C10—H10	120.5
C1—C2—H2A	109.5	C9—C10—H10	120.5
C1—C2—H2B	109.5	C12—C11—C10	120.7 (6)
H2A—C2—H2B	109.5	C12—C11—H11	119.7
C1—C2—H2C	109.5	C10—C11—H11	119.7
H2A—C2—H2C	109.5	C11—C12—C13	120.4 (6)
H2B—C2—H2C	109.5	C11—C12—H12	119.8
C4—C3—N1	116.1 (4)	C13—C12—H12	119.8
C4—C3—H3A	108.3	C12—C13—C8	121.5 (6)
N1—C3—H3A	108.3	C12—C13—H13	119.3
C4—C3—H3B	108.3	C8—C13—H13	119.3
N1—C3—H3B	108.3	Cl3—Fe1—Cl1'	90.7 (3)
H3A—C3—H3B	107.4	Cl3—Fe1—Cl2	111.6 (4)
C3—C4—H4A	109.5	Cl1'—Fe1—Cl2	112.2 (5)
C3—C4—H4B	109.5	Cl3—Fe1—Cl4	124.7 (4)
H4A—C4—H4B	109.5	Cl1'—Fe1—Cl4	108.99 (14)
C3—C4—H4C	109.5	Cl2—Fe1—Cl4	107.5 (4)
H4A—C4—H4C	109.5	Cl3—Fe1—Cl2'	99.9 (3)
H4B—C4—H4C	109.5	Cl1'—Fe1—Cl2'	122.4 (3)
N1—C5—C6	114.9 (4)	Cl2—Fe1—Cl2'	14.5 (4)
N1—C5—H5A	108.5	Cl4—Fe1—Cl2'	110.2 (2)
C6—C5—H5A	108.5	Cl3—Fe1—Cl1	109.1 (4)
N1—C5—H5B	108.5	Cl1'—Fe1—Cl1	21.8 (3)
C6—C5—H5B	108.5	Cl2—Fe1—Cl1	93.9 (7)
H5A—C5—H5B	107.5	Cl4—Fe1—Cl1	105.4 (2)
C5—C6—H6A	109.5	Cl2'—Fe1—Cl1	106.3 (5)
C5—C6—H6B	109.5	Cl3—Fe1—Cl3'	23.7 (2)
H6A—C6—H6B	109.5	Cl1'—Fe1—Cl3'	106.90 (18)
C5—C6—H6C	109.5	Cl2—Fe1—Cl3'	118.3 (4)
H6A—C6—H6C	109.5	Cl4—Fe1—Cl3'	102.3 (2)
H6B—C6—H6C	109.5	Cl2'—Fe1—Cl3'	104.12 (19)
C8—C7—N1	115.5 (4)	Cl1—Fe1—Cl3'	127.9 (4)

supplementary materials

C8—C7—H7A	108.4	C5—N1—C3	106.3 (3)
N1—C7—H7A	108.4	C5—N1—C7	111.0 (4)
C8—C7—H7B	108.4	C3—N1—C7	110.9 (3)
N1—C7—H7B	108.4	C5—N1—C1	111.2 (4)
H7A—C7—H7B	107.5	C3—N1—C1	110.3 (4)
C9—C8—C13	117.2 (5)	C7—N1—C1	107.1 (3)
C9—C8—C7	121.4 (5)		
