

catena-Poly[[bis(nitrato- κ O)cadmium]-bis[μ -1,3-bis[(1H-1,2,4-triazol-1-yl)-methyl]benzene- $\kappa^2N^4:N^4'$]]

Hong-Kun Zhang,^{a*} Xin Wang,^a Shuai Wang^a and
Xiao-Dan Wang^b

^aDepartment of Food and Environmental Engineering, Heilongjiang East University, Harbin 150086, People's Republic of China, and ^bCollege of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China
Correspondence e-mail: zhanghongkun2000@163.com


Received 16 May 2012; accepted 21 May 2012

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; disorder in main residue; R factor = 0.030; wR factor = 0.080; data-to-parameter ratio = 14.9.

In the title compound, $[\text{Cd}(\text{NO}_3)_2(\text{C}_{12}\text{H}_{12}\text{N}_6)_2]_n$, the Cd^{II} cation is located on an inversion center and is six-coordinated by four N atoms from four 1,3-bis[(1H-1,2,4-triazol-1-yl)methyl]benzene (*L*) ligands and two O atoms from two nitrate anions in a slightly distorted octahedral geometry. The ligands link different Cd^{II} ions into a ribbon-like structure along [001]. Two O atoms of the nitrate anion are disordered over two sets of sites with site occupancies of 0.575 (8) and 0.425 (8).

Related literature

For related structures, see: Meng *et al.* (2004). For the synthesis of the ligand, see: Du *et al.* (2008).



Experimental

Crystal data

$[\text{Cd}(\text{NO}_3)_2(\text{C}_{12}\text{H}_{12}\text{N}_6)_2]$	$\gamma = 109.71(3)^\circ$
$M_r = 716.97$	$V = 736.2(2)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 1$
$a = 8.0412(16)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.7303(17)\text{ \AA}$	$\mu = 0.81\text{ mm}^{-1}$
$c = 11.598(2)\text{ \AA}$	$T = 293\text{ K}$
$\alpha = 105.12(3)^\circ$	$0.41 \times 0.28 \times 0.15\text{ mm}$
$\beta = 90.20(3)^\circ$	

Data collection

Rigaku R-AXIS RAPID diffractometer	7285 measured reflections
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	3346 independent reflections
$T_{\min} = 0.734$, $T_{\max} = 0.886$	3216 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$	3 restraints
$wR(F^2) = 0.080$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\max} = 0.77\text{ e \AA}^{-3}$
3346 reflections	$\Delta\rho_{\min} = -0.68\text{ e \AA}^{-3}$
224 parameters	

Table 1
Selected bond lengths (Å).

Cd1—N1	2.330 (2)	Cd1—O1	2.479 (5)
Cd1—N4 ⁱ	2.3278 (19)		

Symmetry code: (i) $-x + 1, -y + 1, -z$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Heilongjiang East University and Heilongjiang University are thanked for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VN2040).

References

- Du, J.-L., Hu, T.-L., Zhang, S.-M., Zheng, T.-F. & Bu, X.-H. (2008). *CrystEngComm*, **10**, 1866–1874.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Meng, X.-L., Song, Y.-L., Hou, H.-W., Han, H.-Y., Xiao, B., Fan, Y.-T. & Zhu, Y. (2004). *Inorg. Chem.* **43**, 3528–3536.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2012). E68, m856 [doi:10.1107/S1600536812023288]

catena-Poly[[bis(nitrato- κO)cadmium]bis[μ -1,3-bis[(1*H*-1,2,4-triazol-1-yl)methyl]benzene- $\kappa^2 N^4:N^4'$]]

Hong-Kun Zhang, Xin Wang, Shuai Wang and Xiao-Dan Wang

Comment

In recent years, much attention has been paid to the use of nitrogen-containing ligands for constructing supramolecular coordination compounds. The reason is that the supramolecular coordination assemblies have not only a variety of architectures but also have potential applications as functional materials. Recently, a series of supramolecular complexes based on the 1,4-bis(1*H*-1,2,4-triazol-1-yl-methyl)-benzene ligand were reported (Meng *et al.*, 2004). In this paper, we report the new title compound, synthesized by the reaction of 1,3-bis((1*H*-1,2,4-triazol-1-yl)methyl)benzene and cadmium dinitrate in an aqueous solution.

In the title compound, $[Cd(NO_3)_2(C_{12}H_{12}N_6)_2]_n$, the eight-coordinated Cd^{II} ion is located on an inversion center and is in an octahedral environment defined by four N atoms from ligands forming the equatorial plane (distances Cd—N1 = 2.330 (2) Å and Cd—N4 = 2.328 (2) Å) and two O atoms from two nitrate anions lying on the polar axis with a Cd—O1 distance of 2.479 (5) Å (Figure 1, Table 1). An infinite ribbon-like structure running along [001] is built up by the *cis*-ligands linking these Cd^{II} ions (Figure 2).

Experimental


The ligand L was synthesized following the reference method (Du *et al.*, 2008). Synthesis of the title compound: L (0.120 g, 0.5 mmol) and Cd(NO₃)₂ (0.152 g, 0.5 mmol) were dissolved in a mixed solution of 3 mL ethanol and 3 mL water. After stirring, the suspension was sealed in a 18 mL Teflon-lined autoclave and heated at 140 °C for 5 days. After slow cooling to room temperature, colorless block crystals were filtered and washed with distilled water (47% yield based on Cd).

Refinement


H atoms bound to C atoms were placed in calculated positions and treated as riding on their parent atoms, with C—H = 0.93 / 0.97 Å (aromatic / methylene) and with $U_{iso}(H) = 1.2U_{eq}(C)$. Two O atoms of the nitrate anion were disordered over two positions with site occupancy of 0.58 for O1 and O2 atoms, and 0.42 for O1' and O2' atoms, respectively. The *SHELXL* 'DFIX' instruction was used to restrain the N—O bond distances of the disordered nitrate anion to close to 1.22 Å.

Computing details

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO* (Rigaku, 1998); data reduction: *CrystalClear* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).

**Figure 1**

The molecular structure of the title compound, showing displacement ellipsoids at the 50% probability level for non-H atoms. Only the main component of the disordered nitrate anion is shown. Symmetry codes: (i) $1-x, 1-y, 1-z$; (ii) $1-x, 1-y, -z$; (iii) $x, y, 1+z$.

**Figure 2**

A portion of the ribbon-like structure in the title compound. H atoms and disordered O atoms have been omitted for clarity.

catena-Poly[[bis(nitrato- κ O)cadmium]bis[μ -1,3-bis[(1H-1,2,4-triazol-1-yl)methyl]benzene- κ^2 N⁴:N^{4'}]]

Crystal data

$[\text{Cd}(\text{NO}_3)_2(\text{C}_{12}\text{H}_{12}\text{N}_6)_2]$

$M_r = 716.97$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 8.0412 (16)$ Å

$b = 8.7303 (17)$ Å

$c = 11.598 (2)$ Å
 $\alpha = 105.12 (3)^\circ$
 $\beta = 90.20 (3)^\circ$
 $\gamma = 109.71 (3)^\circ$
 $V = 736.2 (2)$ Å³
 $Z = 1$
 $F(000) = 362$
 $D_x = 1.617 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 7192 reflections
 $\theta = 3.0\text{--}27.5^\circ$
 $\mu = 0.81 \text{ mm}^{-1}$
 $T = 293$ K
Block, colourless
 $0.41 \times 0.28 \times 0.15$ mm

Data collection

Rigaku R-AXIS RAPID
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scan
Absorption correction: multi-scan
(ABSCOR; Higashi, 1995)
 $T_{\min} = 0.734$, $T_{\max} = 0.886$

7285 measured reflections
3346 independent reflections
3216 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.0^\circ$
 $h = -10 \rightarrow 10$
 $k = -11 \rightarrow 10$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.080$
 $S = 1.03$
3346 reflections
224 parameters
3 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0539P)^2 + 0.1306P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.77 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.68 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.5033 (3)	0.3521 (3)	0.2034 (2)	0.0506 (5)	
H1	0.4228	0.4026	0.1901	0.061*	
C2	0.6741 (3)	0.2753 (3)	0.29299 (19)	0.0456 (5)	
H2	0.7392	0.2575	0.3515	0.055*	
C3	0.7674 (4)	0.1037 (3)	0.1121 (2)	0.0503 (5)	
H3A	0.6804	-0.0067	0.0726	0.060*	
H3B	0.8445	0.0892	0.1697	0.060*	
C4	0.8758 (3)	0.1744 (3)	0.0203 (2)	0.0493 (5)	
C5	1.0410 (4)	0.3011 (5)	0.0530 (3)	0.0798 (10)	

H5	1.0883	0.3424	0.1329	0.096*	
C6	1.1364 (5)	0.3666 (6)	-0.0340 (4)	0.0995 (13)	
H6	1.2456	0.4541	-0.0117	0.119*	
C7	1.0688 (4)	0.3017 (5)	-0.1530 (3)	0.0782 (9)	
H7	1.1344	0.3438	-0.2109	0.094*	
C8	0.9050 (4)	0.1752 (3)	-0.1870 (2)	0.0524 (5)	
C9	0.8091 (3)	0.1123 (3)	-0.0998 (2)	0.0461 (5)	
H9	0.6983	0.0272	-0.1223	0.055*	
C10	0.8280 (4)	0.1018 (4)	-0.3175 (2)	0.0618 (7)	
H10A	0.9232	0.1280	-0.3682	0.074*	
H10B	0.7764	-0.0205	-0.3350	0.074*	
C11	0.7118 (3)	0.2961 (3)	-0.3915 (2)	0.0474 (5)	
H11	0.8197	0.3670	-0.4069	0.057*	
C12	0.4422 (4)	0.1801 (4)	-0.3748 (3)	0.0628 (6)	
H12	0.3203	0.1568	-0.3780	0.075*	
Cd1	0.5000	0.5000	0.5000	0.04272 (9)	
N1	0.5669 (3)	0.3636 (3)	0.31476 (17)	0.0487 (4)	
N2	0.5646 (3)	0.2640 (3)	0.11613 (18)	0.0532 (5)	
N3	0.6753 (2)	0.2162 (2)	0.17601 (16)	0.0421 (4)	
N4	0.5558 (3)	0.3100 (2)	-0.41081 (17)	0.0476 (4)	
N5	0.5189 (4)	0.0906 (3)	-0.3351 (3)	0.0695 (6)	
N6	0.6919 (3)	0.1667 (2)	-0.34697 (17)	0.0494 (4)	
N7	0.9115 (2)	0.7301 (3)	0.5706 (2)	0.0529 (5)	
O1	0.8078 (6)	0.6800 (8)	0.4850 (4)	0.093 (2)	0.575 (8)
O2	0.8601 (8)	0.6969 (6)	0.6706 (4)	0.0892 (18)	0.575 (8)
O1'	0.9178 (14)	0.5978 (8)	0.4970 (11)	0.166 (7)	0.425 (8)
O2'	0.7681 (6)	0.6944 (7)	0.6029 (8)	0.087 (3)	0.425 (8)
O3	1.0506 (2)	0.8468 (2)	0.58725 (18)	0.0585 (4)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0476 (12)	0.0654 (14)	0.0474 (11)	0.0266 (10)	0.0043 (9)	0.0209 (10)
C2	0.0513 (12)	0.0516 (11)	0.0396 (10)	0.0235 (9)	0.0035 (8)	0.0147 (9)
C3	0.0635 (14)	0.0542 (12)	0.0446 (11)	0.0304 (11)	0.0151 (10)	0.0196 (10)
C4	0.0494 (12)	0.0616 (13)	0.0489 (11)	0.0288 (10)	0.0145 (9)	0.0226 (10)
C5	0.0503 (15)	0.115 (3)	0.0644 (16)	0.0128 (16)	0.0023 (12)	0.0295 (17)
C6	0.0533 (18)	0.134 (3)	0.095 (3)	-0.0005 (19)	0.0142 (17)	0.049 (2)
C7	0.0623 (18)	0.110 (2)	0.079 (2)	0.0308 (17)	0.0312 (15)	0.0531 (19)
C8	0.0636 (14)	0.0671 (14)	0.0520 (12)	0.0444 (12)	0.0224 (10)	0.0301 (11)
C9	0.0516 (12)	0.0500 (11)	0.0467 (11)	0.0259 (9)	0.0130 (9)	0.0194 (9)
C10	0.093 (2)	0.0764 (16)	0.0493 (12)	0.0617 (16)	0.0242 (13)	0.0296 (12)
C11	0.0579 (13)	0.0468 (11)	0.0476 (11)	0.0249 (9)	0.0122 (9)	0.0212 (9)
C12	0.0568 (15)	0.0597 (14)	0.0770 (17)	0.0169 (11)	0.0113 (13)	0.0320 (13)
Cd1	0.05123 (15)	0.04842 (14)	0.03877 (13)	0.02770 (10)	0.00625 (8)	0.01561 (9)
N1	0.0545 (11)	0.0579 (11)	0.0424 (9)	0.0288 (9)	0.0050 (8)	0.0164 (8)
N2	0.0510 (11)	0.0723 (13)	0.0429 (9)	0.0262 (9)	0.0031 (8)	0.0209 (9)
N3	0.0433 (9)	0.0485 (9)	0.0392 (8)	0.0179 (7)	0.0077 (7)	0.0173 (7)
N4	0.0564 (11)	0.0451 (9)	0.0490 (10)	0.0229 (8)	0.0085 (8)	0.0191 (8)
N5	0.0727 (15)	0.0610 (13)	0.0887 (17)	0.0231 (11)	0.0196 (13)	0.0442 (13)

N6	0.0685 (13)	0.0499 (10)	0.0450 (9)	0.0330 (9)	0.0155 (9)	0.0219 (8)
N7	0.0389 (10)	0.0562 (11)	0.0699 (13)	0.0168 (8)	0.0083 (9)	0.0280 (10)
O1	0.057 (2)	0.124 (4)	0.078 (3)	0.010 (2)	-0.025 (2)	0.024 (3)
O2	0.077 (3)	0.105 (3)	0.076 (3)	0.004 (3)	0.016 (2)	0.047 (2)
O1'	0.127 (9)	0.108 (6)	0.159 (8)	-0.046 (6)	0.067 (7)	-0.023 (6)
O2'	0.043 (3)	0.076 (3)	0.115 (6)	0.009 (2)	0.028 (3)	-0.003 (3)
O3	0.0391 (8)	0.0582 (9)	0.0754 (12)	0.0123 (7)	0.0042 (8)	0.0201 (9)

Geometric parameters (\AA , $^\circ$)

C1—N2	1.309 (3)	C10—H10B	0.9700
C1—N1	1.353 (3)	C11—N6	1.323 (3)
C1—H1	0.9300	C11—N4	1.324 (3)
C2—N3	1.321 (3)	C11—H11	0.9300
C2—N1	1.323 (3)	C12—N5	1.309 (4)
C2—H2	0.9300	C12—N4	1.358 (3)
C3—N3	1.475 (3)	C12—H12	0.9300
C3—C4	1.504 (3)	Cd1—O2 ⁱ	2.326 (6)
C3—H3A	0.9700	Cd1—O2'	2.326 (6)
C3—H3B	0.9700	Cd1—N1	2.330 (2)
C4—C5	1.385 (4)	Cd1—N4 ⁱⁱ	2.3278 (19)
C4—C9	1.388 (3)	Cd1—N4 ⁱⁱⁱ	2.3278 (19)
C5—C6	1.394 (5)	Cd1—N1 ⁱ	2.330 (2)
C5—H5	0.9300	Cd1—O1	2.479 (5)
C6—C7	1.378 (5)	Cd1—O1 ⁱ	2.479 (5)
C6—H6	0.9300	N2—N3	1.359 (3)
C7—C8	1.378 (5)	N4—Cd1 ^{iv}	2.3278 (19)
C7—H7	0.9300	N5—N6	1.349 (3)
C8—C9	1.388 (3)	N7—O2'	1.179 (4)
C8—C10	1.518 (4)	N7—O1	1.180 (3)
C9—H9	0.9300	N7—O3	1.206 (3)
C10—N6	1.468 (3)	N7—O1'	1.260 (5)
C10—H10A	0.9700	N7—O2	1.304 (5)
N2—C1—N1	114.4 (2)	O2 ⁱ —Cd1—N1	73.58 (14)
N2—C1—H1	122.8	O2'—Cd1—N1	106.42 (14)
N1—C1—H1	122.8	N4 ⁱⁱ —Cd1—N1	88.62 (7)
N3—C2—N1	110.0 (2)	N4 ⁱⁱⁱ —Cd1—N1	91.38 (7)
N3—C2—H2	125.0	O2 ⁱ —Cd1—N1 ⁱ	106.42 (14)
N1—C2—H2	125.0	O2'—Cd1—N1 ⁱ	73.58 (14)
N3—C3—C4	111.58 (19)	N4 ⁱⁱ —Cd1—N1 ⁱ	91.38 (7)
N3—C3—H3A	109.3	N4 ⁱⁱⁱ —Cd1—N1 ⁱ	88.62 (7)
C4—C3—H3A	109.3	N1—Cd1—N1 ⁱ	180.0
N3—C3—H3B	109.3	O2 ⁱ —Cd1—O1	146.62 (19)
C4—C3—H3B	109.3	O2'—Cd1—O1	33.38 (19)
H3A—C3—H3B	108.0	N4 ⁱⁱ —Cd1—O1	79.67 (14)
C5—C4—C9	119.1 (2)	N4 ⁱⁱⁱ —Cd1—O1	100.33 (14)
C5—C4—C3	121.4 (2)	N1—Cd1—O1	76.10 (15)
C9—C4—C3	119.5 (2)	N1 ⁱ —Cd1—O1	103.90 (15)
C4—C5—C6	120.0 (3)	O2 ⁱ —Cd1—O1 ⁱ	33.38 (19)

C4—C5—H5	120.0	O2'—Cd1—O1 ⁱ	146.62 (19)
C6—C5—H5	120.0	N4 ⁱⁱ —Cd1—O1 ⁱ	100.33 (14)
C7—C6—C5	120.0 (3)	N4 ⁱⁱⁱ —Cd1—O1 ⁱ	79.67 (14)
C7—C6—H6	120.0	N1—Cd1—O1 ⁱ	103.90 (15)
C5—C6—H6	120.0	N1 ⁱ —Cd1—O1 ⁱ	76.10 (15)
C8—C7—C6	120.7 (3)	O1—Cd1—O1 ⁱ	180.0
C8—C7—H7	119.7	C2—N1—C1	103.0 (2)
C6—C7—H7	119.7	C2—N1—Cd1	128.22 (15)
C7—C8—C9	119.1 (3)	C1—N1—Cd1	128.74 (16)
C7—C8—C10	121.6 (2)	C1—N2—N3	102.59 (18)
C9—C8—C10	119.3 (3)	C2—N3—N2	109.94 (19)
C4—C9—C8	121.1 (2)	C2—N3—C3	128.28 (19)
C4—C9—H9	119.4	N2—N3—C3	121.62 (18)
C8—C9—H9	119.4	C11—N4—C12	102.7 (2)
N6—C10—C8	113.12 (19)	C11—N4—Cd1 ^{iv}	126.68 (16)
N6—C10—H10A	109.0	C12—N4—Cd1 ^{iv}	130.47 (18)
C8—C10—H10A	109.0	C12—N5—N6	102.9 (2)
N6—C10—H10B	109.0	C11—N6—N5	110.0 (2)
C8—C10—H10B	109.0	C11—N6—C10	128.3 (2)
H10A—C10—H10B	107.8	N5—N6—C10	121.7 (2)
N6—C11—N4	110.1 (2)	O2'—N7—O1	72.1 (5)
N6—C11—H11	124.9	O2'—N7—O3	141.1 (4)
N4—C11—H11	124.9	O1—N7—O3	125.6 (3)
N5—C12—N4	114.3 (3)	O2'—N7—O1'	106.5 (5)
N5—C12—H12	122.9	O1—N7—O1'	66.5 (7)
N4—C12—H12	122.9	O3—N7—O1'	112.4 (4)
O2 ⁱ —Cd1—O2'	180.000 (1)	O2'—N7—O2	50.7 (4)
O2 ⁱ —Cd1—N4 ⁱⁱ	85.9 (2)	O1—N7—O2	119.7 (4)
O2'—Cd1—N4 ⁱⁱ	94.1 (2)	O3—N7—O2	111.6 (3)
O2 ⁱ —Cd1—N4 ⁱⁱⁱ	94.1 (2)	O1'—N7—O2	109.5 (6)
O2'—Cd1—N4 ⁱⁱⁱ	85.9 (2)	N7—O1—Cd1	118.1 (3)
N4 ⁱⁱ —Cd1—N4 ⁱⁱⁱ	180.0	N7—O2—Cd1	129.6 (5)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, -y+1, -z$; (iii) $x, y, z+1$; (iv) $x, y, z-1$.