

## Ethyl (3*E*)-3-[2-(4-bromophenylsulfonyl)-hydrazin-1-ylidene]butanoate

Shahzad Murtaza,<sup>a</sup> Naghma Kausar,<sup>a</sup> Aadil Abbas,<sup>a</sup>  
M. Nawaz Tahir<sup>b\*</sup> and Muhammad Zulfiqar<sup>b</sup>

<sup>a</sup>University of Gujrat, Department of Chemistry, Hafiz Hayat Campus, Gujrat, Pakistan, and <sup>b</sup>University of Sargodha, Department of Physics, Sargodha, Pakistan  
Correspondence e-mail: dmntahir\_uos@yahoo.com


Received 29 April 2012; accepted 29 April 2012

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$ ;  $R$  factor = 0.049;  $wR$  factor = 0.135; data-to-parameter ratio = 17.7.

The asymmetric unit of title compound,  $\text{C}_{12}\text{H}_{15}\text{BrN}_2\text{O}_4\text{S}$ , contains two molecules (*A* and *B*), with slightly different conformations: the bromophenyl rings and the  $\text{SO}_2$  planes of the sulfonyl groups are oriented at dihedral angles of 50.2 (2) (molecule *A*) and 58.24 (7)° (molecule *B*), and the ethyl acetate groups make dihedral angles of 63.99 (19)° (*A*) and 65.35 (16)° (*B*) with their bromophenyl groups. In the crystal, both molecules exist as inversion dimers linked by pairs of  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds, which generate  $R_2^2(14)$  loops. The dimers are linked by  $\text{C}-\text{H}\cdots\text{O}$  interactions.

### Related literature

For a related structure, see: Uramoto *et al.* (1971). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



### Experimental

#### Crystal data

$\text{C}_{12}\text{H}_{15}\text{BrN}_2\text{O}_4\text{S}$   
 $M_r = 363.23$   
Triclinic,  $P\bar{1}$

$a = 11.0550 (3)\text{ \AA}$   
 $b = 11.9763 (4)\text{ \AA}$   
 $c = 13.1146 (3)\text{ \AA}$

$\alpha = 77.457 (1)^\circ$   
 $\beta = 72.163 (2)^\circ$   
 $\gamma = 73.981 (1)^\circ$   
 $V = 1571.92 (8)\text{ \AA}^3$   
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 2.76\text{ mm}^{-1}$   
 $T = 296\text{ K}$   
 $0.30 \times 0.15 \times 0.14\text{ mm}$

#### Data collection

Bruker Kappa APEXII CCD diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)  
 $T_{\min} = 0.635$ ,  $T_{\max} = 0.650$

22120 measured reflections  
6096 independent reflections  
4293 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.135$   
 $S = 1.05$   
6096 reflections

344 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 1.27\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -1.11\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ , °).

$D-\text{H}\cdots A$	$D-\text{H}$	$H\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 $\cdots$ O3 <sup>i</sup>	0.86	2.46	2.909 (5)	114
N3—H3A $\cdots$ O7 <sup>ii</sup>	0.86	2.36	2.8855 (13)	120
C3—H3 $\cdots$ O4 <sup>iii</sup>	0.93	2.53	3.450 (6)	172

Symmetry codes: (i)  $-x + 1, -y, -z + 1$ ; (ii)  $-x + 2, -y + 1, -z$ ; (iii)  $-x + 2, -y, -z + 1$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of a diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan. The authors also acknowledge the technical support provided by Syed Muhammad Hussain Rizvi of Bana International, Karachi, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6771).

### References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Uramoto, M., Otake, N., Ogawa, Y., Yonehara, H., Marumo, F. & Saito, Y. (1971). *Acta Cryst. B* **27**, 236–241.

# supplementary materials

*Acta Cryst.* (2012). E68, o1616 [doi:10.1107/S1600536812019265]

## Ethyl (3*E*)-3-[2-(4-bromophenylsulfonyl)hydrazin-1-ylidene]butanoate

**Shahzad Murtaza, Naghma Kausar, Aadil Abbas, M. Nawaz Tahir and Muhammad Zulfiqar**

### Comment

The title compound, (I) (Fig. 1), has been synthesized for the study of enzyme inhibition and other biological activities owing to the concept that sulfonyl hydrazide moiety is an important part of different drugs.

The crystal structure of bundlin A *p*-bromophenylsulfonylhydrazone (Uramoto *et al.*, 1971) has been published which contains the common moiety 4-bromo-*N'*-(propan-2-ylidene)benzenesulfonohydrazide which is related to the title compound (I, Fig. 1).

In (I), two molecules in the asymmetric unit are present, which differ from each other geometrically. In one molecule, the moieties of bromophenyl A (C1–C6/BR1), ethylidenehydrazine B (N1/N2/C7/C8) and ethyl acetate C (C9/C10/O3/O4/C11/C12) are almost planar with r. m. s. deviation of 0.0255, 0.0078 and 0.0398 Å, respectively. The dihedral angle between A/B, A/C and B/C is 86.14 (13)°, 54.81 (14)° and 65.35 (16)°, respectively. The sulfonyl group D (O1/S1/O2) is of course planar and makes a dihydral angle of 50.23 (21)° with its parent bromophenyl moiety. In second molecule, the similar moieties E (C13–C18/BR2), F (N3/N4/C19/C20) and G (C21/C22/O7/O8/C23/C24) are planar with r. m. s. deviation of 0.0059, 0.0016 and 0.0522 Å, respectively. The dihedral angle between E/F, E/G and F/G is 88.25 (10)°, 58.35 (14)° and 63.99 (19)°, respectively. The sulfonyl group H (O5/S2/O6) makes a dihydral angle of 58.24 (7)° with its parent bromophenyl E moiety. Both molecules are dimerized with themselves due to classical H-bondings of N—H···O type with  $R_2^2(14)$  (Table 1, Fig. 2) ring motifs (Bernstein *et al.*, 1995). The dimers of molecules containing bromine atom BR1 are interlinked due to C—H···O type of H-bondings, where CH is of phenyl ring and O-atom is of ester group.

### Experimental


A mixture of 4-bromobenzenesulphonohydrazide (0.251 g, 1 mmol) and ethyl acetoacetate (0.130 g, 1 mmol) in ethanol (20 ml) was refluxed for 4 h and then cooled to room temperature. After cooling, the solvent was removed under reduced pressure and the colorless solid residue was obtained. The resulting product was re-crystallized in ethanol and obtained colorless prisms of (I). m.p. 388 K.

### Refinement

The H-atoms were positioned geometrically (N—H = 0.86, C—H = 0.93–0.97 Å) and were included in the refinement in the riding model approximation, with  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$ , where  $x = 1.5$  for  $\text{CH}_3$  and  $x = 1.2$  for other H-atoms.


### Computing details

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).



**Figure 1**

View of the title compound with displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

The partial packing, which shows that molecules are dimers and then polymeric chains are formed. The H-atoms not involved in H-bondings are omitted for clarity.

#### Ethyl (3E)-3-[2-(4-bromophenylsulfonyl)hydrazin-1-ylidene]butanoate

##### Crystal data

$C_{12}H_{15}BrN_2O_4S$   
 $M_r = 363.23$   
Triclinic,  $P\bar{1}$   
Hall symbol: -P 1  
 $a = 11.0550 (3)$  Å  
 $b = 11.9763 (4)$  Å  
 $c = 13.1146 (3)$  Å  
 $\alpha = 77.457 (1)^\circ$   
 $\beta = 72.163 (2)^\circ$   
 $\gamma = 73.981 (1)^\circ$   
 $V = 1571.92 (8)$  Å<sup>3</sup>

$Z = 4$   
 $F(000) = 736$   
 $D_x = 1.535 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 4293 reflections  
 $\theta = 2.0\text{--}26.0^\circ$   
 $\mu = 2.76 \text{ mm}^{-1}$   
 $T = 296 \text{ K}$   
Prism, colourless  
 $0.30 \times 0.15 \times 0.14$  mm

*Data collection*

Bruker Kappa APEXII CCD diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 Detector resolution: 7.80 pixels mm<sup>-1</sup>  
 $\omega$  scans  
 Absorption correction: multi-scan (SADABS; Bruker, 2005)  
 $T_{\min} = 0.635$ ,  $T_{\max} = 0.650$

22120 measured reflections  
 6096 independent reflections  
 4293 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$   
 $\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 2.0^\circ$   
 $h = -13 \rightarrow 13$   
 $k = -14 \rightarrow 14$   
 $l = -16 \rightarrow 16$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.135$   
 $S = 1.05$   
 6096 reflections  
 344 parameters  
 0 restraints  
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
 Hydrogen site location: inferred from neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0612P)^2 + 1.5738P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 1.27 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -1.11 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.11769 (5)	-0.38812 (5)	0.43815 (4)	0.0788 (2)
S1	0.78563 (9)	0.03587 (8)	0.17990 (7)	0.0411 (3)
O1	0.8641 (3)	0.1199 (2)	0.1313 (2)	0.0557 (10)
O2	0.7263 (3)	-0.0035 (3)	0.1162 (2)	0.0558 (10)
O3	0.5216 (3)	0.0584 (3)	0.6384 (2)	0.0629 (11)
O4	0.6201 (3)	0.1478 (3)	0.7100 (2)	0.0532 (10)
N1	0.6630 (3)	0.0928 (3)	0.2748 (2)	0.0397 (10)
N2	0.7025 (3)	0.1166 (3)	0.3597 (2)	0.0384 (10)
C1	0.8795 (3)	-0.0857 (3)	0.2460 (3)	0.0381 (11)
C2	1.0045 (4)	-0.0849 (4)	0.2421 (4)	0.0544 (14)
C3	1.0752 (4)	-0.1770 (4)	0.2979 (4)	0.0597 (16)
C4	1.0203 (4)	-0.2686 (3)	0.3560 (3)	0.0461 (12)
C5	0.8954 (4)	-0.2710 (4)	0.3593 (3)	0.0482 (12)
C6	0.8250 (4)	-0.1790 (3)	0.3030 (3)	0.0453 (12)
C7	0.6106 (3)	0.1691 (3)	0.4305 (3)	0.0396 (11)
C8	0.4704 (4)	0.2101 (4)	0.4284 (3)	0.0586 (16)
C9	0.6502 (4)	0.1910 (3)	0.5228 (3)	0.0424 (12)

C10	0.5890 (3)	0.1252 (3)	0.6290 (3)	0.0419 (12)
C11	0.5598 (6)	0.0940 (7)	0.8188 (4)	0.099 (3)
C12	0.6102 (3)	0.1162 (2)	0.89744 (13)	0.109 (3)
Br2	0.35796 (4)	0.36766 (4)	0.06489 (4)	0.0807 (2)
S2	0.68585 (4)	0.53305 (5)	0.30636 (4)	0.0406 (3)
O5	0.60925 (4)	0.63906 (5)	0.34696 (4)	0.0562 (10)
O6	0.73411 (5)	0.43392 (7)	0.37752 (5)	0.0600 (10)
O7	0.99883 (6)	0.64595 (9)	-0.14623 (7)	0.0629 (11)
O8	0.89569 (5)	0.82270 (6)	-0.20911 (5)	0.0571 (10)
N3	0.81726 (5)	0.56299 (7)	0.21809 (5)	0.0401 (10)
N4	0.7935 (3)	0.6522 (3)	0.1321 (2)	0.0384 (9)
C13	0.5960 (3)	0.4894 (3)	0.2375 (3)	0.0378 (11)
C14	0.4690 (4)	0.5485 (4)	0.2418 (4)	0.0541 (16)
C15	0.3982 (4)	0.5113 (4)	0.1912 (4)	0.0646 (16)
C16	0.4555 (4)	0.4169 (4)	0.1368 (3)	0.0513 (14)
C17	0.5823 (4)	0.3571 (3)	0.1316 (3)	0.0498 (14)
C18	0.6528 (4)	0.3938 (3)	0.1828 (3)	0.0448 (12)
C19	0.8945 (4)	0.6858 (3)	0.0688 (3)	0.0411 (12)
C20	1.0308 (4)	0.6400 (4)	0.0792 (4)	0.0603 (16)
C21	0.8693 (4)	0.7805 (3)	-0.0225 (3)	0.0442 (12)
C22	0.9295 (4)	0.7404 (3)	-0.1316 (3)	0.0428 (12)
C23	0.9492 (6)	0.7938 (5)	-0.3193 (4)	0.088 (2)
C24	0.8877 (10)	0.8834 (7)	-0.3902 (5)	0.151 (4)
H1	0.58287	0.10678	0.27334	0.0476*
H2	1.04121	-0.02258	0.20198	0.0649*
H3	1.15997	-0.17705	0.29617	0.0714*
H5	0.85930	-0.33381	0.39886	0.0579*
H6	0.74091	-0.17969	0.30335	0.0542*
H8A	0.46525	0.24267	0.35579	0.0878*
H8B	0.42710	0.26895	0.47500	0.0878*
H8C	0.42897	0.14484	0.45306	0.0878*
H9A	0.62468	0.27440	0.52727	0.0511*
H9B	0.74440	0.16716	0.50890	0.0511*
H11A	0.46620	0.12524	0.83593	0.1181*
H11B	0.57574	0.00993	0.82031	0.1181*
H12A	0.70237	0.08301	0.88193	0.1631*
H12B	0.56803	0.08127	0.96785	0.1631*
H12C	0.59457	0.19941	0.89594	0.1631*
H3A	0.89452	0.52737	0.22352	0.0481*
H14	0.43149	0.61308	0.27857	0.0650*
H15	0.31208	0.54993	0.19406	0.0775*
H17	0.61965	0.29300	0.09419	0.0597*
H18	0.73855	0.35439	0.18051	0.0534*
H20A	1.03026	0.62775	0.15418	0.0903*
H20B	1.08237	0.69588	0.03942	0.0903*
H20C	1.06744	0.56699	0.05082	0.0903*
H21A	0.90367	0.84604	-0.02061	0.0530*
H21B	0.77582	0.80815	-0.01255	0.0530*
H23A	1.04258	0.78818	-0.34174	0.1054*

H23B	0.93377	0.71873	-0.32190	0.1054*
H24A	0.79493	0.89168	-0.36510	0.2268*
H24B	0.91809	0.86258	-0.46188	0.2268*
H24C	0.90870	0.95638	-0.39131	0.2268*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0633 (3)	0.0840 (4)	0.0816 (4)	-0.0112 (3)	-0.0321 (3)	0.0156 (3)
S1	0.0424 (5)	0.0425 (5)	0.0351 (5)	-0.0108 (4)	-0.0027 (4)	-0.0082 (4)
O1	0.0520 (17)	0.0497 (17)	0.0538 (17)	-0.0190 (14)	0.0037 (13)	0.0009 (13)
O2	0.0610 (18)	0.0643 (19)	0.0442 (16)	-0.0064 (14)	-0.0166 (14)	-0.0179 (14)
O3	0.076 (2)	0.073 (2)	0.0516 (17)	-0.0494 (18)	-0.0108 (15)	-0.0003 (15)
O4	0.0545 (16)	0.075 (2)	0.0368 (14)	-0.0327 (15)	-0.0092 (12)	-0.0028 (13)
N1	0.0329 (16)	0.0461 (18)	0.0392 (17)	-0.0084 (14)	-0.0044 (13)	-0.0123 (14)
N2	0.0397 (17)	0.0412 (17)	0.0361 (16)	-0.0136 (14)	-0.0071 (14)	-0.0077 (14)
C1	0.039 (2)	0.039 (2)	0.0376 (19)	-0.0126 (16)	-0.0050 (16)	-0.0109 (16)
C2	0.046 (2)	0.052 (2)	0.067 (3)	-0.023 (2)	-0.015 (2)	0.003 (2)
C3	0.042 (2)	0.068 (3)	0.073 (3)	-0.024 (2)	-0.018 (2)	0.001 (2)
C4	0.044 (2)	0.051 (2)	0.045 (2)	-0.0073 (18)	-0.0155 (18)	-0.0092 (19)
C5	0.050 (2)	0.049 (2)	0.049 (2)	-0.0175 (19)	-0.0138 (19)	-0.0045 (19)
C6	0.039 (2)	0.049 (2)	0.053 (2)	-0.0160 (18)	-0.0125 (18)	-0.0092 (19)
C7	0.041 (2)	0.040 (2)	0.0349 (19)	-0.0133 (16)	-0.0021 (17)	-0.0056 (16)
C8	0.044 (2)	0.074 (3)	0.052 (3)	0.000 (2)	-0.008 (2)	-0.021 (2)
C9	0.045 (2)	0.046 (2)	0.038 (2)	-0.0197 (18)	-0.0028 (17)	-0.0095 (17)
C10	0.038 (2)	0.046 (2)	0.043 (2)	-0.0148 (18)	-0.0075 (17)	-0.0068 (18)
C11	0.107 (4)	0.164 (6)	0.040 (3)	-0.084 (4)	-0.012 (3)	0.012 (3)
C12	0.137 (6)	0.154 (6)	0.052 (3)	-0.068 (5)	-0.029 (4)	0.002 (4)
Br2	0.0752 (4)	0.0956 (4)	0.0968 (4)	-0.0393 (3)	-0.0383 (3)	-0.0181 (3)
S2	0.0412 (5)	0.0483 (6)	0.0339 (5)	-0.0158 (4)	-0.0065 (4)	-0.0066 (4)
O5	0.0492 (16)	0.0646 (19)	0.0545 (17)	-0.0135 (14)	0.0028 (14)	-0.0311 (15)
O6	0.0692 (19)	0.0684 (19)	0.0463 (16)	-0.0249 (16)	-0.0256 (14)	0.0111 (14)
O7	0.069 (2)	0.0466 (18)	0.0585 (18)	0.0060 (15)	-0.0043 (16)	-0.0199 (15)
O8	0.0664 (19)	0.0519 (17)	0.0427 (16)	-0.0043 (14)	-0.0066 (14)	-0.0082 (14)
N3	0.0303 (16)	0.0441 (18)	0.0438 (18)	-0.0091 (13)	-0.0073 (13)	-0.0044 (14)
N4	0.0411 (17)	0.0359 (16)	0.0374 (16)	-0.0107 (14)	-0.0044 (14)	-0.0095 (13)
C13	0.038 (2)	0.040 (2)	0.0370 (19)	-0.0138 (16)	-0.0103 (16)	-0.0019 (16)
C14	0.040 (2)	0.054 (3)	0.069 (3)	-0.0041 (19)	-0.012 (2)	-0.022 (2)
C15	0.038 (2)	0.076 (3)	0.084 (3)	-0.006 (2)	-0.021 (2)	-0.022 (3)
C16	0.051 (2)	0.059 (3)	0.054 (2)	-0.025 (2)	-0.020 (2)	-0.004 (2)
C17	0.056 (3)	0.043 (2)	0.055 (2)	-0.0112 (19)	-0.018 (2)	-0.0119 (19)
C18	0.041 (2)	0.042 (2)	0.051 (2)	-0.0057 (17)	-0.0147 (18)	-0.0069 (18)
C19	0.042 (2)	0.042 (2)	0.040 (2)	-0.0139 (17)	-0.0009 (17)	-0.0161 (17)
C20	0.039 (2)	0.078 (3)	0.063 (3)	-0.021 (2)	-0.008 (2)	-0.006 (2)
C21	0.049 (2)	0.038 (2)	0.041 (2)	-0.0146 (17)	0.0016 (18)	-0.0097 (17)
C22	0.039 (2)	0.042 (2)	0.045 (2)	-0.0122 (18)	-0.0008 (17)	-0.0120 (18)
C23	0.105 (4)	0.096 (4)	0.045 (3)	0.001 (3)	-0.009 (3)	-0.021 (3)
C24	0.223 (10)	0.125 (6)	0.077 (4)	0.040 (6)	-0.060 (6)	-0.027 (4)

Geometric parameters ( $\text{\AA}$ ,  $\text{^{\circ}}$ )

Br1—C4	1.887 (4)	C6—H6	0.9300
Br2—C16	1.901 (5)	C8—H8B	0.9600
S1—O2	1.424 (3)	C8—H8A	0.9600
S1—N1	1.636 (3)	C8—H8C	0.9600
S1—O1	1.427 (3)	C9—H9B	0.9700
S1—C1	1.766 (4)	C9—H9A	0.9700
S2—O6	1.4280 (9)	C11—H11B	0.9700
S2—N3	1.6297 (8)	C11—H11A	0.9700
S2—C13	1.765 (4)	C12—H12A	0.9600
S2—O5	1.4271 (8)	C12—H12B	0.9600
O3—C10	1.200 (5)	C12—H12C	0.9600
O4—C11	1.467 (6)	C13—C18	1.380 (5)
O4—C10	1.311 (5)	C13—C14	1.377 (6)
O7—C22	1.196 (4)	C14—C15	1.375 (7)
O8—C23	1.462 (5)	C15—C16	1.366 (6)
O8—C22	1.319 (4)	C16—C17	1.375 (6)
N1—N2	1.420 (4)	C17—C18	1.375 (6)
N2—C7	1.270 (5)	C19—C20	1.492 (7)
N1—H1	0.8600	C19—C21	1.496 (5)
N3—N4	1.409 (3)	C21—C22	1.503 (5)
N4—C19	1.278 (5)	C23—C24	1.425 (10)
N3—H3A	0.8600	C14—H14	0.9300
C1—C2	1.370 (6)	C15—H15	0.9300
C1—C6	1.380 (5)	C17—H17	0.9300
C2—C3	1.374 (7)	C18—H18	0.9300
C3—C4	1.369 (6)	C20—H20A	0.9600
C4—C5	1.376 (7)	C20—H20B	0.9600
C5—C6	1.375 (6)	C20—H20C	0.9600
C7—C9	1.501 (5)	C21—H21A	0.9700
C7—C8	1.498 (6)	C21—H21B	0.9700
C9—C10	1.503 (5)	C23—H23A	0.9700
C11—C12	1.413 (7)	C23—H23B	0.9700
C2—H2	0.9300	C24—H24A	0.9600
C3—H3	0.9300	C24—H24B	0.9600
C5—H5	0.9300	C24—H24C	0.9600
O1—S1—O2	120.04 (17)	O4—C11—H11B	109.00
O1—S1—N1	108.05 (17)	C12—C11—H11A	109.00
O1—S1—C1	107.54 (18)	H11A—C11—H11B	108.00
O2—S1—N1	104.25 (19)	O4—C11—H11A	109.00
O2—S1—C1	109.86 (19)	C12—C11—H11B	109.00
N1—S1—C1	106.29 (17)	C11—C12—H12B	109.00
O5—S2—O6	120.45 (5)	H12A—C12—H12C	109.00
O5—S2—N3	107.69 (5)	H12B—C12—H12C	109.00
O5—S2—C13	107.71 (12)	C11—C12—H12A	109.00
O6—S2—N3	103.84 (5)	C11—C12—H12C	110.00
O6—S2—C13	108.70 (12)	H12A—C12—H12B	109.00
N3—S2—C13	107.84 (13)	C14—C13—C18	120.7 (4)

C10—O4—C11	116.9 (4)	S2—C13—C14	119.8 (3)
C22—O8—C23	116.2 (3)	S2—C13—C18	119.4 (3)
S1—N1—N2	113.1 (3)	C13—C14—C15	119.4 (4)
N1—N2—C7	114.7 (3)	C14—C15—C16	119.4 (4)
S1—N1—H1	123.00	Br2—C16—C17	119.0 (3)
N2—N1—H1	124.00	C15—C16—C17	121.9 (4)
S2—N3—N4	114.28 (15)	Br2—C16—C15	119.2 (4)
N3—N4—C19	115.1 (3)	C16—C17—C18	118.7 (4)
S2—N3—H3A	123.00	C13—C18—C17	119.8 (4)
N4—N3—H3A	123.00	N4—C19—C20	126.0 (4)
C2—C1—C6	120.9 (4)	C20—C19—C21	118.9 (4)
S1—C1—C6	119.6 (3)	N4—C19—C21	115.1 (4)
S1—C1—C2	119.5 (3)	C19—C21—C22	113.0 (3)
C1—C2—C3	119.4 (4)	O7—C22—O8	124.3 (3)
C2—C3—C4	119.7 (4)	O7—C22—C21	124.4 (3)
C3—C4—C5	121.4 (4)	O8—C22—C21	111.3 (3)
Br1—C4—C5	120.7 (3)	O8—C23—C24	108.9 (5)
Br1—C4—C3	117.9 (4)	C13—C14—H14	120.00
C4—C5—C6	118.9 (4)	C15—C14—H14	120.00
C1—C6—C5	119.8 (4)	C14—C15—H15	120.00
C8—C7—C9	118.9 (3)	C16—C15—H15	120.00
N2—C7—C9	115.2 (3)	C16—C17—H17	121.00
N2—C7—C8	125.9 (3)	C18—C17—H17	121.00
C7—C9—C10	112.1 (3)	C13—C18—H18	120.00
O4—C10—C9	111.9 (3)	C17—C18—H18	120.00
O3—C10—C9	124.1 (3)	C19—C20—H20A	109.00
O3—C10—O4	124.0 (3)	C19—C20—H20B	109.00
O4—C11—C12	111.3 (5)	C19—C20—H20C	109.00
C3—C2—H2	120.00	H20A—C20—H20B	109.00
C1—C2—H2	120.00	H20A—C20—H20C	109.00
C4—C3—H3	120.00	H20B—C20—H20C	110.00
C2—C3—H3	120.00	C19—C21—H21A	109.00
C6—C5—H5	121.00	C19—C21—H21B	109.00
C4—C5—H5	121.00	C22—C21—H21A	109.00
C5—C6—H6	120.00	C22—C21—H21B	109.00
C1—C6—H6	120.00	H21A—C21—H21B	108.00
C7—C8—H8C	109.00	O8—C23—H23A	110.00
H8B—C8—H8C	109.00	O8—C23—H23B	110.00
H8A—C8—H8C	109.00	C24—C23—H23A	110.00
C7—C8—H8B	109.00	C24—C23—H23B	110.00
C7—C8—H8A	110.00	H23A—C23—H23B	108.00
H8A—C8—H8B	109.00	C23—C24—H24A	109.00
C7—C9—H9A	109.00	C23—C24—H24B	109.00
C10—C9—H9B	109.00	C23—C24—H24C	109.00
C10—C9—H9A	109.00	H24A—C24—H24B	110.00
C7—C9—H9B	109.00	H24A—C24—H24C	109.00
H9A—C9—H9B	108.00	H24B—C24—H24C	109.00
O1—S1—N1—N2		N3—N4—C19—C21	
—63.0 (3)		180.0 (3)	

O2—S1—N1—N2	168.3 (3)	S1—C1—C2—C3	176.9 (4)
C1—S1—N1—N2	52.2 (3)	C6—C1—C2—C3	-1.6 (7)
O1—S1—C1—C2	-0.5 (4)	S1—C1—C6—C5	-176.6 (3)
O1—S1—C1—C6	178.0 (3)	C2—C1—C6—C5	1.9 (6)
O2—S1—C1—C2	131.8 (3)	C1—C2—C3—C4	0.4 (7)
O2—S1—C1—C6	-49.8 (4)	C2—C3—C4—Br1	-176.2 (4)
N1—S1—C1—C2	-116.0 (4)	C2—C3—C4—C5	0.4 (7)
N1—S1—C1—C6	62.5 (3)	Br1—C4—C5—C6	176.5 (3)
C13—S2—N3—N4	59.0 (2)	C3—C4—C5—C6	-0.1 (6)
O5—S2—C13—C14	-7.2 (4)	C4—C5—C6—C1	-1.0 (6)
O5—S2—C13—C18	174.8 (3)	N2—C7—C9—C10	-115.5 (4)
O6—S2—C13—C14	124.8 (3)	C8—C7—C9—C10	64.9 (4)
O6—S2—C13—C18	-53.1 (3)	C7—C9—C10—O3	3.5 (5)
O5—S2—N3—N4	-57.04 (16)	C7—C9—C10—O4	-177.7 (3)
O6—S2—N3—N4	174.17 (15)	S2—C13—C14—C15	-177.7 (4)
N3—S2—C13—C18	58.9 (3)	C18—C13—C14—C15	0.2 (7)
N3—S2—C13—C14	-123.2 (3)	S2—C13—C18—C17	178.1 (3)
C11—O4—C10—C9	176.5 (4)	C14—C13—C18—C17	0.2 (6)
C11—O4—C10—O3	-4.8 (6)	C13—C14—C15—C16	-0.5 (7)
C10—O4—C11—C12	174.6 (4)	C14—C15—C16—Br2	-178.9 (4)
C23—O8—C22—O7	-0.5 (6)	C14—C15—C16—C17	0.4 (7)
C22—O8—C23—C24	171.0 (6)	Br2—C16—C17—C18	179.4 (3)
C23—O8—C22—C21	179.7 (4)	C15—C16—C17—C18	0.0 (6)
S1—N1—N2—C7	174.9 (3)	C16—C17—C18—C13	-0.3 (6)
N1—N2—C7—C8	-2.5 (5)	N4—C19—C21—C22	-113.9 (4)
N1—N2—C7—C9	177.9 (3)	C20—C19—C21—C22	66.6 (5)
S2—N3—N4—C19	172.6 (2)	C19—C21—C22—O7	-5.4 (6)
N3—N4—C19—C20	-0.5 (5)	C19—C21—C22—O8	174.4 (3)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O3 <sup>i</sup>	0.86	2.46	2.909 (5)	114
N3—H3A···O7 <sup>ii</sup>	0.86	2.36	2.8855 (13)	120
C3—H3···O4 <sup>iii</sup>	0.93	2.53	3.450 (6)	172

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $-x+2, -y+1, -z$ ; (iii)  $-x+2, -y, -z+1$ .