

4-[(5-Bromo-2-hydroxybenzylidene)-amino]-3-propyl-1*H*-1,2,4-triazole-5(*4H*)-thione

Xin Wu,^a Cai-Xia Yuan,^{b*} Ling Ma,^c Kai-Lu Zhai^d and Miao-Li Zhu^b

^aCollege of Arts and Sciences, Shanxi Agricultural University, Taigu, Shanxi 030801, People's Republic of China, ^bInstitute of Molecular Science, Key Laboratory of Chemical Biology and Molecular Engineering of the Education Ministry, Shanxi University, Taiyuan, Shanxi 030006, People's Republic of China, ^cDepartment of Biochemistry and Molecular Biology, Shanxi Medical University, Taiyuan, Shanxi 030001, People's Republic of China, and ^dJincheng Tap Water Company, Jincheng, Shanxi 048000, People's Republic of China

Correspondence e-mail: cx yuan@sxu.edu.cn, miaoli@sxu.edu.cn


Received 7 April 2012; accepted 2 May 2012

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.043; wR factor = 0.102; data-to-parameter ratio = 14.2.

The asymmetric unit of the title compound, $\text{C}_{12}\text{H}_{13}\text{BrN}_4\text{OS}$, contains two independent molecules in which the dihedral angles between the triazole and benzene rings are 2.9 (3) and 7.5 (3) $^\circ$. The thione group is of the form $\text{R}_2\text{C}=\text{S}$. An intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond occurs in each molecule. The crystal structure features weak $\text{N}-\text{H}\cdots\text{S}$ interactions and $\pi-\pi$ stacking of the benzene rings [centroid–centroid distance = 3.667 (3) \AA].

Related literature

For the pharmacological activity of 1,2,4-triazole-substituted and Schiff base compounds, see: Isloor *et al.* (2009); Ma *et al.* (2011). For copper complexes containing 1,2,4-triazole Schiff base ligands, see: Wen *et al.* (2004).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{13}\text{BrN}_4\text{OS}$

$M_r = 341.23$

Triclinic, $P\bar{1}$	$V = 1399.5 (15)\text{ \AA}^3$
$a = 8.042 (5)\text{ \AA}$	$Z = 4$
$b = 13.187 (8)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 13.408 (8)\text{ \AA}$	$\mu = 3.08\text{ mm}^{-1}$
$\alpha = 97.406 (9)^\circ$	$T = 298\text{ K}$
$\beta = 92.956 (10)^\circ$	$0.30 \times 0.25 \times 0.20\text{ mm}$
$\gamma = 95.916 (9)^\circ$	

Data collection

Bruker SMART 1K CCD diffractometer	14278 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2000)	4939 independent reflections
$T_{\min} = 0.458$, $T_{\max} = 0.578$	3061 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.055$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$	347 parameters
$wR(F^2) = 0.102$	H-atom parameters constrained
$S = 0.93$	$\Delta\rho_{\max} = 0.49\text{ e \AA}^{-3}$
4939 reflections	$\Delta\rho_{\min} = -0.55\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots N4	0.82	1.93	2.644 (4)	145
O2—H2 \cdots N8	0.82	1.90	2.612 (4)	144
N1—H1A \cdots S1 ⁱ	0.86	2.46	3.300 (4)	164
N5—H5 \cdots S2 ⁱⁱ	0.86	2.41	3.252 (4)	165

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $-x, -y + 1, -z$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *pubLCIF* (Westrip, 2010).

The authors acknowledge financial support from the National Natural Science Foundation of China for Youth (grant No. 21001070), the Specialized Research Fund for the Doctoral Program of Higher Education (grant No. 20111401110002) and the Natural Science Foundation of Shanxi Province of China (grant No. 2011021006-2).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2128).

References

- Bruker (2000). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Isloor, A. M., Kalluraya, B. & Shetty, P. (2009). *Eur. J. Med. Chem.* **44**, 3784–3787.
- Ma, L., Lu, L. P., Zhu, M. L., Wang, Q. M., Li, Y., Xing, S., Fu, X. Q., Gao, Z. Q. & Dong, Y. H. (2011). *Dalton Trans.* **40**, 6532–6540.
- Sheldrick, G. M. (2000). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wen, L. R., Li, M., Wang, S. W., Zhang, S. S., Li, X. M. & Ou Yang, P. K. (2004). *Chem. Res. Chin. Univ.* **20**, 722–724.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o1674 [doi:10.1107/S1600536812019745]

4-[(5-Bromo-2-hydroxybenzylidene)amino]-3-propyl-1*H*-1,2,4-triazole-5(4*H*)-thione

Xin Wu, Cai-Xia Yuan, Ling Ma, Kai-Lu Zhai and Miao-Li Zhu

Comment

1,2,4-triazole and its derivatives possess a variety of pharmacological properties (Isloor *et al.*, 2009). Schiff bases derived from substituted salicylaldehydes form numerous metal complexes (Wen *et al.*, 2004). Also, copper complexes containing 1,2,4-triazole Schiff base ligands are potential inhibitors of PTP1B (protein tyrosine phosphatase 1B), TCPTP (T-cell protein tyrosine phosphatase), PTP-MEG2 (megakaryocyte protein tyrosine phosphatase) and SHP-1 (Src homology phosphatase 1) (Ma *et al.*, 2011). In continuation of our work in this area, we report here the synthesis and crystal structure of the title compound, $C_{12}H_{13}BrN_4OS$, (I).

The title compound, (I), crystallizes with two independent molecules in the asymmetric unit (Fig. 1). The dihedral angles between the triazole and phenyl rings are $2.9(3)^\circ$ and $7.5(3)^\circ$, respectively. The thione group is of the form $R_2C=S$ with a $Cdb\S$ distance of $1.673(4)\text{\AA}$ and $1.675(4)\text{\AA}$ indicating significant double bond character. Intramolecular O—H \cdots N hydrogen bonds, weak N—H \cdots S intermolecular interactions (Table 1) and π — π stacking of the benzene rings [centroid–centroid distance = $3.667(3)\text{\AA}$] are observed that may influence crystal packing.

Experimental


0.5 mmol 5-Bromosalicylaldehyde in 10 ml of ethanol was added to a solution of 4-amino-5-propyl-1,2,4-triazole-3-thione (0.5 mmol) in 20 ml of ethanol, and then refluxed for 2 h. The resulting solution was filtered and recrystallized from ethanol. X-ray quality Yellow crystals of the title compound were formed upon slow evaporation of the resulting soluton.

Refinement


All of the H atoms were placed in their calculated positions and the refined using the riding model with C—H lengths of 0.93\AA (CH), 0.97\AA (CH_2), 0.97\AA (CH_3), 0.86\AA (NH) or 0.82\AA . The isotropic displacement parameters for these atoms were set to 1.2 (CH , CH_2 , O, N) or 1.5 (CH_3) times U_{eq} of the parent atom.

Computing details

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT* (Bruker, 2000); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2010).

**Figure 1**

Molecular structure of the title compound, (I), with displacement ellipsoids drawn at the 50% probability level. Dashed lines indicate O—H···N intramolecular hydrogen bonding.

**Figure 2**

Packing of the title compound, (I), viewed along the a axis. Dashed lines indicate $\text{O}-\text{H}\cdots\text{N}$ intramolecular hydrogen bonds and weak $\text{N}-\text{H}\cdots\text{S}$ intermolecular interactions. Remaining hydrogen atoms have been removed for clarity.

4-[(5-Bromo-2-hydroxybenzylidene)amino]-3-propyl-1*H*-1,2,4-triazole- 5(4*H*)-thione

Crystal data

$\text{C}_{12}\text{H}_{13}\text{BrN}_4\text{OS}$
 $M_r = 341.23$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 8.042 (5)$ Å
 $b = 13.187 (8)$ Å
 $c = 13.408 (8)$ Å
 $\alpha = 97.406 (9)^\circ$
 $\beta = 92.956 (10)^\circ$
 $\gamma = 95.916 (9)^\circ$
 $V = 1399.5 (15)$ Å³

$Z = 4$
 $F(000) = 688$
 $D_x = 1.620 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 2374 reflections
 $\theta = 2.5\text{--}23.6^\circ$
 $\mu = 3.08 \text{ mm}^{-1}$
 $T = 298 \text{ K}$
Block, yellow
 $0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker SMART 1K CCD diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: 11.72 pixels mm⁻¹
 ω scans
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 2000)
 $T_{\min} = 0.458$, $T_{\max} = 0.578$

14278 measured reflections
 4939 independent reflections
 3061 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.5^\circ$
 $h = -9 \rightarrow 9$
 $k = -15 \rightarrow 15$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.102$
 $S = 0.93$
 4939 reflections
 347 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0516P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.49 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.55 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.38916 (7)	0.16296 (4)	0.59088 (4)	0.06736 (19)
S1	0.49656 (16)	0.40893 (8)	0.14062 (8)	0.0530 (3)
O1	0.1987 (5)	0.5771 (2)	0.5179 (2)	0.0666 (10)
H1	0.2089	0.5860	0.4590	0.100*
N1	0.4061 (5)	0.5891 (3)	0.0953 (2)	0.0521 (10)
H1A	0.4451	0.5805	0.0366	0.062*
N2	0.3314 (5)	0.6750 (3)	0.1302 (3)	0.0512 (10)
N3	0.3366 (4)	0.5663 (2)	0.2414 (2)	0.0378 (8)
N4	0.3044 (4)	0.5369 (2)	0.3350 (2)	0.0400 (8)
C1	0.4132 (5)	0.5203 (3)	0.1597 (3)	0.0389 (10)
C2	0.2882 (5)	0.6584 (3)	0.2197 (3)	0.0424 (10)
C3	0.1971 (6)	0.7272 (3)	0.2879 (3)	0.0462 (11)
H3A	0.2588	0.7428	0.3531	0.055*
H3B	0.0881	0.6922	0.2975	0.055*
C4	0.1740 (6)	0.8271 (3)	0.2457 (3)	0.0540 (12)
H4A	0.2832	0.8624	0.2370	0.065*

H4B	0.1143	0.8111	0.1799	0.065*
C5	0.0785 (7)	0.8980 (4)	0.3133 (4)	0.0833 (18)
H5A	-0.0316	0.8647	0.3197	0.125*
H5B	0.0695	0.9604	0.2844	0.125*
H5C	0.1368	0.9140	0.3786	0.125*
C6	0.3493 (5)	0.4529 (3)	0.3590 (3)	0.0420 (11)
H6	0.4027	0.4104	0.3131	0.050*
C7	0.3181 (5)	0.4230 (3)	0.4566 (3)	0.0368 (10)
C8	0.2452 (5)	0.4844 (3)	0.5321 (3)	0.0466 (11)
C9	0.2203 (6)	0.4511 (4)	0.6243 (3)	0.0554 (13)
H9	0.1734	0.4926	0.6743	0.066*
C10	0.2647 (6)	0.3566 (4)	0.6423 (3)	0.0562 (13)
H10	0.2475	0.3341	0.7043	0.067*
C11	0.3349 (5)	0.2955 (3)	0.5681 (3)	0.0458 (11)
C12	0.3633 (5)	0.3282 (3)	0.4773 (3)	0.0442 (11)
H12	0.4133	0.2868	0.4287	0.053*
Br2	0.51614 (6)	-0.08187 (4)	-0.31301 (4)	0.06348 (19)
S2	0.14032 (16)	0.37417 (8)	-0.08568 (8)	0.0509 (3)
O2	0.1831 (4)	-0.0214 (2)	0.0781 (2)	0.0544 (8)
H2	0.1680	0.0389	0.0943	0.082*
N6	-0.0208 (5)	0.3668 (3)	0.1823 (2)	0.0499 (9)
N5	0.0102 (5)	0.4074 (2)	0.0951 (2)	0.0516 (10)
H5	-0.0145	0.4674	0.0853	0.062*
N7	0.0963 (4)	0.2607 (2)	0.0741 (2)	0.0374 (8)
N8	0.1549 (4)	0.1671 (2)	0.0448 (2)	0.0385 (8)
C13	0.0814 (5)	0.3467 (3)	0.0267 (3)	0.0418 (10)
C14	0.0324 (5)	0.2768 (3)	0.1686 (3)	0.0413 (10)
C15	0.0280 (5)	0.2020 (3)	0.2411 (3)	0.0441 (11)
H15A	0.1416	0.1892	0.2590	0.053*
H15B	-0.0325	0.1375	0.2094	0.053*
C16	-0.0556 (6)	0.2392 (3)	0.3367 (3)	0.0495 (12)
H16A	0.0035	0.3043	0.3679	0.059*
H16B	-0.1700	0.2505	0.3191	0.059*
C17	-0.0559 (7)	0.1630 (4)	0.4107 (3)	0.0694 (15)
H17A	-0.1174	0.0991	0.3808	0.104*
H17B	-0.1078	0.1893	0.4701	0.104*
H17C	0.0573	0.1519	0.4285	0.104*
C18	0.2235 (5)	0.1495 (3)	-0.0384 (3)	0.0414 (10)
H18	0.2378	0.2006	-0.0799	0.050*
C19	0.2791 (5)	0.0495 (3)	-0.0681 (3)	0.0368 (10)
C20	0.2579 (5)	-0.0309 (3)	-0.0101 (3)	0.0395 (10)
C21	0.3146 (6)	-0.1252 (3)	-0.0421 (3)	0.0497 (12)
H21	0.3009	-0.1784	-0.0030	0.060*
C22	0.3909 (5)	-0.1398 (3)	-0.1314 (3)	0.0490 (11)
H22	0.4284	-0.2029	-0.1529	0.059*
C23	0.4117 (5)	-0.0604 (3)	-0.1893 (3)	0.0428 (10)
C24	0.3565 (5)	0.0322 (3)	-0.1588 (3)	0.0427 (11)
H24	0.3704	0.0846	-0.1989	0.051*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0834 (4)	0.0674 (3)	0.0629 (3)	0.0222 (3)	0.0151 (3)	0.0374 (3)
S1	0.0863 (9)	0.0448 (7)	0.0363 (6)	0.0296 (6)	0.0219 (6)	0.0124 (5)
O1	0.104 (3)	0.053 (2)	0.053 (2)	0.0347 (19)	0.038 (2)	0.0096 (16)
N1	0.081 (3)	0.045 (2)	0.039 (2)	0.027 (2)	0.028 (2)	0.0140 (17)
N2	0.076 (3)	0.042 (2)	0.043 (2)	0.0232 (19)	0.025 (2)	0.0148 (17)
N3	0.054 (2)	0.0365 (19)	0.0274 (18)	0.0125 (16)	0.0092 (16)	0.0128 (14)
N4	0.052 (2)	0.040 (2)	0.0313 (19)	0.0078 (17)	0.0147 (17)	0.0110 (15)
C1	0.052 (3)	0.039 (2)	0.030 (2)	0.011 (2)	0.008 (2)	0.0099 (18)
C2	0.048 (3)	0.043 (2)	0.040 (2)	0.011 (2)	0.010 (2)	0.0129 (19)
C3	0.057 (3)	0.039 (2)	0.049 (3)	0.018 (2)	0.021 (2)	0.015 (2)
C4	0.072 (3)	0.045 (3)	0.051 (3)	0.019 (2)	0.018 (3)	0.017 (2)
C5	0.115 (5)	0.056 (3)	0.097 (4)	0.045 (3)	0.057 (4)	0.028 (3)
C6	0.061 (3)	0.037 (2)	0.031 (2)	0.010 (2)	0.016 (2)	0.0079 (18)
C7	0.047 (3)	0.038 (2)	0.025 (2)	0.0033 (19)	0.0080 (19)	0.0050 (17)
C8	0.052 (3)	0.051 (3)	0.039 (3)	0.009 (2)	0.015 (2)	0.009 (2)
C9	0.074 (4)	0.061 (3)	0.034 (3)	0.013 (3)	0.023 (2)	0.003 (2)
C10	0.062 (3)	0.078 (4)	0.034 (3)	0.008 (3)	0.014 (2)	0.022 (2)
C11	0.052 (3)	0.049 (3)	0.040 (2)	0.004 (2)	0.009 (2)	0.020 (2)
C12	0.060 (3)	0.043 (2)	0.032 (2)	0.009 (2)	0.009 (2)	0.0082 (19)
Br2	0.0799 (4)	0.0558 (3)	0.0581 (3)	0.0235 (3)	0.0309 (3)	-0.0026 (2)
S2	0.0755 (9)	0.0429 (6)	0.0419 (6)	0.0249 (6)	0.0211 (6)	0.0133 (5)
O2	0.085 (2)	0.0426 (17)	0.0434 (18)	0.0200 (17)	0.0240 (17)	0.0158 (14)
N6	0.068 (3)	0.049 (2)	0.037 (2)	0.0223 (19)	0.0166 (19)	0.0066 (17)
N5	0.083 (3)	0.039 (2)	0.041 (2)	0.0294 (19)	0.023 (2)	0.0109 (16)
N7	0.051 (2)	0.0321 (18)	0.0332 (18)	0.0170 (16)	0.0127 (16)	0.0046 (14)
N8	0.051 (2)	0.0298 (18)	0.0369 (19)	0.0156 (16)	0.0104 (17)	0.0020 (14)
C13	0.055 (3)	0.034 (2)	0.038 (2)	0.015 (2)	0.007 (2)	0.0039 (18)
C14	0.049 (3)	0.041 (3)	0.036 (2)	0.013 (2)	0.013 (2)	0.0047 (19)
C15	0.055 (3)	0.044 (2)	0.037 (2)	0.015 (2)	0.010 (2)	0.0093 (19)
C16	0.063 (3)	0.053 (3)	0.034 (2)	0.015 (2)	0.014 (2)	0.002 (2)
C17	0.097 (4)	0.076 (4)	0.042 (3)	0.023 (3)	0.015 (3)	0.019 (3)
C18	0.055 (3)	0.036 (2)	0.037 (2)	0.014 (2)	0.014 (2)	0.0087 (18)
C19	0.044 (3)	0.031 (2)	0.037 (2)	0.0097 (19)	0.005 (2)	0.0057 (18)
C20	0.046 (3)	0.036 (2)	0.039 (2)	0.010 (2)	0.006 (2)	0.0096 (19)
C21	0.067 (3)	0.030 (2)	0.055 (3)	0.010 (2)	0.004 (3)	0.013 (2)
C22	0.060 (3)	0.037 (2)	0.053 (3)	0.021 (2)	0.007 (2)	0.001 (2)
C23	0.047 (3)	0.042 (2)	0.041 (2)	0.013 (2)	0.013 (2)	0.0011 (19)
C24	0.055 (3)	0.039 (2)	0.039 (2)	0.017 (2)	0.014 (2)	0.0085 (19)

Geometric parameters (\AA , ^\circ)

Br1—C11	1.902 (4)	Br2—C23	1.900 (4)
S1—C1	1.673 (4)	S2—C13	1.675 (4)
O1—C8	1.348 (5)	O2—C20	1.351 (4)
O1—H1	0.8200	O2—H2	0.8200
N1—C1	1.334 (5)	N6—C14	1.297 (5)
N1—N2	1.377 (4)	N6—N5	1.371 (4)

N1—H1A	0.8600	N5—C13	1.330 (5)
N2—C2	1.304 (5)	N5—H5	0.8600
N3—C2	1.375 (5)	N7—C13	1.381 (5)
N3—C1	1.385 (5)	N7—N8	1.384 (4)
N3—N4	1.390 (4)	N7—C14	1.391 (5)
N4—C6	1.274 (5)	N8—C18	1.274 (4)
C2—C3	1.479 (5)	C14—C15	1.470 (5)
C3—C4	1.524 (5)	C15—C16	1.527 (5)
C3—H3A	0.9700	C15—H15A	0.9700
C3—H3B	0.9700	C15—H15B	0.9700
C4—C5	1.512 (6)	C16—C17	1.500 (6)
C4—H4A	0.9700	C16—H16A	0.9700
C4—H4B	0.9700	C16—H16B	0.9700
C5—H5A	0.9600	C17—H17A	0.9600
C5—H5B	0.9600	C17—H17B	0.9600
C5—H5C	0.9600	C17—H17C	0.9600
C6—C7	1.442 (5)	C18—C19	1.450 (5)
C6—H6	0.9300	C18—H18	0.9300
C7—C12	1.394 (5)	C19—C20	1.395 (5)
C7—C8	1.404 (5)	C19—C24	1.398 (5)
C8—C9	1.381 (5)	C20—C21	1.392 (5)
C9—C10	1.378 (6)	C21—C22	1.374 (6)
C9—H9	0.9300	C21—H21	0.9300
C10—C11	1.380 (6)	C22—C23	1.383 (6)
C10—H10	0.9300	C22—H22	0.9300
C11—C12	1.366 (5)	C23—C24	1.363 (5)
C12—H12	0.9300	C24—H24	0.9300
C8—O1—H1	109.5	C20—O2—H2	109.5
C1—N1—N2	114.6 (3)	C14—N6—N5	104.5 (3)
C1—N1—H1A	122.7	C13—N5—N6	114.8 (3)
N2—N1—H1A	122.7	C13—N5—H5	122.6
C2—N2—N1	104.0 (3)	N6—N5—H5	122.6
C2—N3—C1	109.1 (3)	C13—N7—N8	133.1 (3)
C2—N3—N4	118.4 (3)	C13—N7—C14	108.8 (3)
C1—N3—N4	132.6 (3)	N8—N7—C14	118.0 (3)
C6—N4—N3	120.6 (3)	C18—N8—N7	121.2 (3)
N1—C1—N3	102.0 (3)	N5—C13—N7	102.2 (3)
N1—C1—S1	126.4 (3)	N5—C13—S2	126.8 (3)
N3—C1—S1	131.5 (3)	N7—C13—S2	131.0 (3)
N2—C2—N3	110.3 (3)	N6—C14—N7	109.7 (3)
N2—C2—C3	125.3 (4)	N6—C14—C15	125.9 (3)
N3—C2—C3	124.4 (3)	N7—C14—C15	124.3 (3)
C2—C3—C4	111.6 (3)	C14—C15—C16	112.5 (3)
C2—C3—H3A	109.3	C14—C15—H15A	109.1
C4—C3—H3A	109.3	C16—C15—H15A	109.1
C2—C3—H3B	109.3	C14—C15—H15B	109.1
C4—C3—H3B	109.3	C16—C15—H15B	109.1
H3A—C3—H3B	108.0	H15A—C15—H15B	107.8

C5—C4—C3	112.6 (3)	C17—C16—C15	111.8 (3)
C5—C4—H4A	109.1	C17—C16—H16A	109.3
C3—C4—H4A	109.1	C15—C16—H16A	109.3
C5—C4—H4B	109.1	C17—C16—H16B	109.3
C3—C4—H4B	109.1	C15—C16—H16B	109.3
H4A—C4—H4B	107.8	H16A—C16—H16B	107.9
C4—C5—H5A	109.5	C16—C17—H17A	109.5
C4—C5—H5B	109.5	C16—C17—H17B	109.5
H5A—C5—H5B	109.5	H17A—C17—H17B	109.5
C4—C5—H5C	109.5	C16—C17—H17C	109.5
H5A—C5—H5C	109.5	H17A—C17—H17C	109.5
H5B—C5—H5C	109.5	H17B—C17—H17C	109.5
N4—C6—C7	120.7 (4)	N8—C18—C19	119.7 (3)
N4—C6—H6	119.7	N8—C18—H18	120.1
C7—C6—H6	119.7	C19—C18—H18	120.1
C12—C7—C8	118.3 (4)	C20—C19—C24	118.3 (3)
C12—C7—C6	118.2 (3)	C20—C19—C18	122.8 (3)
C8—C7—C6	123.5 (4)	C24—C19—C18	118.9 (3)
O1—C8—C9	118.0 (4)	O2—C20—C21	117.4 (3)
O1—C8—C7	121.8 (4)	O2—C20—C19	122.4 (3)
C9—C8—C7	120.3 (4)	C21—C20—C19	120.2 (4)
C10—C9—C8	120.2 (4)	C22—C21—C20	120.2 (4)
C10—C9—H9	119.9	C22—C21—H21	119.9
C8—C9—H9	119.9	C20—C21—H21	119.9
C9—C10—C11	119.7 (4)	C21—C22—C23	119.8 (4)
C9—C10—H10	120.1	C21—C22—H22	120.1
C11—C10—H10	120.1	C23—C22—H22	120.1
C12—C11—C10	120.7 (4)	C24—C23—C22	120.6 (4)
C12—C11—Br1	119.3 (3)	C24—C23—Br2	120.0 (3)
C10—C11—Br1	120.0 (3)	C22—C23—Br2	119.4 (3)
C11—C12—C7	120.7 (4)	C23—C24—C19	120.9 (4)
C11—C12—H12	119.6	C23—C24—H24	119.6
C7—C12—H12	119.6	C19—C24—H24	119.6
C1—N1—N2—C2	-0.8 (5)	C14—N6—N5—C13	-0.2 (5)
C2—N3—N4—C6	-178.9 (4)	C13—N7—N8—C18	6.2 (6)
C1—N3—N4—C6	0.5 (6)	C14—N7—N8—C18	-176.1 (4)
N2—N1—C1—N3	0.2 (5)	N6—N5—C13—N7	0.2 (5)
N2—N1—C1—S1	-179.6 (3)	N6—N5—C13—S2	-178.1 (3)
C2—N3—C1—N1	0.5 (4)	N8—N7—C13—N5	177.7 (4)
N4—N3—C1—N1	-178.9 (4)	C14—N7—C13—N5	-0.1 (4)
C2—N3—C1—S1	-179.8 (3)	N8—N7—C13—S2	-4.1 (7)
N4—N3—C1—S1	0.8 (7)	C14—N7—C13—S2	178.1 (3)
N1—N2—C2—N3	1.1 (5)	N5—N6—C14—N7	0.0 (5)
N1—N2—C2—C3	-177.7 (4)	N5—N6—C14—C15	179.8 (4)
C1—N3—C2—N2	-1.0 (5)	C13—N7—C14—N6	0.1 (5)
N4—N3—C2—N2	178.5 (3)	N8—N7—C14—N6	-178.2 (3)
C1—N3—C2—C3	177.8 (4)	C13—N7—C14—C15	-179.7 (4)
N4—N3—C2—C3	-2.7 (6)	N8—N7—C14—C15	2.1 (6)

N2—C2—C3—C4	−5.9 (6)	N6—C14—C15—C16	2.6 (6)
N3—C2—C3—C4	175.5 (4)	N7—C14—C15—C16	−177.8 (4)
C2—C3—C4—C5	179.0 (4)	C14—C15—C16—C17	−178.8 (4)
N3—N4—C6—C7	179.1 (4)	N7—N8—C18—C19	−178.2 (3)
N4—C6—C7—C12	177.3 (4)	N8—C18—C19—C20	1.4 (6)
N4—C6—C7—C8	−3.0 (7)	N8—C18—C19—C24	−179.0 (4)
C12—C7—C8—O1	179.8 (4)	C24—C19—C20—O2	−179.4 (4)
C6—C7—C8—O1	0.0 (7)	C18—C19—C20—O2	0.3 (6)
C12—C7—C8—C9	0.3 (6)	C24—C19—C20—C21	0.7 (6)
C6—C7—C8—C9	−179.5 (4)	C18—C19—C20—C21	−179.7 (4)
O1—C8—C9—C10	179.6 (4)	O2—C20—C21—C22	179.6 (4)
C7—C8—C9—C10	−0.9 (7)	C19—C20—C21—C22	−0.4 (7)
C8—C9—C10—C11	0.3 (7)	C20—C21—C22—C23	0.2 (7)
C9—C10—C11—C12	1.0 (7)	C21—C22—C23—C24	−0.3 (7)
C9—C10—C11—Br1	−177.8 (4)	C21—C22—C23—Br2	−179.6 (3)
C10—C11—C12—C7	−1.6 (7)	C22—C23—C24—C19	0.6 (7)
Br1—C11—C12—C7	177.1 (3)	Br2—C23—C24—C19	179.9 (3)
C8—C7—C12—C11	1.0 (6)	C20—C19—C24—C23	−0.8 (6)
C6—C7—C12—C11	−179.3 (4)	C18—C19—C24—C23	179.5 (4)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···N4	0.82	1.93	2.644 (4)	145
O2—H2···N8	0.82	1.90	2.612 (4)	144
N1—H1A···S1 ⁱ	0.86	2.46	3.300 (4)	164
N5—H5···S2 ⁱⁱ	0.86	2.41	3.252 (4)	165

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x, -y+1, -z$.