

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(2,4-Difluorophenyl)-2-(1*H*-1,2,4-triazol-1-yl)ethanol

 Victor Kesternich,^a Ronald Nelson-González,^a Marcia Pérez-Fehrmann,^a Alejandro Cárdenas^b and Iván Brito^{c*}

^aDepartamento de Química, Universidad Católica del Norte, Casilla 1280, Antofagasta, Chile, ^bDepartamento de Física, Facultad de Ciencias Básicas, Universidad de Antofagasta, Casilla 170, Antofagasta, Chile, and ^cDepartamento de Química, Facultad de Ciencias Básicas, Universidad de Antofagasta, Casilla 170, Antofagasta, Chile
Correspondence e-mail: ivanbritob@yahoo.com


Received 3 May 2012; accepted 7 May 2012

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.063; wR factor = 0.149; data-to-parameter ratio = 10.5.

In the title compound, $\text{C}_{10}\text{H}_9\text{F}_2\text{N}_3\text{O}$, the dihedral angle between the mean planes of the triazole and benzene rings is $20.6(2)^\circ$. In the crystal, molecules are linked by strong $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds into chains with graph-set notation $C(9)$ along $[100]$. Weak $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{F}$ interactions are also observed.

Related literature

For phenacylazole derivatives, see: Emami *et al.* (2008, 2009). For their biological properties, see: Schiaffella *et al.* (2005). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{10}\text{H}_9\text{F}_2\text{N}_3\text{O}$ $V = 1056.8(3) \text{ \AA}^3$
 $M_r = 225.20$ $Z = 4$
 Orthorhombic, $P2_12_12_1$ $\text{Mo K}\alpha$ radiation
 $a = 5.3770(11) \text{ \AA}$ $\mu = 0.12 \text{ mm}^{-1}$
 $b = 12.598(3) \text{ \AA}$ $T = 295 \text{ K}$
 $c = 15.601(3) \text{ \AA}$ $0.60 \times 0.29 \times 0.08 \text{ mm}$

Data collection

Nonius KappaCCD area-detector 1536 independent reflections
 diffractometer 1344 reflections with $I > 2\sigma(I)$
 9658 measured reflections $R_{\text{int}} = 0.081$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.063$ 146 parameters
 $wR(F^2) = 0.149$ H-atom parameters constrained
 $S = 1.16$ $\Delta\rho_{\text{max}} = 0.16 \text{ e \AA}^{-3}$
 1536 reflections $\Delta\rho_{\text{min}} = -0.16 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{N3}^i$	0.82	2.01	2.826 (4)	171
$\text{C3}-\text{H3}\cdots\text{N2}^{ii}$	0.93	2.62	3.500 (5)	158
$\text{C8}-\text{H8B}\cdots\text{F1}^{iii}$	0.97	2.45	3.340 (5)	153
$\text{C10}-\text{H10}\cdots\text{F2}^{iv}$	0.93	2.48	3.270 (4)	142

Symmetry codes: (i) $x + \frac{1}{2}, -y - \frac{1}{2}, -z + 1$; (ii) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $x - 1, y, z$; (iv) $x, y - 1, z$.

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *DENZO-SMN* (Otwinowski & Minor, 1997); data reduction: *DENZO-SMN*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *publCIF* (Westrip, 2010).

We are grateful to the Consejo Superior de Investigaciones Científicas (CSIC) of Spain for the award of a licence for the use of the Cambridge Structural Database (CSD).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2423).

References

- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
 Emami, S., Behdad, M., Foroumadi, A., Falahati, M., Lotfali, E. & Sharifynia, S. (2009). *Chem. Biol. Drug Des.* **73**, 388–395.
 Emami, S., Foroumadi, A., Falahati, M., Lotfali, E., Rajabalian, S., Ebrahimi, S., Farahyar, S. & Shafie, A. (2008). *Bioorg. Med. Chem. Lett.* **18**, 141–146.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Nonius (2000). *COLLECT* BV, Delft, The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. **276**, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
 Schiaffella, F., Macchiarulo, M., Milanese, L., Vecchiarelli, A., Costantino, G., Pietrella, D. & Fringuelli, R. (2005). *J. Med. Chem.* **48**, 7658–7666.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o1727 [doi:10.1107/S1600536812020661]

1-(2,4-Difluorophenyl)-2-(1*H*-1,2,4-triazol-1-yl)ethanol

Victor Kesternich, Ronald Nelson-González, Marcia Pérez-Fehrmann, Alejandro Cárdenas and Iván Brito

Comment

Phenacyl azole derivatives (Emami *et al.*, 2008; Emami *et al.*, 2009) are very important due to their antifungal properties (Schiaffella *et al.*, 2005). We report herein on the synthesis and crystal structure of the title compound, a member of this important family of compounds.

In the title molecule, Fig. 1, the dihedral angle between the mean planes of the triazole and benzene rings is 20.6 (2)°.

In the crystal, molecules are linked by strong O—H...N hydrogen bonds (Fig. 2 and Table 1) into chains propagating along [100] which have a C(9) graph-set notation (Bernstein *et al.*, 1995). There are also weak C—H...N and C—H...F interactions present (Table 1).

Experimental


To a stirred solution of 1-(2,4-difluorophenyl)-2-(1*H*-1,2,4-triazol-1-yl)ethanone (6.00 g, 26.88 mmol) in methanol (50 ml), sodium borohydride (1.12 g, 29.57 mmol) dissolved in methanol (20 ml), was added drop wise. The reaction mixture was then stirred at room temperature for 30 min. After completion of the reaction, the solvent was removed under vacuum, and 25 ml of cold water was added. Extraction was performed with dichloromethane (3 × 40 ml), and the organic extracts were washed with water (3 × 30 ml). The organic phase was then dried over anhydrous sodium sulfate. After evaporation of the solvent under vacuum, the residue was purified by crystallization in ethanol to provide the title compound as colourless crystals [85% yield; M.p.: 391-393 K].

Refinement


All of the H atoms could be located in difference Fourier maps. In the final cycles of refinement they were included in calculated positions and treated as riding atoms: O—H = 0.82 Å, C—H = 0.93 and 0.97 Å for CH and CH₂ H atoms, respectively, with $U_{iso}(H) = k \times U_{eq}(O,C)$, where $k = 1.5$ for the OH H atom, and $= 1.2$ for other H atoms. In the final cycles of refinement, in the absence of significant anomalous scattering effects, 1126 Friedel pairs were merged and Δf set to zero.

Computing details

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *DENZO-SMN* (Otwinowski & Minor, 1997); data reduction: *DENZO-SMN* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *pubCIF* (Westrip, 2010).

**Figure 1**

A view of the molecular structure of the title molecule, with the atom numbering. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

A partial view along the *c* axis of the crystal packing of the title compound, showing the formation of the O—H...N hydrogen bonded chain [dashed lines; see Table 1 for details; the H-atoms not involved in hydrogen-bonding have been omitted for clarity]

1-(2,4-Difluorophenyl)-2-(1*H*-1,2,4-triazol-1-yl)ethanol

Crystal data

$C_{10}H_9F_2N_3O$

$M_r = 225.20$

Orthorhombic, $P2_12_12_1$

$a = 5.3770$ (11) Å

$b = 12.598$ (3) Å

$c = 15.601$ (3) Å

$V = 1056.8$ (3) Å³

$Z = 4$

$F(000) = 464$

$D_x = 1.415$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

$\mu = 0.12$ mm⁻¹

$T = 295$ K

Plate, colourless

$0.60 \times 0.29 \times 0.08$ mm

Data collection

Nonius KappaCCD area-detector diffractometer	1344 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.081$
Graphite monochromator	$\theta_{\text{max}} = 28.7^\circ$, $\theta_{\text{min}} = 4.2^\circ$
φ and ω scans with κ offsets	$h = 0 \rightarrow 7$
9658 measured reflections	$k = 0 \rightarrow 16$
1536 independent reflections	$l = 0 \rightarrow 21$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.063$	H-atom parameters constrained
$wR(F^2) = 0.149$	$w = 1/[\sigma^2(F_o^2) + (0.0491P)^2 + 0.3301P]$
$S = 1.16$	where $P = (F_o^2 + 2F_c^2)/3$
1536 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
146 parameters	$\Delta\rho_{\text{max}} = 0.16 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.16 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell esds are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	1.0193 (5)	0.0759 (2)	0.69521 (16)	0.0959 (9)
F2	0.7227 (6)	0.42266 (16)	0.67885 (15)	0.1001 (12)
O1	0.5799 (5)	-0.0141 (2)	0.49259 (14)	0.0733 (9)
N1	0.5125 (5)	-0.1766 (2)	0.62303 (17)	0.0583 (8)
N2	0.7087 (6)	-0.2399 (2)	0.6393 (2)	0.0725 (10)
N3	0.3927 (6)	-0.3347 (3)	0.5866 (2)	0.0716 (11)
C1	0.6906 (6)	0.1114 (2)	0.60153 (17)	0.0491 (8)
C2	0.8583 (6)	0.1469 (3)	0.6608 (2)	0.0615 (10)
C3	0.8761 (8)	0.2503 (3)	0.6881 (2)	0.0707 (11)
C4	0.7107 (8)	0.3195 (3)	0.6541 (2)	0.0686 (13)
C5	0.5350 (8)	0.2905 (3)	0.5961 (2)	0.0703 (13)
C6	0.5262 (7)	0.1855 (3)	0.5700 (2)	0.0614 (10)
C7	0.6804 (6)	-0.0045 (2)	0.57500 (18)	0.0528 (9)
C8	0.5227 (7)	-0.0634 (3)	0.6398 (2)	0.0655 (11)
C9	0.3289 (7)	-0.2346 (3)	0.5921 (2)	0.0699 (11)
C10	0.6262 (8)	-0.3331 (3)	0.6162 (3)	0.0744 (14)
H1	0.66000	-0.05800	0.46520	0.1100*
H3	0.99530	0.27150	0.72770	0.0850*
H5	0.42310	0.34000	0.57450	0.0840*

H6	0.40680	0.16460	0.53030	0.0740*
H7	0.84910	-0.03400	0.57530	0.0630*
H8A	0.59030	-0.05180	0.69660	0.0790*
H8B	0.35530	-0.03490	0.63880	0.0790*
H9	0.17440	-0.20790	0.57620	0.0840*
H10	0.72300	-0.39410	0.62000	0.0900*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0860 (16)	0.0923 (16)	0.1095 (16)	0.0289 (14)	-0.0421 (15)	-0.0148 (14)
F2	0.157 (3)	0.0540 (11)	0.0894 (14)	-0.0013 (15)	0.0206 (18)	-0.0154 (10)
O1	0.0848 (18)	0.0700 (15)	0.0650 (12)	0.0173 (14)	-0.0112 (13)	-0.0159 (11)
N1	0.0499 (13)	0.0594 (14)	0.0656 (14)	0.0022 (13)	-0.0011 (13)	-0.0030 (12)
N2	0.0575 (17)	0.0571 (16)	0.103 (2)	0.0034 (13)	-0.0163 (18)	0.0053 (15)
N3	0.0719 (19)	0.0660 (18)	0.0770 (18)	-0.0127 (16)	0.0007 (16)	-0.0090 (14)
C1	0.0462 (15)	0.0494 (14)	0.0517 (13)	0.0050 (12)	0.0040 (12)	-0.0018 (12)
C2	0.0571 (18)	0.0657 (19)	0.0616 (17)	0.0112 (16)	-0.0070 (15)	-0.0052 (15)
C3	0.072 (2)	0.075 (2)	0.0650 (18)	-0.0056 (19)	-0.0057 (19)	-0.0142 (17)
C4	0.091 (3)	0.0569 (17)	0.0578 (16)	0.001 (2)	0.016 (2)	-0.0059 (14)
C5	0.085 (3)	0.0578 (18)	0.0682 (19)	0.0215 (19)	0.007 (2)	0.0092 (15)
C6	0.0638 (19)	0.0613 (17)	0.0590 (16)	0.0096 (17)	-0.0086 (16)	-0.0008 (14)
C7	0.0481 (15)	0.0521 (15)	0.0581 (15)	0.0081 (14)	-0.0015 (14)	-0.0054 (13)
C8	0.0637 (19)	0.0569 (17)	0.076 (2)	0.0040 (17)	0.0088 (18)	-0.0080 (15)
C9	0.0487 (17)	0.080 (2)	0.081 (2)	-0.0026 (18)	-0.0050 (17)	-0.0055 (19)
C10	0.070 (2)	0.0563 (19)	0.097 (3)	0.0018 (18)	-0.008 (2)	0.0063 (18)

Geometric parameters (\AA , $^\circ$)

F1—C2	1.356 (4)	C2—C3	1.374 (5)
F2—C4	1.357 (4)	C3—C4	1.354 (6)
O1—C7	1.400 (4)	C4—C5	1.358 (5)
O1—H1	0.8200	C5—C6	1.385 (5)
N1—C8	1.451 (5)	C7—C8	1.514 (5)
N1—C9	1.320 (5)	C3—H3	0.9300
N1—N2	1.347 (4)	C5—H5	0.9300
N2—C10	1.306 (5)	C6—H6	0.9300
N3—C10	1.338 (5)	C7—H7	0.9800
N3—C9	1.310 (5)	C8—H8A	0.9700
C1—C2	1.367 (4)	C8—H8B	0.9700
C1—C7	1.519 (4)	C9—H9	0.9300
C1—C6	1.377 (5)	C10—H10	0.9300
C7—O1—H1	109.00	N1—C8—C7	112.5 (3)
N2—N1—C8	121.2 (3)	N1—C9—N3	111.2 (3)
C8—N1—C9	129.7 (3)	N2—C10—N3	115.3 (4)
N2—N1—C9	109.1 (3)	C2—C3—H3	122.00
N1—N2—C10	102.4 (3)	C4—C3—H3	122.00
C9—N3—C10	102.1 (3)	C4—C5—H5	121.00
C2—C1—C7	121.5 (3)	C6—C5—H5	121.00

C6—C1—C7	122.1 (3)	C1—C6—H6	119.00
C2—C1—C6	116.3 (3)	C5—C6—H6	119.00
F1—C2—C3	117.3 (3)	O1—C7—H7	109.00
C1—C2—C3	124.5 (3)	C1—C7—H7	109.00
F1—C2—C1	118.3 (3)	C8—C7—H7	109.00
C2—C3—C4	116.3 (3)	N1—C8—H8A	109.00
F2—C4—C3	118.3 (3)	N1—C8—H8B	109.00
C3—C4—C5	123.0 (4)	C7—C8—H8A	109.00
F2—C4—C5	118.7 (3)	C7—C8—H8B	109.00
C4—C5—C6	118.5 (4)	H8A—C8—H8B	108.00
C1—C6—C5	121.4 (3)	N1—C9—H9	124.00
O1—C7—C8	110.8 (3)	N3—C9—H9	124.00
C1—C7—C8	108.0 (2)	N2—C10—H10	122.00
O1—C7—C1	110.3 (2)	N3—C10—H10	122.00
C8—N1—N2—C10	179.9 (3)	C7—C1—C6—C5	178.8 (3)
C9—N1—N2—C10	0.1 (4)	C2—C1—C7—O1	-154.7 (3)
N2—N1—C8—C7	73.8 (4)	C2—C1—C7—C8	84.1 (4)
C9—N1—C8—C7	-106.5 (4)	C6—C1—C7—O1	27.9 (4)
N2—N1—C9—N3	-0.1 (4)	C6—C1—C7—C8	-93.3 (3)
C8—N1—C9—N3	-179.9 (3)	F1—C2—C3—C4	-178.8 (3)
N1—N2—C10—N3	-0.1 (5)	C1—C2—C3—C4	1.3 (5)
C10—N3—C9—N1	0.0 (4)	C2—C3—C4—F2	-179.6 (3)
C9—N3—C10—N2	0.1 (5)	C2—C3—C4—C5	0.0 (6)
C6—C1—C2—F1	178.3 (3)	F2—C4—C5—C6	179.0 (3)
C6—C1—C2—C3	-1.9 (5)	C3—C4—C5—C6	-0.6 (6)
C7—C1—C2—F1	0.7 (4)	C4—C5—C6—C1	0.0 (5)
C7—C1—C2—C3	-179.5 (3)	O1—C7—C8—N1	62.4 (3)
C2—C1—C6—C5	1.2 (5)	C1—C7—C8—N1	-176.7 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots N3 ⁱ	0.82	2.01	2.826 (4)	171
C3—H3 \cdots N2 ⁱⁱ	0.93	2.62	3.500 (5)	158
C8—H8B \cdots F1 ⁱⁱⁱ	0.97	2.45	3.340 (5)	153
C10—H10 \cdots F2 ^{iv}	0.93	2.48	3.270 (4)	142

Symmetry codes: (i) $x+1/2, -y-1/2, -z+1$; (ii) $-x+2, y+1/2, -z+3/2$; (iii) $x-1, y, z$; (iv) $x, y-1, z$.