

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Benzyl-7-(2,4-dichlorophenyl)-4*H*-1,3,4-thiadiazolo[2,3-*c*][1,2,4]triazin-4-one

Hoong-Kun Fun,^{a,*‡} Suhana Arshad,^a B. K. Sarojini,^b
U. A. Imran^b and B.G. Krishna^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Chemistry, P. A. College of Engineering, Mangalore 574 153, India
Correspondence e-mail: hkfun@usm.my


Received 24 April 2012; accepted 10 May 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; disorder in main residue; R factor = 0.060; wR factor = 0.129; data-to-parameter ratio = 14.5.

In the title compound, $\text{C}_{17}\text{H}_{10}\text{Cl}_2\text{N}_4\text{OS}$, the phenyl ring and the H atoms attached to the adjacent C atom are disordered over two positions, with refined site occupancies of 0.509 (8) and 0.491 (8). The planar 4*H*-1,3,4-thiadiazolo[2,3-*c*][1,2,4]-triazine ring system [maximum deviation = 0.048 (3) Å] forms dihedral angles of 76.9 (5), 74.9 (5) and 9.88 (12)°, respectively, with the major and minor parts of the disordered phenyl ring and with the dichloro-substituted benzene ring. In the crystal, pairs of $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, forming inversion dimers with an $R_2^2(18)$ graph-set motif. A short $\text{S}\cdots\text{N}$ contact of 2.801 (3) Å is observed between the dimers.

Related literature

For applications of thiadiazole derivatives, see: Kurtzer (1965); Sandstrom (1968); Eue & Tietz (1970); Holla *et al.* (1988, 1998). For a related structure, see: Zhang *et al.* (2011); Fun *et al.* (2011); Ma & Yang (2008); Yu *et al.* (2007); Jia *et al.* (2011). For the stability of the temperature controller used for data collection, see: Cosier & Glazer (1986). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



‡ Thomson Reuters ResearcherID: A-3561-2009.

Experimental

Crystal data

$\text{C}_{17}\text{H}_{10}\text{Cl}_2\text{N}_4\text{OS}$
 $M_r = 389.25$
Triclinic, $P\bar{1}$
 $a = 4.4961$ (1) Å
 $b = 13.4412$ (5) Å
 $c = 14.4588$ (5) Å
 $\alpha = 70.620$ (2)°
 $\beta = 85.956$ (2)°
 $\gamma = 83.063$ (2)°
 $V = 817.81$ (5) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.54$ mm⁻¹
 $T = 100$ K
 $0.21 \times 0.10 \times 0.09$ mm

Data collection

Bruker SMART APEXII CCD
area-detector diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)
 $T_{\min} = 0.894$, $T_{\max} = 0.955$
8906 measured reflections
4086 independent reflections
2479 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.053$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$
 $wR(F^2) = 0.129$
 $S = 1.04$
4086 reflections
281 parameters
180 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.42$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.38$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}2A\cdots\text{O}1^i$	0.95	2.36	3.226 (5)	151

Symmetry code: (i) $-x + 1, -y + 2, -z + 2$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and SA thank Universiti Sains Malaysia (USM) for the Research University Grant (No. 1001/PFIZIK/811160). SA thanks the Malaysian government and USM for the award of Academic Staff Training Scheme (ASTS). BKS gratefully acknowledges the Department of Atomic Energy (DAE)/BRNS, Government of India, for providing financial assistance in the BRNS Project (grant No. 2011/34/20-BRNS/0846).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5127).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2009). *SADABS*, *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
Eue, L. & Tietz, H. (1970). *Pflanzenschutz-Nachrichten Bayer*, **23**, 208–218.
Fun, H.-K., Chantrapromma, S., Chandrakantha, B., Isloor, A. M. & Shetty, P. (2011). *Acta Cryst.* **E67**, o205–o206.
Holla, B. S., Kalluraya, B. & Sridhar, K. R. (1988). *Rev. Roum. Chim.* **33**, 277–288.
Holla, B. S., Sarojini, B. K. & Gonsalves, R. (1998). *Il Farmaco*, **53**, 395–398.
Jia, W., Wang, Z., Jia, X., Zhang, J. & Wang, W. (2011). *Acta Cryst.* **E67**, o1093.

- Kurtzer, F. (1965). *Adv. Heterocycl. Chem.* **5**, 119–204.
Ma, W.-W. & Yang, M.-H. (2008). *Acta Cryst. E* **64**, m630.
Sandstrom, J. (1968). *Adv. Heterocycl. Chem.* **9**, 165–209.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
Yu, M., Zhang, K., Qian, K.-D., Zhang, L.-X. & Liu, Y.-Z. (2007). *Acta Cryst. E* **63**, o2551.
Zhang, J., He, Q., Jiang, Q., Mu, H. & Wan, R. (2011). *Acta Cryst. E* **67**, o2255.

supplementary materials

Acta Cryst. (2012). E68, o1736–o1737 [doi:10.1107/S1600536812021253]

3-Benzyl-7-(2,4-dichlorophenyl)-4*H*-1,3,4-thiadiazolo[2,3-*c*][1,2,4]triazin-4-one

Hoong-Kun Fun, Suhana Arshad, B. K. Sarojini, U. A. Imran and B.G. Krishna

Comment

Several thiadiazoles find important application in the field of medicine, agriculture and industry (Kurtzer, 1965; Sandstrom, 1968). Thiadiazoles were also used as effective herbicides and fungicides (Eue & Tietz, 1970). The antibacterial activity of some of the thiadiazolotriazinone derivatives were reported (Holla *et al.*, 1998). The presence of substituents like aryloxymethyl and anilinomethyl on thiadiazole moiety were known to enhance the biological activities (Holla *et al.*, 1988). The crystal structures of some of the thiadiazole derivatives have been reported (Zhang *et al.*, 2011; Fun *et al.*, 2011; Ma & Yang, 2008). The present work describes the synthesis and crystal structure of the title compound (C₁₇H₁₀Cl₂N₄O_S), 3-benzyl-7-(2,4-dichlorophenyl)-4*H*-[1,3,4]thiadiazolo[2,3-*c*][1,2,4]triazin-4-one. The compound was prepared by the reaction of 4-amino-6-benzyl-3-mercapto-1,2,4-triazin-5(4*H*)-one with 2,4 dichlorobenzoic acid.

The molecular structure is shown in Fig. 1. The benzene ring and the hydrogen atoms which are attached to atom C11 are disordered over two positions with refined site-occupancies of 0.509 (8): 0.491 (8) ratio. Bond lengths and angles are within normal ranges. The 4*H*-[1,3,4]thiadiazolo [2,3-*c*][1,2,4]triazine ring (S1/N1/N2/N3/N4/C7–C10; maximum deviation = 0.048 (3) Å at atom C9) forms dihedral angles of 76.9 (5), 74.9 (5) and 9.88 (12)° with the major and minor parts of the disordered benzene ring (C12–C17 & C12X–C17X) and the dichloro-substituted benzene ring (C1–C6), respectively.

In the crystal packing (Fig. 2), two neighbouring molecules are linked by intermolecular C2—H2A⋯O1 hydrogen bonds (Table 1) into inversion dimers, forming R₂²(18) graph-set motifs (Bernstein *et al.*, 1995). A short S1⋯N3 contact [2.801 (3) Å, symmetry code: $-x, 1 - y, 2 - z$] is observed between the dimers and comparable with those short contacts of the previously reported structures (Yu *et al.*, 2007; Jia *et al.*, 2011).

Experimental


A mixture of 4-amino-6-benzyl-3-mercapto-1,2,4-triazin-5(4*H*)-one (2.34 g, 0.01 mol), 2,4-dichlorobenzoic acid (1.91 g, 0.01 mol) and phosphorus oxychloride (10 ml) was refluxed on a water bath for about 5 h. Excess of phosphorus oxychloride was removed under reduced pressure. The reaction mixture was cooled and poured onto crushed ice. The resulting solid product was filtered, washed with sodium bicarbonate solution (2%), followed by distilled water. It was dried and recrystallized from aqueous dioxane. The single crystals were grown by slow evaporation from a mixture of ethanol and dichloromethane. (1:1) (*M.p.* = 473–475 K).

Refinement


The benzene ring and the hydrogen atoms which are attached to atom C11 are disordered over two positions with refined site-occupancies of 0.509 (8): 0.491 (8) ratio. All the H atoms were positioned geometrically (C—H = 0.95 and 0.99 Å) and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Similarity and rigid bond restraints were used in the final refinement. Two outliers (0 1 1) and (0 0 1) were omitted.

Computing details

Data collection: *APEX2* (Bruker, 2009); cell refinement: *S SAINT* (Bruker, 2009); data reduction: *S SAINT* (Bruker, 2009); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).


Figure 1

The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atom-numbering scheme. All disordered components are shown.


Figure 2

A packing diagram of the title compound viewed along the *a* axis. Dashed lines represent the intermolecular hydrogen bonds. Only major disordered component is shown.

3-Benzyl-7-(2,4-dichlorophenyl)-4H-1,3,4-thiadiazolo[2,3-c][1,2,4]triazin-4-one
Crystal data

$C_{17}H_{10}Cl_2N_4OS$
 $M_r = 389.25$
 Triclinic, $P\bar{1}$
 Hall symbol: $-P\ 1$
 $a = 4.4961\ (1)\ \text{\AA}$
 $b = 13.4412\ (5)\ \text{\AA}$
 $c = 14.4588\ (5)\ \text{\AA}$
 $\alpha = 70.620\ (2)^\circ$
 $\beta = 85.956\ (2)^\circ$
 $\gamma = 83.063\ (2)^\circ$
 $V = 817.81\ (5)\ \text{\AA}^3$

$Z = 2$
 $F(000) = 396$
 $D_x = 1.581\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 5482 reflections
 $\theta = 2.5\text{--}31.5^\circ$
 $\mu = 0.54\ \text{mm}^{-1}$
 $T = 100\ \text{K}$
 Block, yellow
 $0.21 \times 0.10 \times 0.09\ \text{mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	8906 measured reflections
Radiation source: fine-focus sealed tube	4086 independent reflections
Graphite monochromator	2479 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.053$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 28.5^\circ$, $\theta_{\text{min}} = 2.5^\circ$
$T_{\text{min}} = 0.894$, $T_{\text{max}} = 0.955$	$h = -6 \rightarrow 6$
	$k = -18 \rightarrow 18$
	$l = -18 \rightarrow 19$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.060$	H-atom parameters constrained
$wR(F^2) = 0.129$	$w = 1/[\sigma^2(F_o^2) + (0.0449P)^2 + 0.2058P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
4086 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
281 parameters	$\Delta\rho_{\text{max}} = 0.42 \text{ e } \text{\AA}^{-3}$
180 restraints	$\Delta\rho_{\text{min}} = -0.38 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	-0.1885 (2)	1.03193 (7)	1.30609 (7)	0.0514 (3)	
C12	-0.2647 (2)	0.66171 (7)	1.25574 (6)	0.0410 (3)	
S1	0.00839 (17)	0.62886 (7)	1.07446 (5)	0.0301 (2)	
O1	0.7716 (5)	0.79090 (18)	0.84041 (16)	0.0380 (6)	
N1	0.3542 (6)	0.7790 (2)	0.99367 (18)	0.0301 (6)	
N2	0.3951 (5)	0.7091 (2)	0.94181 (18)	0.0284 (6)	
N3	0.2316 (5)	0.5525 (2)	0.92944 (17)	0.0321 (7)	
N4	0.4208 (5)	0.5638 (2)	0.84780 (18)	0.0330 (7)	
C1	0.1722 (7)	0.9123 (3)	1.1014 (2)	0.0338 (8)	
H1A	0.3000	0.9339	1.0447	0.041*	
C2	0.0985 (8)	0.9801 (3)	1.1546 (2)	0.0376 (8)	
H2A	0.1744	1.0473	1.1351	0.045*	
C3	-0.0885 (7)	0.9488 (3)	1.2373 (2)	0.0360 (8)	
C4	-0.1976 (7)	0.8511 (3)	1.2661 (2)	0.0354 (8)	
H4A	-0.3251	0.8301	1.3230	0.043*	

C5	-0.1205 (7)	0.7840 (3)	1.2119 (2)	0.0296 (7)	
C6	0.0653 (7)	0.8122 (3)	1.1278 (2)	0.0292 (7)	
C7	0.1560 (7)	0.7466 (3)	1.0647 (2)	0.0281 (7)	
C8	0.2259 (7)	0.6250 (3)	0.9714 (2)	0.0294 (7)	
C9	0.6039 (7)	0.7215 (3)	0.8630 (2)	0.0309 (8)	
C10	0.5922 (7)	0.6403 (3)	0.8173 (2)	0.0313 (8)	
C11	0.7890 (7)	0.6472 (3)	0.7261 (2)	0.0371 (9)	
H11A	0.9216	0.5801	0.7374	0.045*	0.509 (8)
H11B	0.9176	0.7053	0.7140	0.045*	0.509 (8)
H11C	0.9466	0.6944	0.7210	0.045*	0.491 (8)
H11D	0.8881	0.5759	0.7305	0.045*	0.491 (8)
C12	0.600 (6)	0.6676 (16)	0.6331 (18)	0.020 (3)	0.509 (8)
C13	0.4824 (18)	0.7651 (7)	0.5769 (5)	0.0307 (18)	0.509 (8)
H13A	0.5246	0.8266	0.5899	0.037*	0.509 (8)
C14	0.302 (4)	0.7744 (17)	0.5012 (10)	0.044 (3)	0.509 (8)
H14A	0.2146	0.8423	0.4631	0.053*	0.509 (8)
C15	0.248 (3)	0.6858 (12)	0.4802 (8)	0.049 (3)	0.509 (8)
H15A	0.1266	0.6927	0.4268	0.059*	0.509 (8)
C16	0.3718 (17)	0.5854 (7)	0.5380 (5)	0.042 (2)	0.509 (8)
H16A	0.3364	0.5239	0.5237	0.051*	0.509 (8)
C17	0.546 (2)	0.5769 (7)	0.6159 (7)	0.034 (2)	0.509 (8)
H17A	0.6267	0.5093	0.6569	0.041*	0.509 (8)
C12X	0.611 (7)	0.6874 (18)	0.6421 (19)	0.023 (3)	0.491 (8)
C13X	0.5639 (18)	0.7976 (7)	0.5985 (6)	0.0300 (18)	0.491 (8)
H13B	0.6560	0.8415	0.6253	0.036*	0.491 (8)
C14X	0.3859 (17)	0.8449 (7)	0.5171 (5)	0.041 (2)	0.491 (8)
H14B	0.3532	0.9199	0.4898	0.050*	0.491 (8)
C15X	0.256 (4)	0.7806 (18)	0.4761 (10)	0.042 (3)	0.491 (8)
H15B	0.1343	0.8127	0.4208	0.050*	0.491 (8)
C16X	0.302 (3)	0.6708 (13)	0.5142 (7)	0.041 (2)	0.491 (8)
H16B	0.2120	0.6277	0.4861	0.049*	0.491 (8)
C17X	0.483 (2)	0.6257 (8)	0.5948 (7)	0.033 (2)	0.491 (8)
H17B	0.5238	0.5507	0.6193	0.040*	0.491 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0638 (7)	0.0300 (5)	0.0509 (6)	0.0054 (5)	0.0255 (5)	-0.0094 (4)
C12	0.0471 (5)	0.0458 (6)	0.0308 (5)	-0.0175 (4)	0.0167 (4)	-0.0126 (4)
S1	0.0230 (4)	0.0477 (5)	0.0195 (4)	-0.0078 (4)	0.0029 (3)	-0.0101 (4)
O1	0.0324 (13)	0.0363 (14)	0.0368 (13)	-0.0029 (11)	0.0121 (10)	-0.0035 (11)
N1	0.0250 (14)	0.0336 (16)	0.0261 (14)	0.0025 (12)	0.0024 (11)	-0.0049 (12)
N2	0.0209 (13)	0.0383 (16)	0.0228 (13)	0.0010 (12)	0.0022 (11)	-0.0076 (12)
N3	0.0220 (14)	0.059 (2)	0.0178 (13)	-0.0098 (14)	0.0029 (10)	-0.0146 (13)
N4	0.0189 (13)	0.065 (2)	0.0189 (13)	-0.0113 (14)	0.0041 (10)	-0.0174 (13)
C1	0.0279 (18)	0.0290 (19)	0.0327 (18)	0.0058 (15)	0.0077 (14)	0.0008 (15)
C2	0.038 (2)	0.0248 (18)	0.0365 (19)	0.0046 (16)	0.0101 (16)	0.0027 (15)
C3	0.0323 (19)	0.0298 (19)	0.038 (2)	0.0105 (15)	0.0043 (15)	-0.0056 (15)
C4	0.0338 (19)	0.034 (2)	0.0289 (18)	0.0028 (16)	0.0108 (14)	-0.0020 (15)

C5	0.0270 (17)	0.0313 (18)	0.0253 (16)	-0.0021 (15)	0.0027 (13)	-0.0031 (14)
C6	0.0231 (17)	0.0342 (19)	0.0235 (16)	0.0048 (14)	0.0010 (13)	-0.0036 (14)
C7	0.0210 (16)	0.0343 (19)	0.0219 (16)	0.0001 (14)	-0.0002 (12)	-0.0007 (13)
C8	0.0194 (16)	0.050 (2)	0.0169 (15)	-0.0035 (15)	-0.0006 (12)	-0.0079 (14)
C9	0.0232 (17)	0.040 (2)	0.0215 (16)	0.0038 (16)	0.0003 (13)	-0.0014 (14)
C10	0.0175 (15)	0.054 (2)	0.0195 (15)	-0.0005 (16)	-0.0001 (12)	-0.0087 (15)
C11	0.0227 (17)	0.064 (3)	0.0236 (17)	-0.0066 (17)	0.0083 (13)	-0.0135 (17)
C12	0.019 (5)	0.034 (7)	0.011 (4)	0.005 (5)	0.002 (3)	-0.013 (4)
C13	0.038 (4)	0.035 (4)	0.018 (3)	0.000 (3)	0.008 (3)	-0.010 (3)
C14	0.033 (6)	0.064 (5)	0.020 (6)	0.013 (5)	-0.002 (4)	0.000 (6)
C15	0.033 (5)	0.080 (7)	0.025 (5)	-0.002 (5)	-0.005 (4)	-0.007 (5)
C16	0.048 (4)	0.058 (5)	0.029 (4)	-0.012 (4)	0.013 (3)	-0.027 (4)
C17	0.041 (5)	0.035 (5)	0.024 (4)	-0.004 (4)	0.007 (3)	-0.006 (4)
C12X	0.019 (4)	0.035 (7)	0.016 (6)	-0.008 (5)	0.008 (4)	-0.011 (4)
C13X	0.028 (4)	0.036 (4)	0.022 (4)	0.003 (3)	0.006 (3)	-0.008 (3)
C14X	0.039 (4)	0.051 (5)	0.022 (3)	0.005 (4)	0.009 (3)	0.000 (3)
C15X	0.019 (5)	0.083 (7)	0.014 (6)	0.009 (5)	0.001 (4)	-0.010 (6)
C16X	0.024 (5)	0.074 (6)	0.023 (5)	-0.004 (5)	0.003 (4)	-0.015 (5)
C17X	0.039 (5)	0.042 (5)	0.019 (4)	-0.005 (4)	0.011 (3)	-0.012 (4)

Geometric parameters (Å, °)

C11—C3	1.730 (4)	C11—H11A	0.9900
C12—C5	1.741 (3)	C11—H11B	0.9900
S1—C8	1.736 (3)	C11—H11C	0.9900
S1—C7	1.749 (3)	C11—H11D	0.9900
O1—C9	1.216 (4)	C12—C13	1.36 (2)
N1—C7	1.306 (4)	C12—C17	1.375 (18)
N1—N2	1.373 (4)	C13—C14	1.370 (14)
N2—C8	1.371 (4)	C13—H13A	0.9500
N2—C9	1.403 (4)	C14—C15	1.37 (3)
N3—C8	1.305 (4)	C14—H14A	0.9500
N3—N4	1.382 (3)	C15—C16	1.402 (16)
N4—C10	1.299 (4)	C15—H15A	0.9500
C1—C2	1.374 (5)	C16—C17	1.382 (12)
C1—C6	1.406 (4)	C16—H16A	0.9500
C1—H1A	0.9500	C17—H17A	0.9500
C2—C3	1.387 (4)	C12X—C13X	1.40 (2)
C2—H2A	0.9500	C12X—C17X	1.425 (19)
C3—C4	1.379 (5)	C13X—C14X	1.395 (10)
C4—C5	1.380 (4)	C13X—H13B	0.9500
C4—H4A	0.9500	C14X—C15X	1.40 (2)
C5—C6	1.395 (4)	C14X—H14B	0.9500
C6—C7	1.474 (4)	C15X—C16X	1.39 (3)
C9—C10	1.458 (5)	C15X—H15B	0.9500
C10—C11	1.518 (4)	C16X—C17X	1.391 (14)
C11—C12X	1.42 (3)	C16X—H16B	0.9500
C11—C12	1.57 (2)	C17X—H17B	0.9500
C8—S1—C7	88.09 (16)	H11A—C11—H11B	107.9

C7—N1—N2	109.0 (3)	C12X—C11—H11C	109.8
C8—N2—N1	117.1 (2)	C10—C11—H11C	109.8
C8—N2—C9	121.1 (3)	C12X—C11—H11D	109.8
N1—N2—C9	121.9 (3)	C10—C11—H11D	109.8
C8—N3—N4	116.4 (3)	H11C—C11—H11D	108.2
C10—N4—N3	121.4 (3)	C13—C12—C17	121.7 (16)
C2—C1—C6	122.4 (3)	C13—C12—C11	124.0 (12)
C2—C1—H1A	118.8	C17—C12—C11	114.2 (16)
C6—C1—H1A	118.8	C12—C13—C14	119.8 (13)
C1—C2—C3	118.8 (3)	C12—C13—H13A	120.1
C1—C2—H2A	120.6	C14—C13—H13A	120.1
C3—C2—H2A	120.6	C13—C14—C15	120.3 (15)
C4—C3—C2	120.7 (3)	C13—C14—H14A	119.9
C4—C3—C11	118.9 (3)	C15—C14—H14A	119.9
C2—C3—C11	120.4 (3)	C14—C15—C16	119.7 (10)
C3—C4—C5	119.6 (3)	C14—C15—H15A	120.1
C3—C4—H4A	120.2	C16—C15—H15A	120.1
C5—C4—H4A	120.2	C17—C16—C15	119.4 (9)
C4—C5—C6	121.8 (3)	C17—C16—H16A	120.3
C4—C5—C12	115.9 (2)	C15—C16—H16A	120.3
C6—C5—C12	122.2 (3)	C12—C17—C16	119.1 (12)
C5—C6—C1	116.6 (3)	C12—C17—H17A	120.5
C5—C6—C7	125.7 (3)	C16—C17—H17A	120.5
C1—C6—C7	117.7 (3)	C13X—C12X—C11	117.8 (13)
N1—C7—C6	118.7 (3)	C13X—C12X—C17X	116.3 (19)
N1—C7—S1	116.4 (2)	C11—C12X—C17X	125.9 (17)
C6—C7—S1	124.9 (2)	C14X—C13X—C12X	122.0 (12)
N3—C8—N2	125.2 (3)	C14X—C13X—H13B	119.0
N3—C8—S1	125.4 (2)	C12X—C13X—H13B	119.0
N2—C8—S1	109.4 (2)	C13X—C14X—C15X	119.1 (10)
O1—C9—N2	121.9 (3)	C13X—C14X—H14B	120.4
O1—C9—C10	127.1 (3)	C15X—C14X—H14B	120.4
N2—C9—C10	111.0 (3)	C16X—C15X—C14X	121.5 (11)
N4—C10—C9	124.8 (3)	C16X—C15X—H15B	119.3
N4—C10—C11	117.6 (3)	C14X—C15X—H15B	119.3
C9—C10—C11	117.6 (3)	C15X—C16X—C17X	118.2 (13)
C12X—C11—C10	109.5 (13)	C15X—C16X—H16B	120.9
C10—C11—C12	112.3 (11)	C17X—C16X—H16B	120.9
C10—C11—H11A	109.2	C16X—C17X—C12X	122.7 (14)
C12—C11—H11A	109.2	C16X—C17X—H17B	118.7
C10—C11—H11B	109.2	C12X—C17X—H17B	118.7
C12—C11—H11B	109.2		
C7—N1—N2—C8	1.3 (4)	N1—N2—C9—O1	4.5 (5)
C7—N1—N2—C9	-177.7 (3)	C8—N2—C9—C10	4.1 (4)
C8—N3—N4—C10	2.7 (4)	N1—N2—C9—C10	-177.0 (3)
C6—C1—C2—C3	-0.1 (5)	N3—N4—C10—C9	-0.7 (5)
C1—C2—C3—C4	0.4 (5)	N3—N4—C10—C11	-179.1 (3)
C1—C2—C3—C11	-179.7 (3)	O1—C9—C10—N4	175.7 (3)

C2—C3—C4—C5	-0.2 (5)	N2—C9—C10—N4	-2.7 (4)
C11—C3—C4—C5	179.8 (3)	O1—C9—C10—C11	-5.8 (5)
C3—C4—C5—C6	-0.2 (5)	N2—C9—C10—C11	175.7 (3)
C3—C4—C5—C12	178.9 (2)	N4—C10—C11—C12X	75.2 (9)
C4—C5—C6—C1	0.4 (5)	C9—C10—C11—C12X	-103.4 (9)
C12—C5—C6—C1	-178.6 (2)	N4—C10—C11—C12	62.9 (9)
C4—C5—C6—C7	-179.1 (3)	C9—C10—C11—C12	-115.7 (8)
C12—C5—C6—C7	1.9 (5)	C10—C11—C12—C13	84 (2)
C2—C1—C6—C5	-0.3 (5)	C10—C11—C12—C17	-92.4 (19)
C2—C1—C6—C7	179.3 (3)	C17—C12—C13—C14	0 (3)
N2—N1—C7—C6	-177.3 (3)	C11—C12—C13—C14	-175.2 (18)
N2—N1—C7—S1	0.6 (3)	C12—C13—C14—C15	-2 (3)
C5—C6—C7—N1	-173.5 (3)	C13—C14—C15—C16	1 (3)
C1—C6—C7—N1	7.0 (4)	C14—C15—C16—C17	0.4 (19)
C5—C6—C7—S1	8.8 (5)	C13—C12—C17—C16	1 (3)
C1—C6—C7—S1	-170.7 (2)	C11—C12—C17—C16	177.5 (13)
C8—S1—C7—N1	-1.7 (3)	C15—C16—C17—C12	-1.9 (19)
C8—S1—C7—C6	176.0 (3)	C10—C11—C12X—C13X	87 (2)
N4—N3—C8—N2	-1.2 (5)	C10—C11—C12X—C17X	-95 (2)
N4—N3—C8—S1	-180.0 (2)	C11—C12X—C13X—C14X	-178.3 (14)
N1—N2—C8—N3	178.5 (3)	C17X—C12X—C13X—C14X	4 (3)
C9—N2—C8—N3	-2.5 (5)	C12X—C13X—C14X—C15X	-1.6 (19)
N1—N2—C8—S1	-2.5 (3)	C13X—C14X—C15X—C16X	0 (2)
C9—N2—C8—S1	176.5 (2)	C14X—C15X—C16X—C17X	0 (2)
C7—S1—C8—N3	-178.9 (3)	C15X—C16X—C17X—C12X	3 (2)
C7—S1—C8—N2	2.2 (2)	C13X—C12X—C17X—C16X	-5 (3)
C8—N2—C9—O1	-174.4 (3)	C11—C12X—C17X—C16X	177.7 (18)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C2—H2 <i>A</i> ...O1 ⁱ	0.95	2.36	3.226 (5)	151

Symmetry code: (i) $-x+1, -y+2, -z+2$.