

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N,N'-Bis[(1*H*-imidazol-1-yl)methyl]-2,2'-(disulfanediyl)dianiline

Hon Man Lee, Chang-Chih Hsieh and Yih-Chern Horng*

National Changhua University of Education, Department of Chemistry, Changhua, Taiwan 50058

Correspondence e-mail: ychorng@cc.ncue.edu.tw


Received 1 June 2012; accepted 4 June 2012

 Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.033; wR factor = 0.128; data-to-parameter ratio = 19.0.

The symmetrical title compound, $\text{C}_{20}\text{H}_{20}\text{N}_6\text{S}_2$, contains a disulfide bond of 2.0884 (6) Å. The C—S—S—C torsion angle is -59.57 (7)°. In the crystal, classical N—H···N and non-classical C—H···N hydrogen bonds link the compounds into chains along the a axis.

Related literature

For transition metal complexes having related ligands, see: Hsieh *et al.* (2009*a,b*).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{20}\text{N}_6\text{S}_2$
 $M_r = 408.56$
 Monoclinic, $P2_1/c$

$a = 11.2009$ (10) Å
 $b = 11.6067$ (10) Å
 $c = 15.5230$ (14) Å

$\beta = 97.162$ (2)°
 $V = 2002.3$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.28$ mm⁻¹
 $T = 150$ K
 $0.38 \times 0.32 \times 0.24$ mm

Data collection

Bruker SMART APEXII diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2003)
 $T_{\min} = 0.888$, $T_{\max} = 0.954$

23330 measured reflections
 4970 independent reflections
 3844 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.128$
 $S = 0.96$
 4970 reflections
 261 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.27$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N6}-\text{H22}\cdots\text{N4}^{\text{i}}$	0.818 (17)	2.252 (17)	2.9825 (19)	148.9 (15)
$\text{C1}-\text{H1}\cdots\text{N2}^{\text{ii}}$	0.93	2.58	3.448 (2)	155

Symmetry codes: (i) $-x, -y, -z + 2$; (ii) $-x + 1, -y, -z + 2$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: DIAMOND (Brandenburg, 2006).

We thank the National Science Council of Taiwan for financial support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2478).

References

- Brandenburg, K. (2006). DIAMOND. Crystal Impact GbR, Bonn, Germany.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Hsieh, C.-C., Chao, W.-J. & Horng, Y.-C. (2009*a*). *Inorg. Chem. Commun.* **12**, 778–781.
 Hsieh, C.-C., Chao, W.-J., Horng, Y.-C. & Lee, H. M. (2009*b*). *J. Chin. Chem. Soc.* **56**, 435–442.
 Sheldrick, G. M. (2003). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2012). E68, o2051 [doi:10.1107/S1600536812025330]

N,N'*-Bis[(1*H*-imidazol-1-yl)methyl]-2,2'-(disulfaneyl)dianiline*Hon Man Lee, Chang-Chih Hsieh and Yih-Chern Horng****Comment**

In our research, we have focused on the synthesis of low-molecular-weight complexes, which exhibited the structural and functional features of the active sites in particular enzymes, such as Ni-containing methyl coenzyme *M* reductase (Hsieh *et al.*, 2009*b*) and Fe-containing nitrile hydratase (Hsieh *et al.*, 2009*a*). The synthesis and crystal structure determination of the title compound are reported here. This compound will be used as a coordination ligand in related studies.

The disulfide bond length is 2.0884 (6) Å and the C—S—S—C torsion angle is 59.56 (7)° (Fig. 1). In the crystal structure the compound does not exhibit crystallographic symmetry. Classical N—H···N and non-classical C—H···N hydrogen bonds (Table 1) link the compounds into one-dimensional chains along the *a*-axis (Fig. 2).

Experimental

Pyrazole (1.0 g, 14.7 mmol) was dissolved in 50 ml CH₂Cl₂ and treated with formaldehyde (37%, 1.19 g, 14.7 mmol). After stirring at room temperature for 10 min, the mixture was added to a batch of 2-aminophenyl disulfide (1.82 g, 7.32 mmol) and stirred for a further 12 h. After completion of the reaction, the solution was extracted with distilled water. Portions of the CH₂Cl₂ extract were collected and dried using anhydrous MgSO₄. The solvent was then removed under vacuum to afford a yellow powder (2.56 g, 86%). Crystals suitable for X-ray diffraction analysis were obtained by slow evaporation of a tetrahydrofuran solution of the compound at -4 °C.

Refinement

C-bound H atoms were positioned geometrically and refined as riding, with C_{aryl}—H = 0.93 and C_{methylene}—H = 0.97 Å; U_{iso}(H) = 1.2U_{eq}(C). N-bound H atoms were located in a difference Fourier map and freely refined (N5—H21 = 0.833 (17) Å and N6—H22 = 0.818 (17) Å).

Computing details

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINTE* (Bruker, 2007); data reduction: *SAINTE* (Bruker, 2007); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *DIAMOND* (Brandenburg, 2006).





Figure 1

The structure of the title compound, showing 50% probability displacement ellipsoids for the non-hydrogen atoms. The H atoms are depicted as spheres of arbitrary radius.

**Figure 2**

A view of the one-dimensional hydrogen-bonded chain, showing the hydrogen bonds as dashed lines.

N,N'-Bis[(1*H*-imidazol-1-yl)methyl]-2,2'-(disulfanediyl)dianiline

Crystal data

$C_{20}H_{20}N_6S_2$

$M_r = 408.56$

Monoclinic, $P2_1/c$

$a = 11.2009$ (10) Å

$b = 11.6067$ (10) Å

$c = 15.5230$ (14) Å

$\beta = 97.162$ (2)°

$V = 2002.3$ (3) Å³

$Z = 4$

$F(000) = 840$

$D_x = 1.355$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6226 reflections

$\theta = 2.2$ – 27.2 °

$\mu = 0.28$ mm⁻¹

$T = 150$ K

Block, yellow

$0.38 \times 0.32 \times 0.24$ mm

Data collection

Bruker SMART APEXII

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 2003)

$T_{\min} = 0.888$, $T_{\max} = 0.954$

23330 measured reflections

4970 independent reflections

3844 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 28.3$ °, $\theta_{\min} = 1.8$ °

$h = -14 \rightarrow 14$

$k = -15 \rightarrow 15$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.033$

$wR(F^2) = 0.128$

$S = 0.96$

4970 reflections

261 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 0.0208P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.27$ e Å⁻³

$\Delta\rho_{\min} = -0.26$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	-0.02094 (3)	0.08013 (3)	0.76606 (3)	0.03264 (13)
S2	0.01131 (3)	0.25438 (3)	0.79328 (3)	0.03404 (13)
N3	0.20015 (10)	0.01306 (11)	1.07381 (8)	0.0302 (3)
N1	0.36181 (11)	-0.17778 (11)	0.86130 (8)	0.0317 (3)
N4	0.14835 (12)	-0.08760 (11)	1.04385 (10)	0.0390 (3)
N2	0.47440 (12)	-0.20760 (14)	0.84774 (10)	0.0460 (4)
N6	0.10991 (11)	0.15689 (12)	0.97273 (8)	0.0338 (3)
N5	0.21637 (11)	-0.02980 (11)	0.82527 (8)	0.0312 (3)
C7	0.33157 (12)	-0.05670 (13)	0.87099 (10)	0.0309 (3)
H7A	0.3316	-0.0392	0.9321	0.037*
H7B	0.3923	-0.0091	0.8491	0.037*
C8	0.13251 (13)	0.12062 (13)	1.06103 (10)	0.0321 (3)
H8A	0.0561	0.1112	1.0836	0.039*
H8B	0.1768	0.1808	1.0946	0.039*
C1	0.30946 (13)	-0.00488 (16)	1.12009 (10)	0.0384 (4)
H1	0.3605	0.0512	1.1469	0.046*
C3	0.22890 (16)	-0.16789 (15)	1.07264 (12)	0.0446 (4)
H3	0.2182	-0.2463	1.0622	0.053*
C2	0.33048 (15)	-0.12061 (16)	1.12002 (11)	0.0455 (4)
H2	0.3982	-0.1594	1.1461	0.055*
C4	0.29052 (19)	-0.26996 (15)	0.86760 (14)	0.0490 (4)
H4	0.2098	-0.2687	0.8762	0.059*
C5	0.3583 (2)	-0.36500 (17)	0.85913 (13)	0.0607 (6)
H5	0.3348	-0.4417	0.8613	0.073*
C6	0.4712 (2)	-0.32236 (18)	0.84645 (13)	0.0595 (6)
H6	0.5365	-0.3686	0.8381	0.071*
C13	0.16083 (13)	0.25746 (12)	0.84587 (10)	0.0288 (3)
C15	0.19876 (12)	-0.00747 (12)	0.73713 (9)	0.0277 (3)
C14	0.19334 (12)	0.21299 (12)	0.93032 (9)	0.0271 (3)
C16	0.09108 (13)	0.04528 (12)	0.69947 (10)	0.0304 (3)
C9	0.31319 (13)	0.22646 (13)	0.96785 (10)	0.0317 (3)
H9	0.3368	0.1982	1.0234	0.038*
C10	0.39688 (14)	0.28116 (14)	0.92343 (11)	0.0372 (4)
H10	0.4754	0.2902	0.9500	0.045*
C11	0.36564 (16)	0.32242 (15)	0.84051 (12)	0.0429 (4)
H11	0.4225	0.3578	0.8105	0.052*
C20	0.28490 (14)	-0.03762 (13)	0.68293 (10)	0.0342 (3)

H20	0.3558	-0.0736	0.7063	0.041*
C17	0.07364 (16)	0.06746 (13)	0.61034 (11)	0.0389 (4)
H17	0.0026	0.1023	0.5859	0.047*
C18	0.15999 (17)	0.03861 (14)	0.55792 (11)	0.0433 (4)
H18	0.1478	0.0544	0.4987	0.052*
C19	0.26499 (16)	-0.01419 (14)	0.59480 (11)	0.0415 (4)
H19	0.3232	-0.0343	0.5597	0.050*
C12	0.24702 (15)	0.31009 (14)	0.80253 (10)	0.0382 (4)
H12	0.2251	0.3379	0.7466	0.046*
H22	0.0410 (15)	0.1489 (14)	0.9491 (11)	0.029 (4)*
H21	0.1641 (15)	-0.0085 (15)	0.8556 (11)	0.036 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.02746 (19)	0.0298 (2)	0.0386 (2)	0.00133 (13)	-0.00418 (14)	0.00020 (15)
S2	0.0400 (2)	0.0265 (2)	0.0330 (2)	0.00832 (14)	-0.00588 (16)	0.00215 (14)
N3	0.0269 (6)	0.0364 (7)	0.0274 (6)	0.0039 (5)	0.0033 (5)	0.0034 (5)
N1	0.0289 (6)	0.0306 (7)	0.0343 (7)	0.0044 (5)	-0.0016 (5)	-0.0033 (5)
N4	0.0324 (7)	0.0366 (7)	0.0478 (8)	0.0018 (5)	0.0038 (6)	0.0016 (6)
N2	0.0330 (7)	0.0521 (9)	0.0504 (9)	0.0144 (6)	-0.0048 (6)	-0.0143 (7)
N6	0.0277 (6)	0.0436 (8)	0.0283 (7)	-0.0016 (5)	-0.0029 (5)	0.0089 (5)
N5	0.0277 (6)	0.0371 (7)	0.0282 (6)	0.0087 (5)	0.0018 (5)	0.0003 (5)
C7	0.0282 (7)	0.0285 (7)	0.0346 (8)	0.0019 (5)	-0.0018 (6)	-0.0058 (6)
C8	0.0311 (7)	0.0373 (8)	0.0278 (7)	0.0064 (6)	0.0034 (6)	0.0037 (6)
C1	0.0323 (7)	0.0538 (10)	0.0279 (8)	0.0103 (7)	-0.0009 (6)	0.0002 (7)
C3	0.0440 (9)	0.0375 (9)	0.0533 (11)	0.0108 (7)	0.0104 (8)	0.0060 (8)
C2	0.0422 (9)	0.0565 (11)	0.0370 (9)	0.0210 (8)	0.0023 (7)	0.0054 (8)
C4	0.0564 (11)	0.0355 (9)	0.0553 (12)	-0.0075 (8)	0.0083 (9)	-0.0083 (8)
C5	0.0962 (17)	0.0302 (9)	0.0521 (12)	0.0047 (10)	-0.0055 (11)	-0.0059 (8)
C6	0.0685 (13)	0.0504 (11)	0.0531 (11)	0.0337 (10)	-0.0175 (10)	-0.0168 (9)
C13	0.0360 (7)	0.0219 (7)	0.0275 (7)	0.0029 (5)	0.0001 (6)	-0.0020 (5)
C15	0.0316 (7)	0.0211 (7)	0.0296 (7)	-0.0043 (5)	0.0005 (5)	-0.0017 (5)
C14	0.0308 (7)	0.0230 (7)	0.0272 (7)	0.0027 (5)	0.0025 (5)	-0.0017 (5)
C16	0.0340 (7)	0.0235 (7)	0.0318 (8)	-0.0052 (5)	-0.0033 (6)	-0.0024 (6)
C9	0.0328 (7)	0.0318 (8)	0.0295 (8)	0.0011 (6)	0.0006 (6)	-0.0012 (6)
C10	0.0331 (8)	0.0364 (8)	0.0422 (9)	-0.0037 (6)	0.0050 (6)	-0.0063 (7)
C11	0.0450 (9)	0.0420 (10)	0.0435 (10)	-0.0103 (7)	0.0125 (7)	0.0004 (7)
C20	0.0358 (7)	0.0285 (8)	0.0385 (9)	-0.0057 (6)	0.0057 (6)	-0.0043 (6)
C17	0.0489 (9)	0.0294 (8)	0.0349 (9)	-0.0063 (7)	-0.0089 (7)	0.0011 (6)
C18	0.0642 (11)	0.0355 (9)	0.0290 (8)	-0.0150 (8)	0.0010 (7)	-0.0001 (7)
C19	0.0542 (10)	0.0348 (8)	0.0378 (9)	-0.0143 (7)	0.0146 (7)	-0.0071 (7)
C12	0.0510 (9)	0.0346 (8)	0.0289 (8)	-0.0041 (7)	0.0051 (7)	0.0033 (6)

Geometric parameters (\AA , $^\circ$)

S1—C16	1.7691 (16)	C4—C5	1.355 (3)
S1—S2	2.0884 (6)	C4—H4	0.9300
S2—C13	1.7697 (15)	C5—C6	1.394 (3)
N3—C1	1.3562 (18)	C5—H5	0.9300

N3—N4	1.3605 (18)	C6—H6	0.9300
N3—C8	1.4609 (18)	C13—C12	1.385 (2)
N1—C4	1.346 (2)	C13—C14	1.414 (2)
N1—N2	1.3493 (18)	C15—C20	1.401 (2)
N1—C7	1.4577 (19)	C15—C16	1.4127 (19)
N4—C3	1.335 (2)	C14—C9	1.4039 (19)
N2—C6	1.333 (3)	C16—C17	1.397 (2)
N6—C14	1.3727 (19)	C9—C10	1.385 (2)
N6—C8	1.4260 (19)	C9—H9	0.9300
N6—H22	0.818 (17)	C10—C11	1.377 (3)
N5—C15	1.3823 (18)	C10—H10	0.9300
N5—C7	1.4276 (17)	C11—C12	1.392 (2)
N5—H21	0.833 (17)	C11—H11	0.9300
C7—H7A	0.9700	C20—C19	1.385 (2)
C7—H7B	0.9700	C20—H20	0.9300
C8—H8A	0.9700	C17—C18	1.381 (3)
C8—H8B	0.9700	C17—H17	0.9300
C1—C2	1.364 (2)	C18—C19	1.385 (3)
C1—H1	0.9300	C18—H18	0.9300
C3—C2	1.389 (3)	C19—H19	0.9300
C3—H3	0.9300	C12—H12	0.9300
C2—H2	0.9300		
C16—S1—S2	102.84 (5)	C4—C5—H5	127.6
C13—S2—S1	104.05 (5)	C6—C5—H5	127.6
C1—N3—N4	111.56 (13)	N2—C6—C5	112.10 (16)
C1—N3—C8	128.45 (14)	N2—C6—H6	123.9
N4—N3—C8	119.71 (12)	C5—C6—H6	123.9
C4—N1—N2	112.46 (15)	C12—C13—C14	119.78 (14)
C4—N1—C7	127.68 (14)	C12—C13—S2	117.49 (12)
N2—N1—C7	119.81 (13)	C14—C13—S2	122.68 (11)
C3—N4—N3	104.09 (14)	N5—C15—C20	121.57 (13)
C6—N2—N1	103.56 (15)	N5—C15—C16	120.00 (13)
C14—N6—C8	123.45 (13)	C20—C15—C16	118.43 (14)
C14—N6—H22	120.0 (11)	N6—C14—C9	121.79 (13)
C8—N6—H22	116.2 (11)	N6—C14—C13	120.39 (13)
C15—N5—C7	122.80 (13)	C9—C14—C13	117.81 (13)
C15—N5—H21	118.7 (12)	C17—C16—C15	119.77 (14)
C7—N5—H21	116.1 (12)	C17—C16—S1	120.99 (12)
N5—C7—N1	111.52 (12)	C15—C16—S1	119.24 (11)
N5—C7—H7A	109.3	C10—C9—C14	121.06 (14)
N1—C7—H7A	109.3	C10—C9—H9	119.5
N5—C7—H7B	109.3	C14—C9—H9	119.5
N1—C7—H7B	109.3	C11—C10—C9	121.06 (15)
H7A—C7—H7B	108.0	C11—C10—H10	119.5
N6—C8—N3	114.18 (12)	C9—C10—H10	119.5
N6—C8—H8A	108.7	C10—C11—C12	118.55 (15)
N3—C8—H8A	108.7	C10—C11—H11	120.7
N6—C8—H8B	108.7	C12—C11—H11	120.7

N3—C8—H8B	108.7	C19—C20—C15	120.38 (15)
H8A—C8—H8B	107.6	C19—C20—H20	119.8
N3—C1—C2	107.18 (15)	C15—C20—H20	119.8
N3—C1—H1	126.4	C18—C17—C16	121.16 (16)
C2—C1—H1	126.4	C18—C17—H17	119.4
N4—C3—C2	112.12 (16)	C16—C17—H17	119.4
N4—C3—H3	123.9	C17—C18—C19	119.00 (15)
C2—C3—H3	123.9	C17—C18—H18	120.5
C1—C2—C3	105.04 (14)	C19—C18—H18	120.5
C1—C2—H2	127.5	C18—C19—C20	121.26 (16)
C3—C2—H2	127.5	C18—C19—H19	119.4
N1—C4—C5	107.17 (19)	C20—C19—H19	119.4
N1—C4—H4	126.4	C13—C12—C11	121.72 (15)
C5—C4—H4	126.4	C13—C12—H12	119.1
C4—C5—C6	104.70 (18)	C11—C12—H12	119.1
C16—S1—S2—C13	-59.57 (8)	C8—N6—C14—C13	-173.15 (14)
C1—N3—N4—C3	-0.24 (18)	C12—C13—C14—N6	-177.59 (14)
C8—N3—N4—C3	-174.71 (13)	S2—C13—C14—N6	5.04 (19)
C4—N1—N2—C6	-0.47 (19)	C12—C13—C14—C9	1.3 (2)
C7—N1—N2—C6	177.01 (14)	S2—C13—C14—C9	-176.10 (10)
C15—N5—C7—N1	-81.64 (17)	N5—C15—C16—C17	-179.88 (13)
C4—N1—C7—N5	-44.9 (2)	C20—C15—C16—C17	-0.9 (2)
N2—N1—C7—N5	138.08 (14)	N5—C15—C16—S1	-0.91 (19)
C14—N6—C8—N3	-80.33 (18)	C20—C15—C16—S1	178.09 (10)
C1—N3—C8—N6	118.43 (16)	S2—S1—C16—C17	-83.60 (12)
N4—N3—C8—N6	-68.13 (18)	S2—S1—C16—C15	97.44 (11)
N4—N3—C1—C2	0.44 (18)	N6—C14—C9—C10	178.62 (14)
C8—N3—C1—C2	174.31 (14)	C13—C14—C9—C10	-0.2 (2)
N3—N4—C3—C2	-0.06 (19)	C14—C9—C10—C11	-1.1 (2)
N3—C1—C2—C3	-0.44 (18)	C9—C10—C11—C12	1.3 (2)
N4—C3—C2—C1	0.3 (2)	N5—C15—C20—C19	-179.97 (14)
N2—N1—C4—C5	0.8 (2)	C16—C15—C20—C19	1.0 (2)
C7—N1—C4—C5	-176.40 (15)	C15—C16—C17—C18	0.1 (2)
N1—C4—C5—C6	-0.8 (2)	S1—C16—C17—C18	-178.89 (12)
N1—N2—C6—C5	-0.1 (2)	C16—C17—C18—C19	0.6 (2)
C4—C5—C6—N2	0.6 (2)	C17—C18—C19—C20	-0.4 (2)
S1—S2—C13—C12	112.56 (11)	C15—C20—C19—C18	-0.4 (2)
S1—S2—C13—C14	-70.02 (12)	C14—C13—C12—C11	-1.1 (2)
C7—N5—C15—C20	16.4 (2)	S2—C13—C12—C11	176.41 (13)
C7—N5—C15—C16	-164.62 (13)	C10—C11—C12—C13	-0.2 (3)
C8—N6—C14—C9	8.0 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N6—H22...N4 ⁱ	0.818 (17)	2.252 (17)	2.9825 (19)	148.9 (15)
C1—H1...N2 ⁱⁱ	0.93	2.58	3.448 (2)	155

Symmetry codes: (i) $-x, -y, -z+2$; (ii) $-x+1, -y, -z+2$.