

N'-(3-Fluorobenzylidene)-2-methylbenzohydrazide

Ying Song,^{a,b} Jian-Long Zhao,^b Jan-Gang Wang,^b Fei Lu,^a Hong Lu^{a*} and Shi-Peng Li^b

^aThe 1st Affiliated Hospital of Henan University of Science & Technology, Luoyang Henan 471003, People's Republic of China, and ^bMedical College, Henan University of Science & Technology, Luoyang Henan 471003, People's Republic of China

Correspondence e-mail: sying20096@126.com


Received 1 June 2012; accepted 5 June 2012

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.041; wR factor = 0.126; data-to-parameter ratio = 13.8.

The asymmetric unit of the title compound, $C_{15}H_{13}FN_2O$, contains two independent molecules with different conformations; the two aromatic rings in the independent molecules form dihedral angles of 85.3 (2) and 10.0 (2) $^\circ$. In the crystal, N—H···O hydrogen bonds link the molecules into chains along [100].

Related literature

For related structures, see: Xu *et al.* (2011); Wang *et al.* (2012); Hu & Liu (2012). For the biological activity of benzohydrazones, see: Zhang *et al.* (2012).



Experimental

Crystal data



$M_r = 256.27$

Triclinic, $P\bar{1}$	$V = 1326.9 (4)\text{ \AA}^3$
$a = 7.8516 (13)\text{ \AA}$	$Z = 4$
$b = 8.1466 (13)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 21.158 (3)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$\alpha = 86.668 (2)$ $^\circ$	$T = 298\text{ K}$
$\beta = 85.806 (2)$ $^\circ$	$0.17 \times 0.15 \times 0.15\text{ mm}$
$\gamma = 79.772 (2)$ $^\circ$	

Data collection

Bruker SMART CCD area-detector diffractometer	9780 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	4844 independent reflections
	4067 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.019$
	$T_{\min} = 0.985$, $T_{\max} = 0.986$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.126$	$\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
$S = 1.06$	$\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$
4844 reflections	
351 parameters	
2 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N4—H4···O1	0.89 (1)	1.93 (1)	2.8115 (15)	168 (2)
N2—H2···O2 ⁱ	0.89 (1)	2.08 (1)	2.9185 (16)	157 (2)

Symmetry code: (i) $x + 1, y, z$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5308).

References

- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hu, H.-N. & Liu, S.-Y. (2012). *Acta Cryst. E68*, o1613.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Wang, D.-Y., Meng, X.-F. & Ma, J.-J. (2012). *Acta Cryst. E68*, o21.
- Xu, C.-B., Wang, Z.-G., Nan, Y., Yuan, L., Wang, R. & Zhang, S.-X. (2011). *Acta Cryst. E67*, o70.
- Zhang, M., Xian, D.-M., Li, H.-H., Zhang, J.-C. & You, Z.-L. (2012). *Aust. J. Chem.* **65**, 343–350.

supplementary materials

Acta Cryst. (2012). E68, o2074 [doi:10.1107/S1600536812025548]

N'-(3-Fluorobenzylidene)-2-methylbenzohydrazide

Ying Song, Jian-Long Zhao, Jan-Gang Wang, Fei Lu, Hong Lu and Shi-Peng Li

Comment

Benzohydrazone derived from the condensation reactions of benzohydrazides with carbonyl-containing compounds have been proved to have antimicrobial and antitumor activities (Zhang *et al.*, 2012). Herewith we present the title compound (I), which is a new benzohydrazone derivative.

The asymmetric unit of the title compound contains two independent molecules, A and B, respectively, with different conformations (Fig. 1) - two aromatic rings in the independent molecules form the dihedral angles of 85.3 (2) $^{\circ}$ (B) and 10.0 (2) $^{\circ}$ (A), respectively. All the bond lengths and angles are normal and correspond to those observed in the related structures (Wang *et al.*, 2012; Hu & Liu, 2012; Xu *et al.*, 2011). In the crystal structure, intermolecular N—H \cdots O hydrogen bonds (Table 1) link the molecules into chains in [100] (Fig. 2).

Experimental


Equimolar quantities (1 mmol each) of 3-fluorohlorobenzaldehyde and 2-methylbenzohydrazide were mixed in 50 mL methanol. The materials were stirred at room temperature for 1 h to give a clear colorless solution. Colourless block-like single crystals were obtained after a few days.

Refinement


Atom H2 located from a Fourier map was isotropically refined, with N—H distance restrained to 0.90 (1) Å. C-bound H atoms were positioned geometrically and treated as riding on their parent atoms, with C—H distances of 0.93–0.96 Å, and with $U_{\text{iso}}(\text{H})$ set to 1.2–1.5 $U_{\text{eq}}(\text{C})$.

Computing details

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT* (Bruker, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

**Figure 1**

Conformations of two independent molecules in (I). Displacement ellipsoids are drawn with 30% probability level.

**Figure 2**

Molecular packing diagram of (I), viewed down the *b* axis. Hydrogen bonds are shown as thin dashed lines. C-bound H atoms omitted for clarity

N'-(3-Fluorobenzylidene)-2-methylbenzohydrazide*Crystal data*

$C_{15}H_{13}FN_2O$	$Z = 4$
$M_r = 256.27$	$F(000) = 536$
Triclinic, $P\bar{1}$	$D_x = 1.283 \text{ Mg m}^{-3}$
$a = 7.8516 (13) \text{ \AA}$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$b = 8.1466 (13) \text{ \AA}$	Cell parameters from 5461 reflections
$c = 21.158 (3) \text{ \AA}$	$\theta = 2.6\text{--}27.0^\circ$
$\alpha = 86.668 (2)^\circ$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 85.806 (2)^\circ$	$T = 298 \text{ K}$
$\gamma = 79.772 (2)^\circ$	Block, colourless
$V = 1326.9 (4) \text{ \AA}^3$	$0.17 \times 0.15 \times 0.15 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	9780 measured reflections
Radiation source: fine-focus sealed tube	4844 independent reflections
Graphite monochromator	4067 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.019$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 25.5^\circ, \theta_{\text{min}} = 2.6^\circ$
$T_{\text{min}} = 0.985, T_{\text{max}} = 0.986$	$h = -9 \rightarrow 9$
	$k = -9 \rightarrow 9$
	$l = -25 \rightarrow 25$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.041$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.126$	$w = 1/[\sigma^2(F_o^2) + (0.0719P)^2 + 0.1885P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\text{max}} = 0.001$
4844 reflections	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
351 parameters	$\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$
2 restraints	
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^* / U_{\text{eq}}$
F1	0.49294 (13)	-0.42660 (13)	0.33943 (6)	0.0741 (3)
F2	0.1795 (2)	-0.2449 (2)	0.48808 (6)	0.1094 (5)
N1	0.86796 (15)	0.01129 (14)	0.27473 (6)	0.0424 (3)

N2	0.95208 (15)	0.13387 (14)	0.24586 (6)	0.0432 (3)
N3	0.38019 (14)	0.05746 (15)	0.28693 (5)	0.0409 (3)
N4	0.46524 (15)	0.09877 (15)	0.23035 (5)	0.0419 (3)
O1	0.70015 (12)	0.31769 (12)	0.24076 (5)	0.0480 (3)
O2	0.27172 (13)	0.01591 (14)	0.17023 (5)	0.0523 (3)
C1	0.72163 (19)	-0.27856 (18)	0.31576 (7)	0.0450 (3)
H1	0.6494	-0.1987	0.2922	0.054*
C2	0.6609 (2)	-0.41226 (19)	0.34545 (8)	0.0501 (4)
C3	0.7605 (2)	-0.5331 (2)	0.38078 (8)	0.0591 (4)
H3	0.7146	-0.6222	0.4004	0.071*
C4	0.9300 (2)	-0.5191 (2)	0.38638 (8)	0.0614 (4)
H4A	1.0003	-0.5993	0.4105	0.074*
C5	0.9978 (2)	-0.38669 (19)	0.35649 (8)	0.0516 (4)
H5	1.1137	-0.3798	0.3600	0.062*
C6	0.89381 (18)	-0.26449 (17)	0.32148 (6)	0.0410 (3)
C7	0.96683 (18)	-0.12343 (17)	0.29124 (6)	0.0417 (3)
H7	1.0863	-0.1325	0.2841	0.050*
C8	0.85819 (17)	0.28570 (17)	0.23110 (6)	0.0384 (3)
C9	0.96279 (17)	0.41235 (16)	0.20355 (6)	0.0396 (3)
C10	0.8958 (2)	0.53546 (18)	0.15847 (7)	0.0484 (4)
C11	1.0015 (2)	0.6487 (2)	0.13572 (9)	0.0619 (5)
H11	0.9606	0.7303	0.1053	0.074*
C12	1.1644 (3)	0.6444 (2)	0.15654 (9)	0.0660 (5)
H12	1.2315	0.7221	0.1401	0.079*
C13	1.2274 (2)	0.5256 (2)	0.20142 (9)	0.0606 (4)
H13	1.3368	0.5227	0.2160	0.073*
C14	1.12632 (19)	0.41004 (18)	0.22478 (7)	0.0472 (3)
H14	1.1688	0.3293	0.2553	0.057*
C15	0.7180 (3)	0.5522 (3)	0.13424 (10)	0.0746 (6)
H15A	0.6322	0.5901	0.1673	0.112*
H15B	0.7008	0.4457	0.1212	0.112*
H15C	0.7075	0.6314	0.0987	0.112*
C16	0.2917 (2)	-0.0634 (2)	0.41323 (7)	0.0553 (4)
H16	0.2941	-0.1400	0.3821	0.066*
C17	0.2317 (2)	-0.0961 (3)	0.47414 (9)	0.0698 (5)
C18	0.2250 (3)	0.0105 (3)	0.52174 (9)	0.0813 (7)
H18	0.1828	-0.0163	0.5625	0.098*
C19	0.2823 (3)	0.1588 (3)	0.50772 (9)	0.0807 (6)
H19	0.2789	0.2338	0.5395	0.097*
C20	0.3458 (2)	0.1984 (3)	0.44652 (8)	0.0649 (5)
H20	0.3855	0.2987	0.4376	0.078*
C21	0.34927 (18)	0.0876 (2)	0.39894 (7)	0.0477 (4)
C22	0.41875 (18)	0.1272 (2)	0.33473 (7)	0.0469 (3)
H22	0.4926	0.2051	0.3289	0.056*
C23	0.40720 (17)	0.06825 (16)	0.17440 (6)	0.0397 (3)
C24	0.52512 (18)	0.10606 (17)	0.11860 (7)	0.0424 (3)
C25	0.4604 (2)	0.1922 (2)	0.06360 (7)	0.0541 (4)
C26	0.5812 (3)	0.2257 (3)	0.01535 (9)	0.0756 (6)
H26	0.5419	0.2832	-0.0215	0.091*

C27	0.7564 (3)	0.1768 (3)	0.02029 (10)	0.0811 (6)
H27	0.8334	0.2020	-0.0129	0.097*
C28	0.8183 (2)	0.0910 (2)	0.07389 (9)	0.0670 (5)
H28	0.9370	0.0567	0.0772	0.080*
C29	0.70265 (19)	0.0561 (2)	0.12286 (8)	0.0506 (4)
H29	0.7441	-0.0017	0.1594	0.061*
C30	0.2710 (2)	0.2514 (3)	0.05541 (9)	0.0746 (6)
H30A	0.2566	0.3456	0.0257	0.112*
H30B	0.2133	0.2834	0.0955	0.112*
H30C	0.2219	0.1629	0.0398	0.112*
H2	1.0634 (14)	0.108 (3)	0.2319 (9)	0.080*
H4	0.551 (2)	0.157 (2)	0.2320 (10)	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0524 (6)	0.0685 (6)	0.1057 (8)	-0.0265 (5)	-0.0017 (5)	0.0042 (6)
F2	0.1345 (12)	0.1237 (11)	0.0735 (8)	-0.0505 (10)	0.0027 (7)	0.0419 (8)
N1	0.0406 (6)	0.0401 (6)	0.0470 (7)	-0.0131 (5)	0.0031 (5)	0.0061 (5)
N2	0.0343 (6)	0.0405 (6)	0.0540 (7)	-0.0104 (5)	0.0045 (5)	0.0083 (5)
N3	0.0335 (6)	0.0500 (7)	0.0384 (6)	-0.0091 (5)	0.0024 (5)	0.0044 (5)
N4	0.0350 (6)	0.0532 (7)	0.0392 (6)	-0.0164 (5)	0.0043 (5)	0.0025 (5)
O1	0.0339 (5)	0.0459 (6)	0.0639 (7)	-0.0102 (4)	0.0031 (4)	0.0015 (5)
O2	0.0464 (6)	0.0652 (7)	0.0511 (6)	-0.0272 (5)	0.0032 (5)	-0.0064 (5)
C1	0.0458 (8)	0.0395 (7)	0.0492 (8)	-0.0073 (6)	-0.0031 (6)	0.0024 (6)
C2	0.0465 (8)	0.0483 (8)	0.0571 (9)	-0.0151 (7)	0.0019 (7)	-0.0028 (7)
C3	0.0700 (11)	0.0468 (9)	0.0621 (10)	-0.0209 (8)	0.0000 (8)	0.0119 (7)
C4	0.0698 (11)	0.0504 (9)	0.0622 (10)	-0.0090 (8)	-0.0134 (8)	0.0197 (8)
C5	0.0493 (9)	0.0503 (9)	0.0549 (9)	-0.0086 (7)	-0.0090 (7)	0.0067 (7)
C6	0.0449 (8)	0.0392 (7)	0.0384 (7)	-0.0082 (6)	0.0003 (6)	0.0006 (6)
C7	0.0376 (7)	0.0441 (8)	0.0433 (7)	-0.0096 (6)	-0.0001 (6)	0.0024 (6)
C8	0.0359 (7)	0.0406 (7)	0.0395 (7)	-0.0109 (6)	0.0009 (5)	0.0008 (5)
C9	0.0374 (7)	0.0371 (7)	0.0441 (7)	-0.0092 (5)	0.0051 (6)	-0.0011 (6)
C10	0.0503 (9)	0.0420 (8)	0.0506 (8)	-0.0069 (6)	0.0034 (7)	0.0050 (6)
C11	0.0706 (12)	0.0449 (9)	0.0662 (11)	-0.0111 (8)	0.0133 (9)	0.0118 (8)
C12	0.0708 (12)	0.0497 (9)	0.0805 (12)	-0.0297 (8)	0.0199 (9)	0.0004 (9)
C13	0.0490 (9)	0.0605 (10)	0.0776 (12)	-0.0259 (8)	0.0060 (8)	-0.0084 (9)
C14	0.0416 (8)	0.0456 (8)	0.0556 (9)	-0.0129 (6)	0.0016 (6)	-0.0008 (7)
C15	0.0657 (12)	0.0769 (13)	0.0791 (13)	-0.0114 (10)	-0.0182 (10)	0.0291 (10)
C16	0.0528 (9)	0.0725 (11)	0.0402 (8)	-0.0119 (8)	-0.0050 (7)	0.0075 (7)
C17	0.0599 (11)	0.0974 (14)	0.0500 (10)	-0.0156 (10)	-0.0042 (8)	0.0214 (10)
C18	0.0639 (12)	0.133 (2)	0.0398 (10)	-0.0032 (12)	0.0036 (8)	0.0069 (11)
C19	0.0677 (12)	0.1221 (19)	0.0475 (10)	0.0027 (12)	-0.0010 (9)	-0.0255 (11)
C20	0.0539 (10)	0.0865 (13)	0.0546 (10)	-0.0092 (9)	-0.0029 (8)	-0.0151 (9)
C21	0.0337 (7)	0.0681 (10)	0.0394 (8)	-0.0041 (6)	-0.0039 (6)	-0.0008 (7)
C22	0.0379 (8)	0.0583 (9)	0.0460 (8)	-0.0137 (6)	0.0001 (6)	-0.0012 (7)
C23	0.0375 (7)	0.0390 (7)	0.0429 (7)	-0.0099 (6)	0.0035 (6)	-0.0019 (6)
C24	0.0434 (8)	0.0437 (7)	0.0407 (7)	-0.0117 (6)	0.0060 (6)	-0.0054 (6)
C25	0.0562 (9)	0.0662 (10)	0.0395 (8)	-0.0121 (8)	0.0042 (7)	-0.0032 (7)
C26	0.0768 (13)	0.0993 (15)	0.0448 (10)	-0.0104 (11)	0.0115 (9)	0.0120 (9)

C27	0.0740 (13)	0.1000 (15)	0.0625 (12)	-0.0154 (11)	0.0323 (10)	0.0091 (11)
C28	0.0482 (9)	0.0761 (12)	0.0714 (12)	-0.0075 (8)	0.0223 (8)	-0.0017 (9)
C29	0.0438 (8)	0.0538 (9)	0.0518 (9)	-0.0068 (7)	0.0084 (7)	-0.0012 (7)
C30	0.0635 (12)	0.1075 (16)	0.0501 (10)	-0.0086 (11)	-0.0095 (8)	0.0069 (10)

Geometric parameters (\AA , $\text{^{\circ}}$)

F1—C2	1.3598 (18)	C13—C14	1.384 (2)
F2—C17	1.357 (2)	C13—H13	0.9300
N1—C7	1.2735 (18)	C14—H14	0.9300
N1—N2	1.3824 (15)	C15—H15A	0.9600
N2—C8	1.3549 (18)	C15—H15B	0.9600
N2—H2	0.894 (9)	C15—H15C	0.9600
N3—C22	1.2689 (19)	C16—C17	1.367 (2)
N3—N4	1.3815 (15)	C16—C21	1.394 (2)
N4—C23	1.3506 (18)	C16—H16	0.9300
N4—H4	0.893 (9)	C17—C18	1.360 (3)
O1—C8	1.2263 (16)	C18—C19	1.372 (3)
O2—C23	1.2249 (16)	C18—H18	0.9300
C1—C2	1.367 (2)	C19—C20	1.394 (3)
C1—C6	1.391 (2)	C19—H19	0.9300
C1—H1	0.9300	C20—C21	1.386 (2)
C2—C3	1.366 (2)	C20—H20	0.9300
C3—C4	1.370 (3)	C21—C22	1.467 (2)
C3—H3	0.9300	C22—H22	0.9300
C4—C5	1.387 (2)	C23—C24	1.4984 (18)
C4—H4A	0.9300	C24—C29	1.389 (2)
C5—C6	1.386 (2)	C24—C25	1.404 (2)
C5—H5	0.9300	C25—C26	1.390 (2)
C6—C7	1.4675 (19)	C25—C30	1.499 (3)
C7—H7	0.9300	C26—C27	1.373 (3)
C8—C9	1.4966 (18)	C26—H26	0.9300
C9—C14	1.388 (2)	C27—C28	1.371 (3)
C9—C10	1.403 (2)	C27—H27	0.9300
C10—C11	1.390 (2)	C28—C29	1.378 (2)
C10—C15	1.503 (2)	C28—H28	0.9300
C11—C12	1.377 (3)	C29—H29	0.9300
C11—H11	0.9300	C30—H30A	0.9600
C12—C13	1.369 (3)	C30—H30B	0.9600
C12—H12	0.9300	C30—H30C	0.9600
C7—N1—N2	115.08 (12)	H15A—C15—H15B	109.5
C8—N2—N1	119.04 (11)	C10—C15—H15C	109.5
C8—N2—H2	120.3 (13)	H15A—C15—H15C	109.5
N1—N2—H2	120.0 (13)	H15B—C15—H15C	109.5
C22—N3—N4	114.56 (12)	C17—C16—C21	118.33 (17)
C23—N4—N3	120.56 (11)	C17—C16—H16	120.8
C23—N4—H4	121.3 (13)	C21—C16—H16	120.8
N3—N4—H4	117.7 (13)	F2—C17—C18	118.24 (17)
C2—C1—C6	118.59 (14)	F2—C17—C16	118.05 (19)

C2—C1—H1	120.7	C18—C17—C16	123.7 (2)
C6—C1—H1	120.7	C17—C18—C19	117.97 (18)
F1—C2—C3	118.08 (14)	C17—C18—H18	121.0
F1—C2—C1	118.55 (14)	C19—C18—H18	121.0
C3—C2—C1	123.37 (15)	C18—C19—C20	120.86 (19)
C2—C3—C4	117.99 (14)	C18—C19—H19	119.6
C2—C3—H3	121.0	C20—C19—H19	119.6
C4—C3—H3	121.0	C21—C20—C19	119.7 (2)
C3—C4—C5	120.65 (15)	C21—C20—H20	120.2
C3—C4—H4A	119.7	C19—C20—H20	120.2
C5—C4—H4A	119.7	C20—C21—C16	119.46 (15)
C6—C5—C4	120.37 (15)	C20—C21—C22	119.93 (16)
C6—C5—H5	119.8	C16—C21—C22	120.58 (14)
C4—C5—H5	119.8	N3—C22—C21	121.35 (13)
C5—C6—C1	119.02 (13)	N3—C22—H22	119.3
C5—C6—C7	119.63 (13)	C21—C22—H22	119.3
C1—C6—C7	121.36 (13)	O2—C23—N4	123.05 (12)
N1—C7—C6	120.57 (13)	O2—C23—C24	124.02 (13)
N1—C7—H7	119.7	N4—C23—C24	112.93 (11)
C6—C7—H7	119.7	C29—C24—C25	120.14 (13)
O1—C8—N2	122.54 (12)	C29—C24—C23	118.20 (13)
O1—C8—C9	122.65 (12)	C25—C24—C23	121.65 (13)
N2—C8—C9	114.79 (11)	C26—C25—C24	117.09 (16)
C14—C9—C10	119.96 (13)	C26—C25—C30	119.45 (16)
C14—C9—C8	118.96 (12)	C24—C25—C30	123.45 (14)
C10—C9—C8	121.05 (12)	C27—C26—C25	122.25 (18)
C11—C10—C9	117.18 (14)	C27—C26—H26	118.9
C11—C10—C15	118.83 (15)	C25—C26—H26	118.9
C9—C10—C15	123.98 (14)	C28—C27—C26	120.26 (16)
C12—C11—C10	122.50 (16)	C28—C27—H27	119.9
C12—C11—H11	118.8	C26—C27—H27	119.9
C10—C11—H11	118.8	C27—C28—C29	119.17 (17)
C13—C12—C11	119.88 (15)	C27—C28—H28	120.4
C13—C12—H12	120.1	C29—C28—H28	120.4
C11—C12—H12	120.1	C28—C29—C24	121.09 (16)
C12—C13—C14	119.22 (16)	C28—C29—H29	119.5
C12—C13—H13	120.4	C24—C29—H29	119.5
C14—C13—H13	120.4	C25—C30—H30A	109.5
C13—C14—C9	121.24 (15)	C25—C30—H30B	109.5
C13—C14—H14	119.4	H30A—C30—H30B	109.5
C9—C14—H14	119.4	C25—C30—H30C	109.5
C10—C15—H15A	109.5	H30A—C30—H30C	109.5
C10—C15—H15B	109.5	H30B—C30—H30C	109.5

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N4—H4···O1	0.89 (1)	1.93 (1)	2.8115 (15)	168 (2)

supplementary materials

N2—H2 \cdots O2 ⁱ	0.89 (1)	2.08 (1)	2.9185 (16)	157 (2)
--------------------------------	----------	----------	-------------	---------

Symmetry code: (i) $x+1, y, z$.