

N-(4-Methoxy-2-nitrophenyl)-N-(methylsulfonyl)methanesulfonamide

Sammer Yousuf,^{a*} Hina Siddiqui,^a Rabia Farooq^a and M. Iqbal Choudhary^{a,b}

^aH.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan, and

^bDepartment of Biochemistry, Faculty of Sciences, King Abdul Aziz University, Jeddah 21589, Saudi Arabia

Correspondence e-mail: dr.sammer.yousuf@gmail.com


Received 29 May 2012; accepted 31 May 2012

Key indicators: single-crystal X-ray study; $T = 273\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.040; wR factor = 0.113; data-to-parameter ratio = 18.2.

In the title compound, $\text{C}_9\text{H}_{12}\text{N}_2\text{O}_7\text{S}_2$, the nitro substituent is slightly twisted from the benzene ring [dihedral angle = $14.69(10)^\circ$]. The molecular geometry is stabilized by intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming $S(6)$ ring motifs. In the crystal, molecules are linked by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds into layers parallel to $(10\bar{2})$.

Related literature

For the biological activities of sulfonamides, see: Alsughayer *et al.* (2011); Joshi & Khosla (2003); Scozzafava *et al.* (2003); Drews (2000); Peixoto & Beverley (1987). For crystal structures of closely related compounds, see: Boechat *et al.* (2010); Zia-ur-Rehman *et al.* (2009).



Experimental

Crystal data

$\text{C}_9\text{H}_{12}\text{N}_2\text{O}_7\text{S}_2$
 $M_r = 324.33$

Monoclinic, $P2_1/c$
 $a = 9.4976(7)\text{ \AA}$

$b = 7.5987(6)\text{ \AA}$
 $c = 19.2434(15)\text{ \AA}$
 $\beta = 103.672(2)^\circ$
 $V = 1349.43(18)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.43\text{ mm}^{-1}$
 $T = 273\text{ K}$
 $0.55 \times 0.47 \times 0.11\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.799$, $T_{\max} = 0.955$

9504 measured reflections
3362 independent reflections
2711 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.113$
 $S = 1.05$
3362 reflections

185 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.31\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.30\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C1—H1B \cdots O2 ⁱ	0.93	2.46	3.368 (2)	165
C8—H8B \cdots O4	0.96	2.58	3.225 (3)	125
C8—H8C \cdots O5 ⁱⁱ	0.96	2.58	3.250 (3)	127
C9—H9B \cdots O1	0.96	2.59	3.226 (3)	124
C9—H9B \cdots O1 ⁱⁱⁱ	0.96	2.47	3.173 (3)	130

Symmetry codes: (i) $-x + 2$, $y - \frac{1}{2}$, $-z + \frac{1}{2}$; (ii) x , $y + 1$, z ; (iii) $-x + 1$, $-y$, $-z$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*, *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5148).

References

- Alsughayer, A., Elassar, A. A., Mustafa, S. & Sagheer, F. (2011). *J. Biomater. Nanobiotechnol.* **2**, 144–149.
- Boechat, N., Santos Lages, A. dos, Kover, W. B., Tiekkink, E. R. T., Wardell, J. L. & Wardell, S. M. S. V. (2010). *Acta Cryst. E66*, o506–o507.
- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Drews, J. (2000). *Science*, **287**, 1960–1964.
- Joshi, S. & Khosla, N. (2003). *Bioorg. Med. Chem. Lett.* **13**, 3747–3751.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Peixoto, M. P. & Beverley, S. M. (1987). *Antimicrob. Agents Chemother.* **31**, 1575–1578.
- Scozzafava, A., Owa, T., Mastrolorenzo, A. & Supuran, C. (2003). *Curr. Med. Chem.* **10**, 925–953.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Zia-ur-Rehman, M., Sepehrianazar, A., Ali, M., Siddiqui, W. A. & Çaylak, N. (2009). *Acta Cryst. E65*, o941.

supplementary materials

Acta Cryst. (2012). E68, o2090 [doi:10.1107/S160053681202483X]

N-(4-Methoxy-2-nitrophenyl)-N-(methylsulfonyl)methanesulfonamide

Sammer Yousuf, Hina Siddiqui, Rabia Farooq and M. Iqbal Choudhary

Comment

Compounds containing the sulfonamide moiety have attracted a wide interest due to their interesting chemical and biological properties, which makes them promising candidates in drug discovery. Sulfonamides posses wide variety of biological activities including anti-bacterial, anti-leishmanial, anti-inflammatory, anti-cancer, and carbonic anhydrase inhibitory activities (Alsughayer *et al.*, 2011; Joshi & Khosla, 2003; Scozzafava *et al.*, 2003; Drews, 2000; Peixoto & Beverley, 1987). The title compound was prepared as a part of our ongoing research to synthesize different sulfonamide derivatives to study their bioactive potential and structure activity relationship (SAR). In the title compound (Fig. 1), the nitro group was found to be slightly twisted with the dihedral angle of 14.69 (10) $^{\circ}$ between the NO₂ group and the benzene ring. The S1—N2—C3—C4 and S2—N2—C3—C4 torsion angles are 82.83 (19) and -92.42 (17) $^{\circ}$, respectively. The molecule is stabilized by two intramolecular C8—H8B···O4 and C9—H9B···O1 interactions to form two S(6) ring motifs. In the crystal structure, the molecules are linked to form a two-dimensional network through C1—H1B···O2ⁱ, C8—H8C···O5ⁱⁱ and C9—H9B···O1ⁱⁱⁱ intermolecular hydrogen bonds (Fig. 2 and Table 1). The bond lengths and angles are within the normal range and similar to other closely related structures (Boechat *et al.*, 2010; Zia-ur-Rehman *et al.*, 2009).

Experimental


To a stirring solution of methanesulfonyl chloride (1.0 g, 8.7 mmol) in CH₂Cl₂ (20 ml) at 0 °C, 3 ml Et₃N and 4-methoxy-2-nitroaniline (1.1 eq., 1.61 g m, 9.6 mmol) were added along with catalytic amount of dimethylamino pyridine (DMAP). Progress of the reaction was monitored by thin layer chromatography in 7:3 hexanes: ethyl acetate solvent system. After complete consumption of starting material (2 hrs), workup was performed with H₂O (10 ml), organic layer was separated and aqueous layer was extracted with CH₂Cl₂ (2 × 10 ml). Organic layers were further washed with brine (10 ml), and dried over MgSO₄, filtered, and concentrated in vacuum to obtain the crude product (0.9 g, 90% yield). Flash chromatography was performed hexanes: ethyl acetate (7:3), to obtain crystalline compound I, in 55% yield. Crystals were found suitable for single-crystal X-ray diffraction studies. All the starting materials and solvents were purchased from commercial suppliers and used for reaction without purification.

Refinement

H atoms were positioned geometrically with C—H = 0.96 or 0.93 Å, and constrained to ride on their parent atoms with U_{iso}(H)= 1.2U_{eq}(C) or 1.5U_{eq}(C_{methyl}). A rotating group model was applied to the methyl groups.

Computing details

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT (Bruker, 2000); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL (Sheldrick, 2008), PARST (Nardelli, 1995) and PLATON (Spek, 2009).

**Figure 1**

The molecular structure of the title compound with displacement ellipsoids drawn at 30% probability level.




Figure 2

The crystal packing of the title compound. Only hydrogen atoms involved in hydrogen bonding are shown.

N*-(4-Methoxy-2-nitrophenyl)-*N*-(methylsulfonyl)methanesulfonamideCrystal data*

$C_9H_{12}N_2O_7S_2$	$F(000) = 672$
$M_r = 324.33$	$D_x = 1.596 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3096 reflections
$a = 9.4976 (7) \text{ \AA}$	$\theta = 2.2\text{--}27.7^\circ$
$b = 7.5987 (6) \text{ \AA}$	$\mu = 0.43 \text{ mm}^{-1}$
$c = 19.2434 (15) \text{ \AA}$	$T = 273 \text{ K}$
$\beta = 103.672 (2)^\circ$	Block, yellow
$V = 1349.43 (18) \text{ \AA}^3$	$0.55 \times 0.47 \times 0.11 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART APEX CCD area-detector diffractometer	9504 measured reflections
Radiation source: fine-focus sealed tube	3362 independent reflections
Graphite monochromator	2711 reflections with $I > 2\sigma(I)$
ω scan	$R_{\text{int}} = 0.023$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2000)	$\theta_{\max} = 28.3^\circ, \theta_{\min} = 2.2^\circ$
$T_{\min} = 0.799, T_{\max} = 0.955$	$h = -12 \rightarrow 12$
	$k = -10 \rightarrow 9$
	$l = -25 \rightarrow 23$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.040$	$w = 1/[\sigma^2(F_o^2) + (0.0641P)^2 + 0.2074P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.113$	$(\Delta/\sigma)_{\max} = 0.001$
$S = 1.05$	$\Delta\rho_{\max} = 0.31 \text{ e \AA}^{-3}$
3362 reflections	$\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$
185 parameters	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
0 restraints	Extinction coefficient: 0.0101 (14)
Primary atom site location: structure-invariant direct methods	
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.80706 (5)	0.21065 (6)	-0.02024 (2)	0.03866 (15)
S2	0.67671 (5)	0.31016 (6)	0.09782 (2)	0.03825 (15)
O1	0.55361 (15)	0.22596 (19)	0.05410 (8)	0.0529 (4)
O2	0.71498 (16)	0.2788 (2)	0.17294 (7)	0.0514 (4)
O3	0.95185 (15)	0.2109 (2)	-0.02879 (7)	0.0522 (4)
O4	0.70649 (16)	0.3355 (2)	-0.05856 (7)	0.0561 (4)
O5	0.80101 (17)	-0.07491 (19)	0.12618 (9)	0.0608 (4)
O6	0.99411 (19)	-0.20742 (17)	0.18046 (9)	0.0602 (4)
O7	1.36416 (14)	0.2232 (2)	0.25913 (8)	0.0503 (4)
N1	0.92993 (18)	-0.07629 (18)	0.15472 (8)	0.0381 (3)
N2	0.82155 (15)	0.24662 (18)	0.06736 (7)	0.0329 (3)
C1	1.14724 (19)	0.0885 (2)	0.20463 (9)	0.0345 (4)
H1B	1.1802	-0.0117	0.2313	0.041*
C2	1.01332 (18)	0.0890 (2)	0.15816 (8)	0.0311 (3)
C3	0.96028 (17)	0.2371 (2)	0.11724 (8)	0.0315 (3)
C4	1.04925 (19)	0.3835 (2)	0.12521 (10)	0.0406 (4)
H4B	1.0171	0.4833	0.0982	0.049*
C5	1.1838 (2)	0.3863 (2)	0.17185 (10)	0.0428 (4)
H5A	1.2407	0.4872	0.1764	0.051*
C6	1.23354 (19)	0.2382 (3)	0.21179 (9)	0.0366 (4)
C7	1.4561 (2)	0.3748 (3)	0.27101 (13)	0.0630 (6)
H7A	1.5458	0.3458	0.3040	0.095*
H7B	1.4090	0.4680	0.2905	0.095*
H7C	1.4752	0.4123	0.2265	0.095*
C8	0.6647 (3)	0.5369 (3)	0.08270 (13)	0.0576 (6)
H8A	0.5830	0.5830	0.0981	0.086*
H8B	0.6528	0.5598	0.0326	0.086*
H8C	0.7517	0.5927	0.1091	0.086*
C9	0.7335 (2)	-0.0003 (3)	-0.03754 (11)	0.0507 (5)
H9A	0.7196	-0.0262	-0.0875	0.076*
H9B	0.6421	-0.0054	-0.0246	0.076*
H9C	0.7986	-0.0849	-0.0100	0.076*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0379 (3)	0.0453 (3)	0.0321 (2)	-0.00244 (18)	0.00691 (18)	0.00241 (17)
S2	0.0339 (2)	0.0363 (2)	0.0449 (3)	-0.00356 (16)	0.00994 (18)	-0.00732 (18)
O1	0.0338 (7)	0.0560 (9)	0.0673 (10)	-0.0104 (6)	0.0088 (6)	-0.0157 (7)
O2	0.0573 (9)	0.0568 (9)	0.0437 (8)	-0.0021 (7)	0.0191 (6)	-0.0065 (6)
O3	0.0448 (8)	0.0697 (10)	0.0460 (8)	-0.0059 (7)	0.0185 (6)	-0.0001 (7)
O4	0.0582 (9)	0.0632 (9)	0.0421 (7)	0.0097 (7)	0.0020 (6)	0.0133 (7)
O5	0.0512 (9)	0.0431 (8)	0.0834 (11)	-0.0183 (7)	0.0065 (8)	0.0071 (7)
O6	0.0801 (12)	0.0310 (7)	0.0648 (10)	-0.0038 (7)	0.0078 (8)	0.0130 (7)
O7	0.0351 (7)	0.0564 (9)	0.0529 (8)	-0.0008 (6)	-0.0029 (6)	-0.0013 (7)
N1	0.0530 (9)	0.0287 (7)	0.0343 (7)	-0.0084 (6)	0.0137 (6)	-0.0001 (6)
N2	0.0306 (7)	0.0356 (7)	0.0310 (7)	-0.0017 (6)	0.0044 (5)	-0.0005 (6)

C1	0.0407 (9)	0.0316 (8)	0.0311 (8)	0.0033 (7)	0.0082 (6)	0.0031 (6)
C2	0.0388 (9)	0.0254 (8)	0.0307 (8)	-0.0054 (6)	0.0114 (6)	-0.0010 (6)
C3	0.0323 (8)	0.0300 (8)	0.0317 (8)	-0.0036 (6)	0.0062 (6)	0.0016 (6)
C4	0.0416 (10)	0.0297 (8)	0.0466 (10)	-0.0064 (7)	0.0029 (7)	0.0078 (7)
C5	0.0403 (10)	0.0347 (9)	0.0508 (11)	-0.0112 (7)	0.0054 (8)	0.0039 (8)
C6	0.0336 (9)	0.0417 (9)	0.0337 (9)	-0.0008 (7)	0.0065 (7)	-0.0032 (7)
C7	0.0358 (11)	0.0725 (15)	0.0743 (15)	-0.0108 (10)	0.0002 (10)	-0.0152 (13)
C8	0.0612 (14)	0.0352 (10)	0.0733 (15)	0.0068 (9)	0.0096 (11)	-0.0051 (10)
C9	0.0521 (12)	0.0529 (12)	0.0456 (11)	-0.0100 (9)	0.0089 (9)	-0.0160 (9)

Geometric parameters (\AA , $^{\circ}$)

S1—O4	1.4223 (15)	C1—H1B	0.9300
S1—O3	1.4231 (14)	C2—C3	1.397 (2)
S1—N2	1.6809 (14)	C3—C4	1.383 (2)
S1—C9	1.748 (2)	C4—C5	1.377 (2)
S2—O1	1.4210 (14)	C4—H4B	0.9300
S2—O2	1.4247 (14)	C5—C6	1.382 (3)
S2—N2	1.6882 (15)	C5—H5A	0.9300
S2—C8	1.747 (2)	C7—H7A	0.9600
O5—N1	1.218 (2)	C7—H7B	0.9600
O6—N1	1.211 (2)	C7—H7C	0.9600
O7—C6	1.359 (2)	C8—H8A	0.9600
O7—C7	1.431 (3)	C8—H8B	0.9600
N1—C2	1.478 (2)	C8—H8C	0.9600
N2—C3	1.437 (2)	C9—H9A	0.9600
C1—C2	1.371 (2)	C9—H9B	0.9600
C1—C6	1.389 (2)	C9—H9C	0.9600
O4—S1—O3	119.21 (9)	C5—C4—C3	122.23 (16)
O4—S1—N2	107.30 (8)	C5—C4—H4B	118.9
O3—S1—N2	105.27 (8)	C3—C4—H4B	118.9
O4—S1—C9	108.87 (10)	C4—C5—C6	119.50 (17)
O3—S1—C9	109.43 (10)	C4—C5—H5A	120.3
N2—S1—C9	105.93 (9)	C6—C5—H5A	120.3
O1—S2—O2	120.15 (9)	O7—C6—C5	125.38 (17)
O1—S2—N2	106.83 (8)	O7—C6—C1	114.93 (17)
O2—S2—N2	105.73 (8)	C5—C6—C1	119.70 (16)
O1—S2—C8	109.48 (11)	O7—C7—H7A	109.5
O2—S2—C8	108.96 (10)	O7—C7—H7B	109.5
N2—S2—C8	104.52 (10)	H7A—C7—H7B	109.5
C6—O7—C7	117.75 (16)	O7—C7—H7C	109.5
O6—N1—O5	123.14 (15)	H7A—C7—H7C	109.5
O6—N1—C2	117.92 (15)	H7B—C7—H7C	109.5
O5—N1—C2	118.94 (14)	S2—C8—H8A	109.5
C3—N2—S1	120.43 (11)	S2—C8—H8B	109.5
C3—N2—S2	118.43 (11)	H8A—C8—H8B	109.5
S1—N2—S2	120.96 (8)	S2—C8—H8C	109.5
C2—C1—C6	119.80 (16)	H8A—C8—H8C	109.5
C2—C1—H1B	120.1	H8B—C8—H8C	109.5

C6—C1—H1B	120.1	S1—C9—H9A	109.5
C1—C2—C3	121.66 (15)	S1—C9—H9B	109.5
C1—C2—N1	115.57 (14)	H9A—C9—H9B	109.5
C3—C2—N1	122.77 (15)	S1—C9—H9C	109.5
C4—C3—C2	117.11 (15)	H9A—C9—H9C	109.5
C4—C3—N2	118.25 (15)	H9B—C9—H9C	109.5
C2—C3—N2	124.63 (15)		
O4—S1—N2—C3	-139.57 (14)	C1—C2—C3—C4	0.4 (2)
O3—S1—N2—C3	-11.63 (15)	N1—C2—C3—C4	-179.79 (15)
C9—S1—N2—C3	104.25 (14)	C1—C2—C3—N2	179.70 (15)
O4—S1—N2—S2	35.55 (12)	N1—C2—C3—N2	-0.5 (3)
O3—S1—N2—S2	163.50 (10)	S1—N2—C3—C4	82.83 (19)
C9—S1—N2—S2	-80.63 (12)	S2—N2—C3—C4	-92.42 (17)
O1—S2—N2—C3	-150.32 (13)	S1—N2—C3—C2	-96.42 (17)
O2—S2—N2—C3	-21.24 (15)	S2—N2—C3—C2	88.33 (19)
C8—S2—N2—C3	93.69 (15)	C2—C3—C4—C5	-0.8 (3)
O1—S2—N2—S1	34.46 (12)	N2—C3—C4—C5	179.87 (17)
O2—S2—N2—S1	163.53 (10)	C3—C4—C5—C6	0.7 (3)
C8—S2—N2—S1	-81.53 (12)	C7—O7—C6—C5	2.6 (3)
C6—C1—C2—C3	0.0 (3)	C7—O7—C6—C1	-177.73 (17)
C6—C1—C2—N1	-179.76 (15)	C4—C5—C6—O7	179.46 (18)
O6—N1—C2—C1	-14.8 (2)	C4—C5—C6—C1	-0.2 (3)
O5—N1—C2—C1	165.47 (17)	C2—C1—C6—O7	-179.86 (15)
O6—N1—C2—C3	165.42 (17)	C2—C1—C6—C5	-0.2 (3)
O5—N1—C2—C3	-14.3 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C1—H1B···O2 ⁱ	0.93	2.46	3.368 (2)	165
C8—H8B···O4	0.96	2.58	3.225 (3)	125
C8—H8C···O5 ⁱⁱ	0.96	2.58	3.250 (3)	127
C9—H9B···O1	0.96	2.59	3.226 (3)	124
C9—H9B···O1 ⁱⁱⁱ	0.96	2.47	3.173 (3)	130

Symmetry codes: (i) $-x+2, y-1/2, -z+1/2$; (ii) $x, y+1, z$; (iii) $-x+1, -y, -z$.