

3,4-Dimethoxy-N-((E)-4-{[1-(prop-2-en-1-yl)-1H-1,2,3-triazol-4-yl]methoxy}-benzylidene)aniline

Mehmet Akkurt,^{a*} Aliasghar Jarrahpour,^b Hashem Sharghi,^b Mehdi Mohammadi Chermahini,^b Pezhman Shiri^b and Orhan Büyükgüngör^c

^aDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^bDepartment of Chemistry, College of Sciences, Shiraz University, 71454 Shiraz, Iran, and ^cDepartment of Physics, Faculty of Arts and Sciences, Ondokuz Mayıs University, 55139 Samsun, Turkey
Correspondence e-mail: akkurt@erciyes.edu.tr


Received 7 June 2012; accepted 13 June 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.041; wR factor = 0.107; data-to-parameter ratio = 15.5.

In the title compound, $\text{C}_{21}\text{H}_{22}\text{N}_4\text{O}_3$, the triazole ring is planar [maximum deviation = 0.004 (1) \AA] and makes dihedral angles of 26.21 (8) and 38.66 (8) $^\circ$ with the two benzene rings. In the crystal, molecules are linked by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming zigzag chains along [111]. In addition, a weak $\text{C}-\text{H}\cdots\pi$ interaction is also observed.

Related literature

For background on the importance of triazole derivatives and their uses, see: da Silva *et al.* (2011); Hranjec *et al.* (2011); Shi & Zhou (2011); Yap & Weinreb (2006); Yu *et al.* (2006).

**Experimental***Crystal data*

$\text{C}_{21}\text{H}_{22}\text{N}_4\text{O}_3$
 $M_r = 378.43$
Triclinic, $P\bar{1}$
 $a = 5.8144 (4)\text{ \AA}$
 $b = 11.9677 (8)\text{ \AA}$

$c = 14.1019 (10)\text{ \AA}$
 $\alpha = 85.803 (6)^\circ$
 $\beta = 80.497 (6)^\circ$
 $\gamma = 80.434 (5)^\circ$
 $V = 953.30 (12)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.73 \times 0.41 \times 0.21\text{ mm}$

Data collection

Stoe IPDS 2 diffractometer
Absorption correction: integration (*X-RED32*; Stoe & Cie, 2002)
 $T_{\min} = 0.957$, $T_{\max} = 0.981$
15258 measured reflections
3919 independent reflections
2795 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.050$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.107$
 $S = 1.04$
3919 reflections
253 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.14\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.18\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg2$ is the centroid of the C1–C6 benzene ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C19—H19A \cdots O2 ⁱ	0.97	2.54	3.385 (2)	146
C16—H16A \cdots Cg2 ⁱⁱ	0.97	2.87	3.6647 (18)	140

Symmetry codes: (i) $x + 1, y - 1, z + 1$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDS 2 diffractometer (purchased under grant F.279 of the University Research Fund). AJ, HS, MMC and PS also thank the Shiraz University Research Council for financial support (grant No. 89-GR-SC-23).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2500).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hranjec, M., Starcevic, K., Pavelic, S. K., Lucin, P., Pavelic, K. & Zamola, G. K. (2011). *Eur. J. Med. Chem.* **46**, 2274–2279.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shi, Y. & Zhou, C. H. (2011). *Bioorg. Med. Chem. Lett.* **21**, 956–960.
- Silva, C. M. da, da Silva, D. L., Modolo, L. V., Alves, R. B., de Resende, M. A., Martins, C. V. B. & de Fatima, A. (2011). *J. Adv. Res.* **2**, 1–8.
- Stoe & Cie (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.
- Yap, A. H. & Weinreb, S. M. (2006). *Tetrahedron Lett.* **47**, 3035–3038.
- Yu, H.-X., Ma, J.-F., Xu, G.-H., Li, S.-L., Yang, J., Liu, Y.-Y. & Cheng, Y.-X. (2006). *J. Organomet. Chem.* **691**, 3531–3539.

supplementary materials

Acta Cryst. (2012). E68, o2127 [doi:10.1107/S1600536812026761]

3,4-Dimethoxy-N-((E)-4-{[1-(prop-2-en-1-yl)-1H-1,2,3-triazol-4-yl]methoxy}-benzylidene)aniline

Mehmet Akkurt, Aliashghar Jarrahpour, Hashem Sharghi, Mehdi Mohammadi Chermahini, Pezhman Shiri and Orhan Büyükgüngör

Comment

Schiff bases are aldehyde- or ketone-like compounds in which the carbonyl group is replaced by an imine or azomethine group. They are widely used for industrial purposes and also exhibit a broad range of biological activities (da Silva *et al.*, 2011). These compounds represent an important class of organic compounds, especially in the medicinal and pharmaceutical field. Thus, development and synthesis of novel Schiff base derivatives as potential chemotherapeutics still attracts attention of organic and medicinal chemists (Hranjec *et al.*, 2011). 1,2,3-Triazoles are nitrogen heterocycles, which have a number of important industrial, agrochemical, and pharmaceutical uses (Yap & Weinreb, 2006). Triazole derivatives also display a broad range of biological activity, showing potential applications as antitumor, antibacterial, antifungal and antiviral agents (Yu *et al.*, 2006). Recently, some triazole derivatives have been reported to exhibit good anti-MRSA potency (Shi & Zhou, 2011). Therefore, compound (I), which has the two mentioned features, was synthesized and its X-ray studies is reported here.

The title compound, (Fig. 1), has a non-planar conformation. The N2—N4/C17—C18 triazole ring is essentially planar with a maximum deviaton of 0.004 (1) Å. The dihedral angles between the rings in (I) are: A/B = 26.21 (8), A/C = 38.66 (8) and B/C = 64.87 (7)°, where A, B and C define the C1—C6 and C10—C15 benzene rings, and the N2—N4/C17—C18 triazole ring, respectively.

The crystal packing is stabilized by intermolecular C—H···O hydrogen bonding interactions (Table 1) forming zigzag chains along the [1 -1 1] direction (Fig. 2). Furthermore, a weak intermolecular C—H···π interaction contributes to the stability of the crystal structure.

Experimental


Reaction of 4-((1-allyl-1*H*-1, 2, 3-triazol-4-yl)methoxy)benzaldehyde with 3,4-dimethoxyaniline in refluxing ethanol gave the title compound. Recrystallization from ethanol gave brown crystals in 85% yield. [mp: 363–365 K]. IR (KBr, cm^{-1}): 1573 (N=N), 1612 (C=N). $^1\text{H-NMR}$ (250 MHz, CDCl_3) δ (p.p.m): 3.87–3.90 (2OMe, s, 6H), 4.97 (d, 2H, $J=6.0$ Hz), 5.25–5.37 (m, 4H), 5.93–6.1(m, 1H), 6.84–7.89(m, ArH, 7H), 7.62(s, 1H, Htriazole), 8.41(HC=N, s, 1H). $^{13}\text{C-NMR}$ (CDCl_3) δ (p.p.m): 62.8 (N—CH₂), 55.9–56.0 (2OMe), 62.0 (O—CH₂), 105.7–157.9 (C=C aromatic carbons and triazole), 160.96(C=N).

Refinement


H atoms were positioned geometrically with C—H = 0.93, 0.96 and 0.97 Å, and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{C})$.

Computing details

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA* (Stoe & Cie, 2002); data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

**Figure 1**

The title molecule shown with the displacement ellipsoids for non-H atoms at the 30% probability level.

**Figure 2**

Crystal packing and hydrogen bonding diagram of the title compound shown down the a axis. H atoms not participating in hydrogen bonding have been omitted for clarity.

3,4-Dimethoxy-N-((E)-4-{{[1-(prop-2-en-1-yl)-1*H*-1,2,3-triazol-4-yl]methoxy}benzylidene)aniline

Crystal data

$C_{21}H_{22}N_4O_3$
 $M_r = 378.43$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 5.8144 (4)$ Å
 $b = 11.9677 (8)$ Å
 $c = 14.1019 (10)$ Å
 $\alpha = 85.803 (6)^\circ$
 $\beta = 80.497 (6)^\circ$
 $\gamma = 80.434 (5)^\circ$
 $V = 953.30 (12)$ Å³

$Z = 2$
 $F(000) = 400$
 $D_x = 1.318 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 16791 reflections
 $\theta = 2.2\text{--}27.3^\circ$
 $\mu = 0.09 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
Prism, brown
 $0.73 \times 0.41 \times 0.21$ mm

Data collection

Stoe IPDS 2
 diffractometer
 Radiation source: sealed X-ray tube, 12 x 0.4
 mm long-fine focus
 Plane graphite monochromator
 Detector resolution: 6.67 pixels mm⁻¹
 ω scans
 Absorption correction: integration
 $(X\text{-RED}32;$ Stoe & Cie, 2002)

$T_{\min} = 0.957, T_{\max} = 0.981$
 15258 measured reflections
 3919 independent reflections
 2795 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.050$
 $\theta_{\max} = 26.5^\circ, \theta_{\min} = 2.2^\circ$
 $h = -7 \rightarrow 7$
 $k = -15 \rightarrow 15$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.107$
 $S = 1.04$
 3919 reflections
 253 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0603P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.14 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating - R -factor-obs etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.04640 (19)	0.88296 (10)	0.07531 (10)	0.0725 (4)
O2	0.35022 (17)	0.75647 (9)	0.03759 (8)	0.0615 (4)
O3	0.67186 (18)	0.09568 (8)	0.56736 (8)	0.0618 (4)
N1	0.1657 (2)	0.50851 (10)	0.32214 (9)	0.0553 (4)
N2	1.0236 (2)	-0.14989 (10)	0.76913 (9)	0.0557 (4)
N3	0.7884 (2)	-0.14280 (12)	0.78848 (11)	0.0693 (5)
N4	0.6981 (2)	-0.05436 (12)	0.73864 (11)	0.0661 (5)
C1	0.1137 (2)	0.60697 (11)	0.26194 (11)	0.0503 (4)
C2	-0.1013 (2)	0.67614 (12)	0.28129 (11)	0.0561 (5)
C3	-0.1602 (3)	0.77029 (13)	0.22197 (12)	0.0581 (5)
C4	-0.0060 (2)	0.79558 (11)	0.14080 (11)	0.0526 (5)
C5	0.2124 (2)	0.72537 (11)	0.12021 (10)	0.0495 (4)
C6	0.2718 (2)	0.63275 (11)	0.17977 (11)	0.0508 (4)
C7	-0.2711 (3)	0.95187 (13)	0.08467 (16)	0.0732 (6)
C8	0.5684 (3)	0.68677 (14)	0.00812 (13)	0.0653 (6)
C9	0.3712 (3)	0.48356 (12)	0.34398 (11)	0.0537 (5)

C10	0.4505 (2)	0.38187 (12)	0.40066 (11)	0.0504 (4)
C11	0.6468 (2)	0.37782 (12)	0.44608 (12)	0.0554 (5)
C12	0.7232 (2)	0.28590 (12)	0.50323 (11)	0.0550 (5)
C13	0.6079 (2)	0.19201 (11)	0.51366 (11)	0.0497 (4)
C14	0.4171 (2)	0.19244 (12)	0.46571 (11)	0.0539 (5)
C15	0.3388 (2)	0.28612 (12)	0.41111 (11)	0.0534 (5)
C16	0.8385 (3)	0.10194 (13)	0.62971 (13)	0.0608 (5)
C17	0.8768 (2)	-0.00575 (12)	0.68750 (11)	0.0523 (4)
C18	1.0838 (2)	-0.06698 (12)	0.70649 (11)	0.0545 (5)
C19	1.1770 (3)	-0.23609 (13)	0.81922 (14)	0.0659 (6)
C20	1.1179 (3)	-0.35129 (13)	0.81555 (13)	0.0613 (5)
C21	1.1076 (3)	-0.42391 (14)	0.88837 (14)	0.0708 (6)
H2	-0.20840	0.65930	0.33500	0.0670*
H3	-0.30470	0.81690	0.23690	0.0700*
H6	0.41750	0.58680	0.16560	0.0610*
H7A	-0.30230	0.98770	0.14520	0.0880*
H7B	-0.38990	0.90580	0.08210	0.0880*
H7C	-0.27330	1.00890	0.03310	0.0880*
H8A	0.54070	0.61200	-0.00200	0.0780*
H8B	0.66660	0.68280	0.05710	0.0780*
H8C	0.64580	0.71830	-0.05080	0.0780*
H9	0.47820	0.53310	0.32260	0.0640*
H11	0.72860	0.43920	0.43750	0.0660*
H12	0.85140	0.28650	0.53480	0.0660*
H14	0.34230	0.12890	0.47070	0.0650*
H15	0.20890	0.28590	0.38050	0.0640*
H16A	0.78020	0.16410	0.67210	0.0730*
H16B	0.98690	0.11620	0.59190	0.0730*
H18	1.23550	-0.05380	0.68110	0.0650*
H19A	1.16170	-0.21710	0.88590	0.0790*
H19B	1.34010	-0.23560	0.79000	0.0790*
H20	1.08630	-0.37280	0.75750	0.0740*
H21A	1.13830	-0.40490	0.94730	0.0850*
H21B	1.06940	-0.49510	0.88170	0.0850*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0606 (6)	0.0700 (7)	0.0781 (8)	0.0077 (5)	-0.0144 (6)	0.0221 (6)
O2	0.0562 (6)	0.0660 (6)	0.0546 (7)	0.0023 (5)	-0.0065 (5)	0.0151 (5)
O3	0.0753 (7)	0.0498 (5)	0.0673 (7)	-0.0155 (5)	-0.0325 (6)	0.0124 (5)
N1	0.0600 (7)	0.0532 (6)	0.0553 (8)	-0.0148 (5)	-0.0159 (6)	0.0100 (6)
N2	0.0519 (7)	0.0592 (7)	0.0563 (8)	-0.0087 (5)	-0.0153 (6)	0.0104 (6)
N3	0.0569 (8)	0.0777 (9)	0.0704 (9)	-0.0128 (6)	-0.0112 (7)	0.0244 (7)
N4	0.0511 (7)	0.0728 (8)	0.0710 (9)	-0.0081 (6)	-0.0111 (6)	0.0185 (7)
C1	0.0548 (8)	0.0477 (7)	0.0528 (8)	-0.0149 (6)	-0.0180 (6)	0.0053 (6)
C2	0.0505 (8)	0.0608 (8)	0.0587 (9)	-0.0154 (7)	-0.0097 (7)	0.0033 (7)
C3	0.0459 (7)	0.0590 (8)	0.0685 (10)	-0.0034 (6)	-0.0123 (7)	-0.0003 (7)
C4	0.0514 (8)	0.0505 (7)	0.0571 (9)	-0.0059 (6)	-0.0183 (6)	0.0066 (7)
C5	0.0496 (7)	0.0528 (7)	0.0479 (8)	-0.0092 (6)	-0.0148 (6)	0.0040 (6)

C6	0.0493 (7)	0.0500 (7)	0.0538 (9)	-0.0045 (6)	-0.0158 (6)	0.0039 (6)
C7	0.0631 (10)	0.0586 (9)	0.0952 (14)	0.0050 (7)	-0.0242 (9)	0.0085 (9)
C8	0.0573 (9)	0.0723 (10)	0.0603 (10)	-0.0005 (7)	-0.0056 (7)	0.0072 (8)
C9	0.0585 (8)	0.0535 (8)	0.0524 (9)	-0.0174 (6)	-0.0138 (7)	0.0070 (6)
C10	0.0540 (8)	0.0515 (7)	0.0457 (8)	-0.0107 (6)	-0.0083 (6)	0.0052 (6)
C11	0.0584 (8)	0.0516 (7)	0.0599 (9)	-0.0183 (6)	-0.0156 (7)	0.0094 (7)
C12	0.0539 (8)	0.0539 (8)	0.0611 (9)	-0.0143 (6)	-0.0191 (7)	0.0072 (7)
C13	0.0539 (8)	0.0476 (7)	0.0478 (8)	-0.0083 (6)	-0.0112 (6)	0.0054 (6)
C14	0.0597 (8)	0.0498 (7)	0.0565 (9)	-0.0184 (6)	-0.0154 (7)	0.0062 (7)
C15	0.0536 (8)	0.0573 (8)	0.0529 (9)	-0.0147 (6)	-0.0167 (6)	0.0060 (7)
C16	0.0618 (9)	0.0590 (8)	0.0662 (10)	-0.0142 (7)	-0.0253 (7)	0.0123 (7)
C17	0.0522 (8)	0.0542 (7)	0.0512 (8)	-0.0095 (6)	-0.0124 (6)	0.0046 (6)
C18	0.0496 (8)	0.0582 (8)	0.0554 (9)	-0.0111 (6)	-0.0094 (6)	0.0083 (7)
C19	0.0655 (9)	0.0647 (9)	0.0701 (11)	-0.0091 (7)	-0.0285 (8)	0.0174 (8)
C20	0.0631 (9)	0.0614 (9)	0.0583 (10)	0.0010 (7)	-0.0173 (7)	-0.0014 (8)
C21	0.0779 (11)	0.0622 (9)	0.0745 (12)	-0.0105 (8)	-0.0244 (9)	0.0093 (9)

Geometric parameters (\AA , $^\circ$)

O1—C4	1.3603 (19)	C16—C17	1.478 (2)
O1—C7	1.416 (2)	C17—C18	1.3572 (18)
O2—C5	1.3658 (17)	C19—C20	1.482 (2)
O2—C8	1.415 (2)	C20—C21	1.296 (3)
O3—C13	1.3651 (17)	C2—H2	0.9300
O3—C16	1.426 (2)	C3—H3	0.9300
N1—C1	1.4215 (18)	C6—H6	0.9300
N1—C9	1.265 (2)	C7—H7A	0.9600
N2—N3	1.3392 (17)	C7—H7B	0.9600
N2—C18	1.3313 (19)	C7—H7C	0.9600
N2—C19	1.468 (2)	C8—H8A	0.9600
N3—N4	1.311 (2)	C8—H8B	0.9600
N4—C17	1.3528 (19)	C8—H8C	0.9600
C1—C2	1.3780 (18)	C9—H9	0.9300
C1—C6	1.405 (2)	C11—H11	0.9300
C2—C3	1.384 (2)	C12—H12	0.9300
C3—C4	1.381 (2)	C14—H14	0.9300
C4—C5	1.4008 (18)	C15—H15	0.9300
C5—C6	1.3732 (19)	C16—H16A	0.9700
C9—C10	1.458 (2)	C16—H16B	0.9700
C10—C11	1.3906 (18)	C18—H18	0.9300
C10—C15	1.3956 (19)	C19—H19A	0.9700
C11—C12	1.372 (2)	C19—H19B	0.9700
C12—C13	1.3883 (19)	C20—H20	0.9300
C13—C14	1.3904 (18)	C21—H21A	0.9300
C14—C15	1.370 (2)	C21—H21B	0.9300
C4—O1—C7	119.09 (13)	C4—C3—H3	120.00
C5—O2—C8	118.37 (12)	C1—C6—H6	120.00
C13—O3—C16	116.35 (11)	C5—C6—H6	120.00
C1—N1—C9	118.68 (12)	O1—C7—H7A	109.00

N3—N2—C18	110.57 (12)	O1—C7—H7B	109.00
N3—N2—C19	120.68 (13)	O1—C7—H7C	109.00
C18—N2—C19	128.62 (13)	H7A—C7—H7B	109.00
N2—N3—N4	107.23 (12)	H7A—C7—H7C	109.00
N3—N4—C17	108.59 (12)	H7B—C7—H7C	110.00
N1—C1—C2	119.65 (13)	O2—C8—H8A	109.00
N1—C1—C6	121.52 (11)	O2—C8—H8B	109.00
C2—C1—C6	118.75 (13)	O2—C8—H8C	109.00
C1—C2—C3	120.94 (14)	H8A—C8—H8B	110.00
C2—C3—C4	120.51 (14)	H8A—C8—H8C	109.00
O1—C4—C3	126.10 (13)	H8B—C8—H8C	109.00
O1—C4—C5	114.93 (13)	N1—C9—H9	118.00
C3—C4—C5	118.95 (13)	C10—C9—H9	118.00
O2—C5—C4	114.42 (12)	C10—C11—H11	119.00
O2—C5—C6	125.10 (12)	C12—C11—H11	119.00
C4—C5—C6	120.47 (13)	C11—C12—H12	120.00
C1—C6—C5	120.36 (12)	C13—C12—H12	120.00
N1—C9—C10	124.02 (14)	C13—C14—H14	120.00
C9—C10—C11	119.63 (13)	C15—C14—H14	120.00
C9—C10—C15	122.79 (13)	C10—C15—H15	119.00
C11—C10—C15	117.58 (13)	C14—C15—H15	119.00
C10—C11—C12	121.91 (13)	O3—C16—H16A	110.00
C11—C12—C13	119.56 (12)	O3—C16—H16B	110.00
O3—C13—C12	124.27 (12)	C17—C16—H16A	110.00
O3—C13—C14	116.29 (12)	C17—C16—H16B	110.00
C12—C13—C14	119.44 (13)	H16A—C16—H16B	108.00
C13—C14—C15	120.28 (12)	N2—C18—H18	127.00
C10—C15—C14	121.14 (12)	C17—C18—H18	127.00
O3—C16—C17	109.84 (12)	N2—C19—H19A	109.00
N4—C17—C16	123.06 (13)	N2—C19—H19B	109.00
N4—C17—C18	108.21 (13)	C20—C19—H19A	109.00
C16—C17—C18	128.49 (13)	C20—C19—H19B	109.00
N2—C18—C17	105.40 (12)	H19A—C19—H19B	108.00
N2—C19—C20	112.12 (14)	C19—C20—H20	118.00
C19—C20—C21	123.90 (17)	C21—C20—H20	118.00
C1—C2—H2	120.00	C20—C21—H21A	120.00
C3—C2—H2	120.00	C20—C21—H21B	120.00
C2—C3—H3	120.00	H21A—C21—H21B	120.00
C7—O1—C4—C5	173.80 (13)	C2—C3—C4—C5	-0.9 (2)
C7—O1—C4—C3	-4.8 (2)	O1—C4—C5—C6	-178.56 (12)
C8—O2—C5—C6	2.3 (2)	O1—C4—C5—O2	0.26 (17)
C8—O2—C5—C4	-176.48 (13)	C3—C4—C5—O2	178.94 (13)
C16—O3—C13—C12	12.0 (2)	C3—C4—C5—C6	0.1 (2)
C16—O3—C13—C14	-169.30 (13)	C4—C5—C6—C1	0.3 (2)
C13—O3—C16—C17	176.09 (12)	O2—C5—C6—C1	-178.40 (12)
C1—N1—C9—C10	-176.54 (14)	N1—C9—C10—C15	20.6 (2)
C9—N1—C1—C2	-139.67 (14)	N1—C9—C10—C11	-160.07 (15)
C9—N1—C1—C6	43.7 (2)	C9—C10—C15—C14	-179.39 (14)

C18—N2—N3—N4	−0.57 (17)	C9—C10—C11—C12	177.46 (14)
N3—N2—C19—C20	50.9 (2)	C11—C10—C15—C14	1.3 (2)
C18—N2—C19—C20	−133.53 (16)	C15—C10—C11—C12	−3.2 (2)
C19—N2—N3—N4	175.73 (14)	C10—C11—C12—C13	2.5 (2)
C19—N2—C18—C17	−175.21 (14)	C11—C12—C13—C14	0.2 (2)
N3—N2—C18—C17	0.71 (17)	C11—C12—C13—O3	178.86 (14)
N2—N3—N4—C17	0.19 (17)	C12—C13—C14—C15	−2.1 (2)
N3—N4—C17—C18	0.25 (18)	O3—C13—C14—C15	179.18 (13)
N3—N4—C17—C16	−174.51 (15)	C13—C14—C15—C10	1.3 (2)
C6—C1—C2—C3	−0.9 (2)	O3—C16—C17—N4	−51.4 (2)
C2—C1—C6—C5	0.1 (2)	O3—C16—C17—C18	134.98 (16)
N1—C1—C2—C3	−177.66 (13)	N4—C17—C18—N2	−0.58 (17)
N1—C1—C6—C5	176.80 (12)	C16—C17—C18—N2	173.81 (15)
C1—C2—C3—C4	1.3 (2)	N2—C19—C20—C21	−137.14 (18)
C2—C3—C4—O1	177.60 (14)		

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the C1—C6 benzene ring.

D—H···A	D—H	H···A	D···A	D—H···A
C19—H19A···O2 ⁱ	0.97	2.54	3.385 (2)	146
C16—H16A···Cg2 ⁱⁱ	0.97	2.87	3.6647 (18)	140

Symmetry codes: (i) $x+1, y-1, z+1$; (ii) $-x+1, -y+1, -z+1$.