

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(Z)-4-(2-Hydroxyanilino)pent-3-en-2-oneBenghanem Fatiha,^a Keraghel Saida,^{a*} Chahmana Safia,^a Ourari Ali^a and Brelot Lydia^b

^aLaboratoire d'Electrochimie, d'Ingenierie Moléculaire et de Catalyse Redox, Departement de Génie des Procédés, Faculté de Technologie, Université Ferhat Abbas, Sétif, Algeria, and ^bInstitut de Chimie de Strasbourg, UMR 7177 CNRS-UdS, Service de Radiocristallographie, 1 rue Blaise Pascal, 67008 Strasbourg Cedex, France

Correspondence e-mail: s_marouani20012002@yahoo.fr


Received 8 June 2012; accepted 19 June 2012

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.047; wR factor = 0.126; data-to-parameter ratio = 9.9.

In the title compound, $\text{C}_{11}\text{H}_{13}\text{NO}_2$, the dihedral angle between the planes defined by the 2-hydroxyphenylamino group and the pent-3-en-2-one mean plane [maximum deviations = 0.0275 (19) and 0.054 (2) Å, respectively] is 31.01 (10)°. There are intramolecular bifurcated $\text{N}-\text{H}\cdots(\text{O},\text{O})$ hydrogen bonds involving the amine NH group and the adjacent carbonyl and hydroxy O atoms. In the crystal, molecules are linked *via* $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming chains propagating along [100].

Related literature

For transition metal complexes of Schiff bases, see: Salavati-Niasari (2006); Xiong *et al.* (2007); Basu *et al.* (2010). For the biological activity of Schiff bases, see: Jarrahpour *et al.* (2007); El-Masry *et al.* (2000); Singh & Dash (1988). For the use of Schiff bases as intermediates in many industrial processes, see: Salavati-Niasari & Nezamoddin Mirsattari (2007); Katsuki (1995); Ahamad *et al.* (2010); Da Silva *et al.* (2010); Soltani *et al.* (2010). For the tautomeric properties and conformations of the title compound, see: Kabak *et al.* (1998). For the photoconductivity of the title compound, see: Parekh *et al.* (2007), and for its thermochromic properties, see: Moustakali-Mavridis *et al.* (1978); Hadjoudis *et al.* (1987). For hydrogen bonding and graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{13}\text{NO}_2$
 $M_r = 191.22$
 Orthorhombic, $P2_12_12_1$
 $a = 8.7826$ (4) Å
 $b = 10.3999$ (5) Å
 $c = 11.1827$ (3) Å
 $V = 1021.41$ (7) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 173$ K
 $0.50 \times 0.30 \times 0.20$ mm

Data collection

Nonius KappaCCD diffractometer
 7005 measured reflections
 1359 independent reflections
 1260 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.105$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.126$
 $S = 1.07$
 1359 reflections
 137 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.29$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O1}$	0.96 (3)	2.28 (3)	2.633 (2)	100.9 (18)
$\text{N1}-\text{H1N}\cdots\text{O2}$	0.96 (3)	1.85 (3)	2.641 (2)	139 (2)
$\text{O1}-\text{H1}\cdots\text{O2}^i$	0.93 (3)	1.72 (3)	2.637 (2)	172 (3)

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, -z$.

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997); data reduction: *DENZO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

The authors would like to thank Professor J. P. Gisselbrecht, Laboratoire d'Electrochimie et de Chimie Physique du Corps Solide, Strasbourg University, France, for his valuable contributions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2455).

References

- Ahamad, I., Gupta, C., Prasad, R. & Quraishi, A. (2010). *J. Appl. Electrochem.* **40**, 2171–2183.
 Basu, S., Gupta, G., Das, B. & Rao, K. M. (2010). *J. Organomet. Chem.* **695**, 2098–2104.
 Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Da Silva, A. B. D., Elia, E. & Gunha Ponciano Gomes, J. A. D. (2010). *Corros. Sci.* **52**, 788–793.
 El-Masry, A. H., Fahmy, H. H. & Abdelwahed, S. H. A. (2000). *Molecules*, **5**, 1429–1438.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Hadjoudis, E., Vitterakis, M., Moustakali, I. & Mavridis, I. (1987). *Tetrahedron*, **43**, 1345–1360.
 Jarrahpour, A., Khalilil, D., Clercq, E. D., Salmi, C. & Michel, J. (2007). *Molecules*, **12**, 1720–1730.
 Kabak, M., Elmali, A. & Elerman, Y. (1998). *J. Mol. Struct.* **470**, 295–300.
 Katsuki, T. (1995). *Coord. Chem. Rev.* **140**, 189–214.

- Moustakali-Mavridis, I., Hadjoudis, E. & Mavridis, A. (1978). *Acta Cryst.* **B34**, 3709–3715.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Parekh, B. B., Purohit, D. H., Sagayaraj, P., Joshi, H. S. & Josh, M. J. (2007). *Cryst. Res. Technol.* **42**, 407–415.
- Salavati-Niasari, M. (2006). *Microporous Mesoporous Mater.* **95**, 248–256.
- Salavati-Niasari, M. & Nezamoddin Mirsattari, S. (2007). *J. Mol. Catal. A Chem.* **268**, 50–58.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Singh, W. M. & Dash, B. C. (1988). *Pesticides*, **22**, 33–37.
- Soltani, N., Behpour, M., Ghoreishi, S. M. & Naeini, H. (2010). *Corros. Sci.* **52**, 1351–1361.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Xiong, D., Fu, Z., Zhong, S., Jiang, X. & Yin, D. (2007). *Catal. Lett.* **113**, 155–159.

supplementary materials

Acta Cryst. (2012). E68, o2188–o2189 [doi:10.1107/S1600536812027894]

(Z)-4-(2-Hydroxyanilino)pent-3-en-2-one**Benghanem Fatiha, Keraghel Saida, Chahmana Safia, Ourari Ali and BreLOT Lydia****Comment**

Schiff bases have played an important role in the development of coordination chemistry. They have a great capacity for complexation of transition metal (Salavati-Niasari, 2006; Xiong *et al.*, 2007; Basu *et al.*, 2010). A literature survey revealed that this kind of compound possesses diverse biological activities such as antimicrobial (Jarrahpour *et al.*, 2007; El-Masry *et al.*, 2000) and antifungal (Singh & Dash, 1988). They also serve as intermediates in many industrial processes (Salavati-Niasari & Nezamoddin Mirsattari, 2007; Katsuki, 1995) and are used as corrosion inhibitors (Ahamad *et al.*, 2010; Da Silva *et al.*, 2010; Soltani *et al.*, 2010). In order to expand this field of research the title compound, a novel Schiff base derived from a non-aromatic aldehyde, has been synthesized and its crystal structure is reported herein.

The tautomeric properties and conformations of the title compound were investigated by (Kabak *et al.*, 1998). The enol tautomer form of the crystal of the title compound is established by the present crystal structure analysis. The positive photoconductivity of this compound has been studied by (Parekh *et al.*, 2007). The crystal exhibits positive photoconductivity and poor NLO responses, and it is photochromic not thermochromic in the solid state (Moustakali-Mavridis *et al.*, 1978; Hadjoudis *et al.*, 1987).

The molecular structure of the title molecule is shown in Fig. 1. It was prepared by the condensation of acetylacetone and 2-aminophenol and crystallized in the chiral space group $P2_12_12_1$.

The molecule is not planar and the dihedral angle between the two planes defined by O1,C1-C6,N1 and O2,C7-C11 is equal to 31.01 (10)°. The small value of bond N1—C7 (1.346 (2) Å) in comparison to bond N1—C6 (1.416 (3) Å) results in a significant change in the bond angle C7—N1—C6 of 130.76 (17)°. The difference in the C—N bond distances is assumed to be due to the presence of the carbonyl group located at the C10 position. Shortening of the C7—N1 bond length and the large value of the C7—N1—C6 bond angle leads to the existence of an N1—H1···O2 intramolecular hydrogen bond (Table 1) with an S(6) ring motif (Bernstein *et al.*, 1995). The second intramolecular N1—H1···O1 hydrogen bond gives an S(5) ring motif.

In the crystal, molecules are linked via O—H···O hydrogen bonds to form chains propagating along [100] - see Table 1 and Fig. 2.

Experimental

To a ethanol solution (5 ml) of (0.109 g, 1 mmol) of 2-aminophenol was slowly added a ethanol solution (5 ml) of acetylacetone (0.1 g, 1 mmol). The mixture was refluxed with constant stirring under a nitrogen atmosphere for 5 h. The mixture was removed and allowed to cool to rt and the solvent to evaporate slowly. After 20 days yellow crystals of the title compound were obtained. They were washed with ethanol then with diethyl ether.

Refinement

The OH and NH H atoms were located in a difference Fourier map and freely refined. The C-bound H atoms were included in calculated positions and refined using a riding model: C—H = 0.95 and 0.98 Å for CH and CH₃ H atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$ where $k = 1.5$ for methyl H atoms and $= 1.2$ for other H atoms. In the final cycles of refinement, in the absence of significant anomalous scattering effects, the Friedel pairs were merged and $\Delta f''$ set to zero.

Computing details

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).




Figure 1

A view of the molecular structure of the title molecule, with the atom numbering. Displacement ellipsoids are drawn at the 50% probability level. The intramolecular bifurcated N-H...O/O hydrogen bond are shown as dashed lines (see Table 1 for details).




Figure 2

The crystal packing of the title compound viewed along the *b* axis. The N-H \cdots O and O-H \cdots O hydrogen bonds are shown as dashed lines (see Table 1 for details).

(Z)-4-(2-Hydroxyanilino)pent-3-en-2-one

Crystal data

$C_{11}H_{13}NO_2$

$M_r = 191.22$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 8.7826$ (4) Å

$b = 10.3999$ (5) Å

$c = 11.1827$ (3) Å

$V = 1021.41$ (7) Å³

$Z = 4$

$F(000) = 408$

$D_x = 1.244$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4219 reflections

$\theta = 1.0$ – 27.5°

$\mu = 0.09$ mm⁻¹

$T = 173$ K

Prism, yellow

$0.50 \times 0.30 \times 0.20$ mm

Data collection

Nonius KappaCCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

7005 measured reflections

1359 independent reflections

1260 reflections with $I > 2\sigma(I)$

$R_{int} = 0.105$

$\theta_{max} = 27.5^\circ$, $\theta_{min} = 2.7^\circ$

$h = -10 \rightarrow 11$
 $k = -12 \rightarrow 13$

$l = -14 \rightarrow 12$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.126$
 $S = 1.07$
 1359 reflections
 137 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0792P)^2 + 0.0946P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.29 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell esds are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.58388 (17)	0.70724 (15)	0.03852 (14)	0.0320 (4)
O2	0.95582 (17)	0.68677 (15)	0.15060 (12)	0.0331 (4)
N1	0.69854 (19)	0.57884 (17)	0.22040 (14)	0.0248 (4)
C1	0.4942 (2)	0.61181 (19)	0.08258 (17)	0.0257 (5)
C2	0.3527 (2)	0.5805 (2)	0.03503 (19)	0.0322 (6)
C3	0.2679 (3)	0.4816 (2)	0.0850 (2)	0.0387 (7)
C4	0.3242 (3)	0.4126 (2)	0.1812 (2)	0.0402 (7)
C5	0.4665 (3)	0.4415 (2)	0.2282 (2)	0.0349 (6)
C6	0.5510 (2)	0.54332 (18)	0.18138 (17)	0.0256 (5)
C7	0.7611 (2)	0.57778 (19)	0.33025 (16)	0.0253 (5)
C8	0.6698 (3)	0.5315 (3)	0.43570 (18)	0.0392 (7)
C9	0.9069 (2)	0.6227 (2)	0.34810 (17)	0.0276 (5)
C10	0.9993 (2)	0.6778 (2)	0.25800 (17)	0.0275 (5)
C11	1.1530 (3)	0.7300 (3)	0.2911 (2)	0.0400 (7)
H1	0.542 (3)	0.738 (3)	-0.032 (3)	0.044 (7)*
H1N	0.762 (3)	0.619 (3)	0.162 (2)	0.033 (6)*
H2	0.31400	0.62690	-0.03160	0.0390*
H3	0.17070	0.46110	0.05310	0.0460*
H4	0.26530	0.34520	0.21510	0.0480*
H5	0.50630	0.39190	0.29240	0.0420*
H8A	0.56570	0.56470	0.42990	0.0590*
H8B	0.71680	0.56240	0.50980	0.0590*
H8C	0.66760	0.43730	0.43600	0.0590*

H9	0.94840	0.61620	0.42630	0.0330*
H11A	1.23200	0.68290	0.24730	0.0600*
H11B	1.16920	0.71960	0.37730	0.0600*
H11C	1.15810	0.82140	0.27030	0.0600*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0344 (8)	0.0336 (8)	0.0281 (7)	−0.0051 (7)	−0.0054 (6)	0.0101 (6)
O2	0.0357 (7)	0.0395 (8)	0.0241 (7)	−0.0059 (7)	0.0004 (6)	0.0062 (6)
N1	0.0271 (8)	0.0257 (8)	0.0217 (7)	−0.0021 (7)	0.0009 (6)	0.0040 (6)
C1	0.0274 (9)	0.0234 (9)	0.0263 (9)	0.0023 (8)	0.0021 (8)	0.0009 (7)
C2	0.0305 (10)	0.0324 (10)	0.0336 (10)	0.0031 (10)	−0.0033 (8)	−0.0031 (9)
C3	0.0308 (10)	0.0342 (11)	0.0511 (13)	−0.0023 (10)	−0.0051 (10)	−0.0075 (10)
C4	0.0403 (12)	0.0300 (11)	0.0503 (14)	−0.0086 (11)	−0.0004 (10)	0.0027 (10)
C5	0.0407 (12)	0.0263 (10)	0.0377 (11)	−0.0044 (10)	−0.0006 (9)	0.0057 (8)
C6	0.0271 (9)	0.0236 (9)	0.0261 (9)	0.0004 (8)	−0.0002 (7)	−0.0003 (7)
C7	0.0319 (9)	0.0236 (8)	0.0205 (9)	0.0043 (8)	0.0006 (7)	0.0043 (7)
C8	0.0406 (12)	0.0513 (13)	0.0258 (10)	−0.0059 (12)	0.0040 (9)	0.0093 (10)
C9	0.0324 (10)	0.0281 (10)	0.0223 (8)	0.0029 (9)	−0.0028 (8)	0.0042 (7)
C10	0.0304 (9)	0.0250 (9)	0.0272 (9)	0.0023 (9)	0.0004 (7)	−0.0007 (7)
C11	0.0374 (11)	0.0434 (13)	0.0393 (11)	−0.0088 (12)	−0.0011 (10)	−0.0005 (10)

Geometric parameters (Å, °)

O1—C1	1.359 (2)	C9—C10	1.415 (3)
O2—C10	1.264 (2)	C10—C11	1.501 (3)
O1—H1	0.93 (3)	C2—H2	0.9500
N1—C6	1.416 (2)	C3—H3	0.9500
N1—C7	1.346 (2)	C4—H4	0.9500
N1—H1N	0.96 (3)	C5—H5	0.9500
C1—C2	1.390 (3)	C8—H8A	0.9800
C1—C6	1.406 (3)	C8—H8B	0.9800
C2—C3	1.387 (3)	C8—H8C	0.9800
C3—C4	1.385 (3)	C9—H9	0.9500
C4—C5	1.389 (4)	C11—H11A	0.9800
C5—C6	1.395 (3)	C11—H11B	0.9800
C7—C8	1.505 (3)	C11—H11C	0.9800
C7—C9	1.378 (3)		
C1—O1—H1	109.3 (18)	C3—C2—H2	120.00
C6—N1—C7	130.74 (16)	C2—C3—H3	120.00
C6—N1—H1N	115.9 (15)	C4—C3—H3	120.00
C7—N1—H1N	112.9 (15)	C3—C4—H4	120.00
O1—C1—C2	123.38 (18)	C5—C4—H4	120.00
O1—C1—C6	116.69 (16)	C4—C5—H5	120.00
C2—C1—C6	119.93 (18)	C6—C5—H5	120.00
C1—C2—C3	120.0 (2)	C7—C8—H8A	109.00
C2—C3—C4	120.4 (2)	C7—C8—H8B	109.00
C3—C4—C5	120.2 (2)	C7—C8—H8C	109.00

C4—C5—C6	120.1 (2)	H8A—C8—H8B	109.00
N1—C6—C1	115.76 (16)	H8A—C8—H8C	110.00
N1—C6—C5	124.68 (18)	H8B—C8—H8C	109.00
C1—C6—C5	119.39 (18)	C7—C9—H9	118.00
C8—C7—C9	119.35 (17)	C10—C9—H9	118.00
N1—C7—C8	120.02 (17)	C10—C11—H11A	109.00
N1—C7—C9	120.60 (17)	C10—C11—H11B	109.00
C7—C9—C10	124.56 (17)	C10—C11—H11C	109.00
O2—C10—C9	122.23 (17)	H11A—C11—H11B	109.00
O2—C10—C11	118.64 (18)	H11A—C11—H11C	110.00
C9—C10—C11	119.12 (17)	H11B—C11—H11C	110.00
C1—C2—H2	120.00		
C6—N1—C7—C9	-177.06 (19)	C1—C2—C3—C4	-0.8 (3)
C7—N1—C6—C1	148.6 (2)	C2—C3—C4—C5	-0.3 (3)
C7—N1—C6—C5	-36.2 (3)	C3—C4—C5—C6	2.3 (3)
C6—N1—C7—C8	0.8 (3)	C4—C5—C6—N1	-178.26 (19)
C6—C1—C2—C3	-0.1 (3)	C4—C5—C6—C1	-3.2 (3)
O1—C1—C6—N1	-2.5 (3)	N1—C7—C9—C10	3.0 (3)
O1—C1—C6—C5	-177.92 (18)	C8—C7—C9—C10	-174.9 (2)
O1—C1—C2—C3	179.93 (18)	C7—C9—C10—O2	-2.2 (3)
C2—C1—C6—N1	177.60 (17)	C7—C9—C10—C11	176.5 (2)
C2—C1—C6—C5	2.1 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1N \cdots O1	0.96 (3)	2.28 (3)	2.633 (2)	100.9 (18)
N1—H1N \cdots O2	0.96 (3)	1.85 (3)	2.641 (2)	139 (2)
O1—H1 \cdots O2 ⁱ	0.93 (3)	1.72 (3)	2.637 (2)	172 (3)

Symmetry code: (i) $x-1/2, -y+3/2, -z$.