

(Z)-4-Chloro-N-(1-{2-[3-(4-chlorobenzoyl)ureido]ethyl}imidazolidin-2-ylidene)benzamide

Dalina Adan,^a Suhaila Sapari,^a Siti Nadiah Halim^b and Bohari M. Yamin^{a*}

^aSchool of Chemical Sciences and Food Technology, Universiti Kebangsaan Malaysia, UKM 43500 Bangi Selangor, Malaysia, and ^bDepartment of Chemistry, Universiti Malaya, Kuala Lumpur, 50603 Malaysia.
Correspondence e-mail: bohari@ukm.my


Received 18 June 2012; accepted 21 June 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.040; wR factor = 0.090; data-to-parameter ratio = 14.7.

The title compound, $\text{C}_{20}\text{H}_{19}\text{Cl}_2\text{N}_5\text{O}_2\text{S}$, was obtained from the reaction of 4-chlorobenzoyl isothiocyanate with diethylenetriamine. The imidazolidine ring is slightly twisted with an $\text{N}-\text{C}-\text{C}-\text{N}$ torsion angle of $15.4(4)^\circ$, while the thiourea moiety maintains its *trans-cis* geometry. The molecule is stabilized by intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. The crystal structure features $\text{N}-\text{H}\cdots\text{O}$, $\text{N}-\text{H}\cdots\text{S}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and $\pi-\pi$ interactions between benzene rings with a centroid–centroid distance of $3.607(3)\text{ \AA}$.

Related literature

For the structures of bis(*N*-benzoylthioureas) derived from aliphatic diamines, see: Ding *et al.* (2008). For those derived from cyclohexane diamine, see: Jumal *et al.* (2011). For those derived from aromatic diamines, see: Osman & Yamin (2011); Dong *et al.* (2008).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{19}\text{Cl}_2\text{N}_5\text{O}_2\text{S}$

$M_r = 464.36$

Monoclinic, Cc
 $a = 24.488(7)\text{ \AA}$
 $b = 6.645(2)\text{ \AA}$
 $c = 13.108(4)\text{ \AA}$
 $\beta = 97.16(2)^\circ$
 $V = 2116.2(11)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.43\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.22 \times 0.08 \times 0.07\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.911$, $T_{\max} = 0.970$

8499 measured reflections
4165 independent reflections
2855 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.090$
 $S = 0.93$
4165 reflections
283 parameters
5 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$
Absolute structure: Flack (1983)
1987 Friedel pairs
Flack parameter: 0.06 (6)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2A \cdots O1	0.87 (1)	2.00 (4)	2.637 (4)	130 (4)
N2—H2A \cdots S1 ⁱ	0.87 (1)	2.85 (4)	3.438 (3)	127 (4)
N4—H4A \cdots O2	0.86 (1)	1.87 (2)	2.614 (3)	143 (3)
N5—H5A \cdots O1 ⁱⁱ	0.86 (1)	2.09 (2)	2.898 (4)	157 (5)
C17—H17A \cdots O2 ⁱⁱⁱ	0.93	2.46	3.118 (4)	128

Symmetry codes: (i) $x, -y, z - \frac{1}{2}$; (ii) $x, -y, z + \frac{1}{2}$; (iii) $x, y - 1, z$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*, *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2009).

The authors thank Universiti Kebangsaan Malaysia and the Ministry of Higher Education, Malaysia, for research grant No. UKM-GUP-NBT-08-27-110.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2061).

References

- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ding, Y.-J., Chang, X.-B., Yang, X.-Q. & Dong, W.-K. (2008). *Acta Cryst. E64*, o658.
- Dong, W.-K., Yan, H.-B., Chai, L.-Q., Lv, Z.-W. & Zhao, C.-Y. (2008). *Acta Cryst. E64*, o1097.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Jumal, J., Ibrahim, A. R. & Yamin, B. M. (2011). *Acta Cryst. E67*, o1256.
- Nardelli, M. (1995). *J. Appl. Cryst. 28*, 659.
- Osman, U. M. & Yamin, B. M. (2011). *Acta Cryst. E67*, o2583.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2012). E68, o2226 [doi:10.1107/S1600536812028218]

(Z)-4-Chloro-N-(1-{2-[3-(4-chlorobenzoyl)ureido]ethyl}imidazolidin-2-ylidene)benzamide

Dalina Adan, Suhaila Sapari, Siti Nadiah Halim and Bohari M. Yamin

Comment

Some bis(thiourea) compounds with several bridges such as 3,3'-dibenzoyl-1,1'-(butane-1,4-diyl)-dithiourea (Ding *et al.*, 2008) 1,2-bis(*N'*-benzoylthioureido)cyclohexane (Jumal *et al.*, 2011) and 1-(4-methylbenzoyl)-3-{2-[3-(4-methylbenzoyl)thioureido]phenyl}thiourea (Osman & Yamin, 2011) have been synthesized by the established reaction between carbonoyl isothiocyanate with the diamines. However, the reaction of 4-chlorobenzoyl isothiocyanate with diethylenetriamine(DIEN) did not give the expected bis or tris(thiourea) but instead, the title compound was obtained indicating a cyclization by the middle nitrogen atom of DIEN to the thiono carbon forming the azomethine funtional group (Fig.1). The molecule consists of the substituted 4-chlorobenzoylimidazolidin-2-ylidene and 4-chlorobenzoylthioureido moiety connected by the ethylene bridge. The N2—C8 and C8—N1 bond lengths of 1.321 (6) and 1.344 (6) Å respectively which are slightly shorter than C8—N3 bond length (1.364 (6) Å) indicates some degree of delocalization along the bonds. The five membered ring N2/N3/C8/C9/C10 is slightly twisted about the C9—C10 bond with these atoms having deviations of 0.095 (5) Å and -0.091 Å from the mean plane of the 5 atoms. The thiourea moiety S1/N4/N5/C13 is planar (maximum deviation 0.003 (3) Å) for atom C13 from the mean plane. The *cis-trans* geometry of the thiourea moiety is maintained. The benzene rings of the two hanging arms are nearly co-planar with dihedral angle of 4.88 (19)° between them so forming an intramolecular π–π interaction with a centroid to centroid distance of 3.607 (3)°. There are four intramolecular short contacts, N2—H2A···O1, N4—H4A···O2, in the molecule. In the crystal structure, the molecules are linked by N5—H5A···O1(x,-y,1/2+z), N2···H2A···S1(x,-y,-1/2+z) to form a one dimensional zig-zag chain which runs parallel to the *c* axis, Figure 2. These chains are linked and C17—H17A···O2(x,-1+y,z) hydrogen bond forming a two-dimensional sheet which runs parallel to the *bc* plane.

Experimental


A solution of 4-chlorobenzoylisothiocyanate (1.9162 g, 0.015 mol) in 30 ml acetone was added into a flask containing 20 ml acetone solution of diethylenetriamine (0.543 g, 0.005 mol). The solution mixture was stirred in ice bath for 1 h. The resulting solution was poured onto an ice cubes to yield a yellow precipitate. The precipitated was filtered off, washed with distilled water and left to dried in atmosphere. Good quality crystal were obtained by recrystallization from ethanol.

Refinement


All H atoms attached to C and N atoms were fixed geometrically and treated as riding with C—H= 0.93 Å(aromatic) or 0.97 Å(methylene)and N—H= 0.86 Å with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C or N})$. There were 2174 Friedel Pairs.

Computing details

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT* (Bruker, 2000); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008), *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2009).

**Figure 1**


The molecular structure of (I), with displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

Molecular packing of (I) viewed down b axis. The dashed lines indicate intermolecular hydrogen bonds. Black cross-hatched atoms are S, blue atoms are N and green atoms are Cl.

(Z)-4-Chloro-N-(1-{2-[3-(4-chlorobenzoyl)ureido]ethyl}imidazolidin-2-ylidene)benzamide

Crystal data



$$M_r = 464.36$$

Monoclinic, Cc

Hall symbol: C -2yc

$$a = 24.488 (7) \text{ \AA}$$

$$b = 6.645 (2) \text{ \AA}$$

$$c = 13.108 (4) \text{ \AA}$$

$$\beta = 97.16 (2)^\circ$$

$$V = 2116.2 (11) \text{ \AA}^3$$

$$Z = 4$$

$$F(000) = 960$$

$$D_x = 1.457 \text{ Mg m}^{-3}$$

$$\text{Mo } K\alpha \text{ radiation, } \lambda = 0.71073 \text{ \AA}$$

Cell parameters from 2230 reflections

$$\theta = 3.1\text{--}26.4^\circ$$

$$\mu = 0.43 \text{ mm}^{-1}$$

$$T = 296 \text{ K}$$

Slab, colourless

$$0.22 \times 0.08 \times 0.07 \text{ mm}$$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: 83.66 pixels mm⁻¹
 ω scan
 Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.911$, $T_{\max} = 0.970$

8499 measured reflections
 4165 independent reflections
 2855 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$
 $\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 3.1^\circ$
 $h = -30 \rightarrow 30$
 $k = -8 \rightarrow 8$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.090$
 $S = 0.93$
 4165 reflections
 283 parameters
 5 restraints
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0374P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.18 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack (1983) ???? Friedel pairs
 Flack parameter: 0.06 (6)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.84274 (4)	0.3989 (2)	0.19364 (9)	0.0858 (4)
Cl2	0.83019 (4)	-0.17344 (19)	0.37999 (10)	0.0887 (4)
S1	0.47526 (3)	0.21472 (13)	0.36669 (7)	0.0574 (3)
O1	0.59035 (9)	0.0679 (4)	0.0292 (2)	0.0654 (7)
O2	0.63898 (9)	0.5325 (3)	0.35719 (19)	0.0536 (6)
N1	0.56767 (10)	0.3697 (4)	0.1039 (2)	0.0456 (7)
N2	0.48754 (11)	0.1718 (5)	0.0424 (2)	0.0564 (8)
H2A	0.5083 (15)	0.083 (5)	0.018 (3)	0.100 (16)*
N3	0.47666 (10)	0.4668 (4)	0.1099 (2)	0.0478 (7)
N4	0.53148 (11)	0.5442 (4)	0.3285 (2)	0.0479 (7)
H4A	0.5640 (7)	0.584 (5)	0.319 (3)	0.053 (10)*
N5	0.58290 (10)	0.2711 (4)	0.3890 (2)	0.0426 (6)
H5A	0.583 (2)	0.153 (3)	0.415 (4)	0.13 (2)*
C1	0.70080 (13)	0.1240 (6)	0.1077 (3)	0.0509 (9)
H1B	0.6892	-0.0066	0.0914	0.061*

C2	0.75617 (14)	0.1603 (6)	0.1344 (3)	0.0550 (9)
H2B	0.7817	0.0561	0.1368	0.066*
C3	0.77275 (14)	0.3549 (6)	0.1573 (3)	0.0566 (10)
C4	0.73592 (13)	0.5075 (5)	0.1551 (3)	0.0534 (9)
H4B	0.7481	0.6377	0.1710	0.064*
C5	0.68087 (14)	0.4709 (5)	0.1295 (2)	0.0496 (8)
H5B	0.6559	0.5764	0.1288	0.060*
C6	0.66193 (13)	0.2761 (5)	0.1045 (2)	0.0434 (8)
C7	0.60276 (13)	0.2283 (5)	0.0752 (2)	0.0444 (8)
C8	0.51449 (13)	0.3336 (5)	0.0849 (2)	0.0433 (8)
C9	0.42894 (14)	0.2020 (6)	0.0255 (3)	0.0606 (9)
H9A	0.4095	0.0871	0.0488	0.073*
H9B	0.4164	0.2259	-0.0466	0.073*
C10	0.42110 (14)	0.3873 (6)	0.0898 (3)	0.0624 (10)
H10A	0.3964	0.4832	0.0520	0.075*
H10B	0.4068	0.3519	0.1532	0.075*
C11	0.48875 (14)	0.6591 (5)	0.1608 (3)	0.0538 (9)
H11A	0.4633	0.7594	0.1294	0.065*
H11B	0.5256	0.7006	0.1504	0.065*
C12	0.48476 (14)	0.6519 (5)	0.2745 (3)	0.0516 (9)
H12A	0.4839	0.7878	0.3011	0.062*
H12B	0.4509	0.5851	0.2862	0.062*
C13	0.53131 (12)	0.3544 (5)	0.3597 (2)	0.0420 (7)
C14	0.63297 (12)	0.3540 (5)	0.3762 (2)	0.0412 (7)
C15	0.68047 (12)	0.2137 (5)	0.3839 (2)	0.0419 (7)
C16	0.67515 (12)	0.0123 (5)	0.3578 (2)	0.0443 (8)
H16A	0.6403	-0.0424	0.3402	0.053*
C17	0.72101 (13)	-0.1086 (5)	0.3576 (3)	0.0483 (8)
H17A	0.7173	-0.2439	0.3398	0.058*
C18	0.77199 (13)	-0.0256 (6)	0.3840 (3)	0.0516 (8)
C19	0.77881 (13)	0.1758 (5)	0.4111 (3)	0.0524 (8)
H19A	0.8137	0.2295	0.4294	0.063*
C20	0.73277 (12)	0.2930 (5)	0.4101 (2)	0.0473 (8)
H20A	0.7366	0.4285	0.4274	0.057*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0511 (5)	0.1137 (9)	0.0944 (8)	-0.0131 (6)	0.0161 (5)	-0.0142 (7)
Cl2	0.0558 (6)	0.0981 (9)	0.1106 (9)	0.0265 (6)	0.0045 (5)	-0.0239 (8)
S1	0.0419 (4)	0.0488 (5)	0.0824 (7)	0.0017 (4)	0.0109 (4)	0.0106 (5)
O1	0.0544 (13)	0.0527 (15)	0.0894 (19)	-0.0025 (11)	0.0106 (12)	-0.0293 (14)
O2	0.0536 (13)	0.0330 (13)	0.0734 (17)	-0.0069 (10)	0.0045 (11)	0.0035 (12)
N1	0.0427 (15)	0.0405 (16)	0.0538 (18)	0.0054 (12)	0.0067 (13)	-0.0050 (13)
N2	0.0456 (16)	0.0532 (19)	0.071 (2)	0.0027 (14)	0.0082 (14)	-0.0145 (16)
N3	0.0490 (15)	0.0394 (16)	0.0549 (18)	0.0093 (12)	0.0058 (13)	-0.0024 (14)
N4	0.0479 (16)	0.0333 (15)	0.0626 (19)	0.0049 (12)	0.0068 (14)	0.0006 (13)
N5	0.0381 (14)	0.0378 (15)	0.0518 (17)	0.0011 (12)	0.0055 (12)	0.0054 (14)
C1	0.054 (2)	0.050 (2)	0.049 (2)	0.0073 (16)	0.0064 (16)	-0.0063 (17)
C2	0.053 (2)	0.065 (2)	0.049 (2)	0.0126 (17)	0.0112 (16)	-0.0059 (19)

C3	0.050 (2)	0.074 (3)	0.047 (2)	-0.0066 (19)	0.0136 (17)	-0.004 (2)
C4	0.056 (2)	0.049 (2)	0.058 (2)	-0.0078 (17)	0.0134 (17)	-0.0080 (18)
C5	0.059 (2)	0.046 (2)	0.046 (2)	0.0006 (16)	0.0140 (16)	-0.0008 (16)
C6	0.0538 (19)	0.044 (2)	0.0344 (19)	0.0026 (15)	0.0130 (15)	-0.0008 (15)
C7	0.0492 (18)	0.0414 (19)	0.043 (2)	0.0021 (14)	0.0071 (15)	-0.0056 (16)
C8	0.0509 (19)	0.0421 (19)	0.0378 (18)	0.0025 (15)	0.0090 (15)	-0.0011 (15)
C9	0.0486 (19)	0.062 (2)	0.072 (3)	-0.0022 (16)	0.0121 (18)	-0.008 (2)
C10	0.048 (2)	0.068 (3)	0.070 (3)	0.0153 (17)	0.0064 (18)	-0.003 (2)
C11	0.061 (2)	0.0326 (18)	0.068 (3)	0.0122 (15)	0.0082 (18)	0.0048 (17)
C12	0.063 (2)	0.0361 (19)	0.055 (2)	0.0175 (15)	0.0035 (17)	-0.0065 (16)
C13	0.0421 (16)	0.0401 (19)	0.044 (2)	0.0069 (13)	0.0053 (14)	-0.0033 (15)
C14	0.0422 (17)	0.0419 (19)	0.0393 (19)	-0.0029 (14)	0.0041 (13)	-0.0008 (16)
C15	0.0432 (16)	0.0407 (18)	0.0420 (19)	-0.0046 (13)	0.0060 (14)	0.0014 (15)
C16	0.0411 (16)	0.0443 (19)	0.048 (2)	-0.0046 (14)	0.0074 (14)	0.0008 (16)
C17	0.0473 (18)	0.0401 (18)	0.058 (2)	-0.0011 (15)	0.0101 (16)	-0.0010 (17)
C18	0.0441 (17)	0.060 (2)	0.051 (2)	0.0098 (16)	0.0082 (15)	-0.0001 (18)
C19	0.0387 (17)	0.059 (2)	0.058 (2)	-0.0068 (16)	0.0023 (15)	-0.0031 (19)
C20	0.0448 (18)	0.0454 (19)	0.051 (2)	-0.0051 (15)	0.0033 (15)	-0.0003 (16)

Geometric parameters (\AA , $^\circ$)

C11—C3	1.745 (4)	C4—H4B	0.9300
C12—C18	1.737 (3)	C5—C6	1.400 (4)
S1—C13	1.669 (3)	C5—H5B	0.9300
O1—C7	1.243 (4)	C6—C7	1.486 (4)
O2—C14	1.224 (3)	C9—C10	1.518 (5)
N1—C8	1.317 (4)	C9—H9A	0.9700
N1—C7	1.358 (4)	C9—H9B	0.9700
N2—C8	1.346 (4)	C10—H10A	0.9700
N2—C9	1.438 (4)	C10—H10B	0.9700
N2—H2A	0.865 (10)	C11—C12	1.505 (5)
N3—C8	1.351 (4)	C11—H11A	0.9700
N3—C10	1.453 (4)	C11—H11B	0.9700
N3—C11	1.455 (4)	C12—H12A	0.9700
N4—C13	1.326 (4)	C12—H12B	0.9700
N4—C12	1.456 (4)	C14—C15	1.484 (4)
N4—H4A	0.864 (10)	C15—C16	1.384 (4)
N5—C14	1.374 (4)	C15—C20	1.388 (4)
N5—C13	1.389 (4)	C16—C17	1.381 (4)
N5—H5A	0.857 (10)	C16—H16A	0.9300
C1—C2	1.378 (5)	C17—C18	1.369 (4)
C1—C6	1.385 (4)	C17—H17A	0.9300
C1—H1B	0.9300	C18—C19	1.389 (5)
C2—C3	1.378 (5)	C19—C20	1.369 (4)
C2—H2B	0.9300	C19—H19A	0.9300
C3—C4	1.355 (5)	C20—H20A	0.9300
C4—C5	1.370 (4)		
C8—N1—C7	117.8 (3)	N3—C10—C9	102.4 (3)
C8—N2—C9	112.4 (3)	N3—C10—H10A	111.3

C8—N2—H2A	115 (3)	C9—C10—H10A	111.3
C9—N2—H2A	131 (3)	N3—C10—H10B	111.3
C8—N3—C10	111.9 (3)	C9—C10—H10B	111.3
C8—N3—C11	125.5 (3)	H10A—C10—H10B	109.2
C10—N3—C11	122.3 (3)	N3—C11—C12	113.1 (3)
C13—N4—C12	125.8 (3)	N3—C11—H11A	109.0
C13—N4—H4A	112 (2)	C12—C11—H11A	109.0
C12—N4—H4A	118 (2)	N3—C11—H11B	109.0
C14—N5—C13	127.0 (3)	C12—C11—H11B	109.0
C14—N5—H5A	117 (3)	H11A—C11—H11B	107.8
C13—N5—H5A	116 (3)	N4—C12—C11	110.7 (3)
C2—C1—C6	122.1 (3)	N4—C12—H12A	109.5
C2—C1—H1B	119.0	C11—C12—H12A	109.5
C6—C1—H1B	119.0	N4—C12—H12B	109.5
C3—C2—C1	118.2 (3)	C11—C12—H12B	109.5
C3—C2—H2B	120.9	H12A—C12—H12B	108.1
C1—C2—H2B	120.9	N4—C13—N5	115.3 (3)
C4—C3—C2	121.4 (3)	N4—C13—S1	125.5 (2)
C4—C3—Cl1	120.6 (3)	N5—C13—S1	119.3 (2)
C2—C3—Cl1	118.0 (3)	O2—C14—N5	123.0 (3)
C3—C4—C5	120.3 (3)	O2—C14—C15	120.6 (3)
C3—C4—H4B	119.8	N5—C14—C15	116.3 (3)
C5—C4—H4B	119.8	C16—C15—C20	118.9 (3)
C4—C5—C6	120.6 (3)	C16—C15—C14	122.8 (3)
C4—C5—H5B	119.7	C20—C15—C14	118.0 (3)
C6—C5—H5B	119.7	C17—C16—C15	120.8 (3)
C1—C6—C5	117.4 (3)	C17—C16—H16A	119.6
C1—C6—C7	119.7 (3)	C15—C16—H16A	119.6
C5—C6—C7	122.8 (3)	C18—C17—C16	118.8 (3)
O1—C7—N1	127.1 (3)	C18—C17—H17A	120.6
O1—C7—C6	118.7 (3)	C16—C17—H17A	120.6
N1—C7—C6	114.2 (3)	C17—C18—C19	122.0 (3)
N1—C8—N2	130.2 (3)	C17—C18—Cl2	119.4 (3)
N1—C8—N3	121.8 (3)	C19—C18—Cl2	118.7 (3)
N2—C8—N3	107.9 (3)	C20—C19—C18	118.2 (3)
N2—C9—C10	102.7 (3)	C20—C19—H19A	120.9
N2—C9—H9A	111.2	C18—C19—H19A	120.9
C10—C9—H9A	111.2	C19—C20—C15	121.3 (3)
N2—C9—H9B	111.2	C19—C20—H20A	119.3
C10—C9—H9B	111.2	C15—C20—H20A	119.3
H9A—C9—H9B	109.1		
C6—C1—C2—C3	-0.7 (5)	C11—N3—C10—C9	173.4 (3)
C1—C2—C3—C4	0.7 (5)	N2—C9—C10—N3	15.4 (4)
C1—C2—C3—Cl1	178.7 (3)	C8—N3—C11—C12	-99.4 (4)
C2—C3—C4—C5	-0.1 (5)	C10—N3—C11—C12	73.8 (4)
Cl1—C3—C4—C5	-178.0 (3)	C13—N4—C12—C11	-100.0 (4)
C3—C4—C5—C6	-0.6 (5)	N3—C11—C12—N4	73.6 (4)
C2—C1—C6—C5	0.0 (5)	C12—N4—C13—N5	166.5 (3)

C2—C1—C6—C7	180.0 (3)	C12—N4—C13—S1	−14.0 (5)
C4—C5—C6—C1	0.6 (4)	C14—N5—C13—N4	−9.4 (5)
C4—C5—C6—C7	−179.4 (3)	C14—N5—C13—S1	171.1 (3)
C8—N1—C7—O1	2.4 (5)	C13—N5—C14—O2	16.6 (5)
C8—N1—C7—C6	−176.9 (3)	C13—N5—C14—C15	−161.9 (3)
C1—C6—C7—O1	−18.3 (4)	O2—C14—C15—C16	−149.1 (3)
C5—C6—C7—O1	161.6 (3)	N5—C14—C15—C16	29.5 (5)
C1—C6—C7—N1	161.0 (3)	O2—C14—C15—C20	25.3 (5)
C5—C6—C7—N1	−19.0 (4)	N5—C14—C15—C20	−156.2 (3)
C7—N1—C8—N2	1.9 (5)	C20—C15—C16—C17	−0.1 (5)
C7—N1—C8—N3	−179.3 (3)	C14—C15—C16—C17	174.3 (3)
C9—N2—C8—N1	−173.8 (3)	C15—C16—C17—C18	0.1 (5)
C9—N2—C8—N3	7.3 (4)	C16—C17—C18—C19	0.2 (5)
C10—N3—C8—N1	−175.0 (3)	C16—C17—C18—Cl2	−177.9 (2)
C11—N3—C8—N1	−1.2 (5)	C17—C18—C19—C20	−0.6 (5)
C10—N3—C8—N2	4.0 (4)	Cl2—C18—C19—C20	177.5 (3)
C11—N3—C8—N2	177.8 (3)	C18—C19—C20—C15	0.7 (5)
C8—N2—C9—C10	−14.6 (4)	C16—C15—C20—C19	−0.4 (5)
C8—N3—C10—C9	−12.6 (4)	C14—C15—C20—C19	−174.9 (3)

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N2—H2A···O1	0.87 (1)	2.00 (4)	2.637 (4)	130 (4)
N2—H2A···S1 ⁱ	0.87 (1)	2.85 (4)	3.438 (3)	127 (4)
N4—H4A···O2	0.86 (1)	1.87 (2)	2.614 (3)	143 (3)
N5—H5A···O1 ⁱⁱ	0.86 (1)	2.09 (2)	2.898 (4)	157 (5)
C17—H17A···O2 ⁱⁱⁱ	0.93	2.46	3.118 (4)	128

Symmetry codes: (i) $x, -y, z-1/2$; (ii) $x, -y, z+1/2$; (iii) $x, y-1, z$.